



Tropical Journal of Natural Product Research

Available online at <https://www.tjnpr.org>

Short Communication

Preliminary Assessment of the Association between DArT-SEQ SNP and some Nutritional Traits in African Yam Bean

Olubusayo O. Oluwole^{1,4}, Oluwaseyi E. Olomitutu^{2,4}, Rajneesh Paliwal⁴, Olaniyi A. Oyatomi⁴, Michael T. Abberton⁴, Olawole O. Obembe^{1,3*}¹Department of Biological Sciences, Covenant University, Canaan land, Ota, Ogun state, Nigeria²Department of Agronomy, University of Ibadan, Oyo state, Nigeria³Plant Science Research Cluster, Biotechnology Research Centre, Department of Biological Sciences, Covenant University, Canaan land, Ota, Ogun state, Nigeria⁴Genetic Resources Centre, International Institute of Tropical Agriculture, Ibadan, Nigeria

ARTICLE INFO

Article history:

Received 28 August 2020

Revised 21 October 2020

Accepted 23 November 2020

Published online 30 November 2020

Copyright: © 2020 Oluwole *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

African yam bean (AYB) is an underutilized tropical legume rich in protein and starch content. This study's aim was to use Diversity Arrays Technology (DArT) sequencing to study the genome-wide association studies (GWAS) of nutritional traits. The study was carried out on 137 accessions in IITA, Ibadan. Seeds were harvested at maturity and analysed for protein, oil, and starch using Kjeltex, Soxtec, and Phenol-sulfuric acid assay method. Mixed Linear Model (MLM) was used for GWAS analysis by employing the TASSEL software. AYB protein content ranged from 17.13-27.15%, while the oil and starch content ranged from 0.74-2.27% and 60.58-69.84%, respectively. A total of 3.6K SNP was generated using a high-throughput DArTseq genotype-by-sequencing SNP approach in the AYB population. These generated SNPs were filtered for quality control, and the remaining useful 2.48K SNPs were used for the preliminary GWAS. We identified a promising source of genetic material for important grain nutritional traits that could be used in AYB breeding research. Identification of QTL/gene of protein, oil, and starch will contribute to our understanding of the genetic basis of natural variation in AYB grain nutritional traits.

Keywords: African yam bean, Single Nucleotide Polymorphism, Underutilized legume, Nutrition.

Introduction

The void between the amount of food supply and the growing human population is now a global threat; and this gap can be bridged by exploiting underutilized food sources.^{1,2} Food insecurity and malnutrition are major challenges in Africa which need to be addressed via a multi-sectoral approach.^{3,4} In Africa, malnutrition resulting from habitual consumption of diets low in essential nutrients is a significant challenge. Malnutrition in children and adults is characterized by impaired cognitive mental development, belly protrusions, physical stunted growth, poor pregnancy outcomes, reduced productivity, and increased mortality rates.^{5,6} Several methods of resolving malnutrition are being employed in Africa. Among these are food import, genetic biofortification, supplement distribution, and dietary diversification.⁷ Despite these, there are calls from different quarters for crop improvement strategy to improve the nutritional content of commonly eaten food crops to reduce nutrient deficiencies in sub-Saharan Africa.^{8,9} In achieving this, so much emphasis is placed on the major staple crops, neglecting indigenous underutilized low-cost energy-protein legume and vegetable crops, including African yam bean, *Cucurbita pepo*, *Solanecio*

biafrae, and *Telfairia occidentalis*.¹⁰⁻¹³ African yam bean (AYB) is an underutilized tuberous legume, indigenous to tropical Africa. It can alleviate malnutrition due to its rich nutritional content. The neglect of AYB is considered a disadvantage as it limits access to cheap protein sources by category of Africans under high poverty and malnourished index margin.¹⁴ The seed of AYB is high nutritional content: protein range between 20% - 29%; 50% carbohydrate; 2-5% crude fat content.^{15,16,17} Significant variations have been reported for these nutritional contents,^{18,19,14} suggesting possibilities for improvement. However, quantitatively inherited traits are controlled by several genes, environment, and interaction between these genes and the environment,²⁰ rendering them subject to continual change and making conventional selection difficult. Hence, there is a need for a genome-wide level understanding of the genetic basis of variation for these traits for the development of improved genotype. Due to the availability of high-throughput genomic platforms, genome-wide association study (GWAS) has become a standard tool for identifying genes and loci responsible for the quantitative variation of complex traits.²¹ Once genes and loci are identified and validated, they could be fixed to provide AYB germplasm with durable nutritional content. There is no study on the identification of Quantitative Trait Locus (QTL) for any trait in AYB. This work was proposed as a preliminary study for the identification of putative QTLs controlling protein, starch, and oil contents in AYB.

*Corresponding author E mail:

olawole.obembe@covenantuniversity.edu.ng

Tel: +2348071351953

Citation: Oluwole OO, Olomitutu OE, Paliwal R, Oyatomi OA, Abberton MT, Obembe OO. Preliminary Assessment of the Association between DArT-SEQ SNP and some Nutritional Traits in African Yam Bean. Trop J Nat Prod Res. 2020; 4(11):877-879. doi.org/10.26538/tjnpr/v4i11.5

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria

Materials and Methods

Genotyping of african yam bean germplasms

A total of 137 accessions of AYB were obtained from the Genetic Resources Centre (GRC), International Institute of Tropical Agriculture

ture (IITA), Ibadan, Nigeria. Young leaves of each accession were collected, and genomic DNA was extracted using the CTAB method.²² The quality of the DNA was measured with nanodrop spectrophotometer and agarose gel electrophoresis and its concentration was adjusted to 50-100 ng/mL in volume of 50 mL before sending them for DNA sequencing. All the 137 DNA extracts were sent for the SNP genotyping using genotype-by-sequencing (GBS) method at Diversity Array Technology (DART) Pty Ltd, Canberra, Australia. Genotyping characterization was done using DARTseq high-depth GBS technology (<https://www.diversityarrays.com/technology-and-resources/dartseq/>) and further raw sequenced FASTQ data of each samples were processed using proprietary DART analytical pipeline (Diversity Array Technology, Australia) and generated total over 3600 genome-wide SNPs in 137 African yam bean accessions.

Field trials and assessment of some AYB nutritional contents

All the 137 AYB accessions were planted in IITA, Ibadan. Seeds were bulked at harvest and analysed for the protein, oil, and starch content. Protein, oil, and starch analyses were performed using Kjeltex, Soxtec, and Phenol-sulfuric assay method.^{23,24,25} Correlation analysis was performed using SAS v9.4.

Association mapping

Mixed Linear Model (MLM) was used for the preliminary GWAS analysis using TASSEL software. A $-\log(p\text{-value})$ of 2.03 was used for the significant threshold. Total generated 3.6K SNPs were further filtered using call rate ($\geq 80\%$), average reproducibility ($\geq 95\%$), minor allele frequency (MAF) (≥ 0.01) and SNP missing data 30% and maintained total 2485 quality SNPs for GWAS analysis of the seed oil, protein, and starch content. The association between SNP markers and quality traits was analysed using Mixed Linear Model (MLM) that provide advantage in association analysis over general linear model (GLM) and avoid spurious association by using population structure (Q) and relatedness kinship (K) matrix as covariates in GWAS analysis. TASSEL software²⁶ was used for the preliminary GWAS analysis of quality traits of African yam bean and a $-\log(P\text{ value})$ of > 2.03 was used for the significant threshold to confirm association between SNP and traits.

Results and Discussion

AYB protein content ranged from 17.13-27.15%, while the oil and starch content ranged from 0.74-2.27% and 60.58-69.84%, respectively. The correlation between seed protein and oil was observed to be significantly negative with -0.04953 (Table 1).

A total of 3.6K SNP was generated using a high-throughput DARTseq genotype-by-sequencing (GBS) approach in the AYB population. These generated SNPs were filtered for quality control, and the remaining quality 2,485 SNPs were used for the preliminary GWAS. Out of the 2,485 SNPs used for this study, 405 SNPs aligned with a common bean, the reference genome, while 2,065 SNPs did not align with the common bean reference genome. A total of 50 putative QTLs were observed to be associated with AYB seed protein, oil, and starch content using a significant threshold of $-\log(p\text{-value})$ of 2.03 and P-value (< 0.009). Out of the 50 putative QTLs, 31 QTLs were associated with protein trait, while oil and starch content showed 8 and 11 QTLs, respectively (Figure 1).

The efforts in underutilized crops for genetic linkage mapping and QTL/gene identification of value traits of biotic, abiotic, yield related, agronomical and nutritional traits are key factor for initiating crop improvement program using genomic assisted breeding tools.²⁷ The seed protein, oil, and starch are quantitative traits that are controlled by multiple genes and their interaction with the environment.²¹ We need to identify the genes responsible for each trait to understand how each of these traits is segregated,²⁸ hence the GWAS of AYB. The GWAS for AYB seed protein and oil revealed that some QTLs were found on chromosome 1 that aligned on common bean reference genome, while others were unknown because the reference genome of AYB is not available yet.

Table 1: Pearson Correlation Coefficients

Nutritional Traits	Oil	Protein	Starch
Oil	1		
Protein	-0.04953	1	
	0.4142		
Starch	0.02633	0.17495	1
	0.6643	0.0037	




Figure 1: Genome-wide SNP markers associated with nutritional traits in a set of 137 AYB accessions.

Manhattan plots show GWAS results for protein, oil, and starch. The SNPs are shown on the X-axis and the genome-wide scan $-\log_{10}(P\text{-values})$ are shown on the Y-axis. The horizontal blue lines indicates significant association with traits based on threshold at $-\log_{10}(P\text{ value}) \geq 2.03$.

Conclusion

The identified regions may be of particular interest to geneticists and breeders for the alteration of the seed protein, oil, and starch levels. So far, this is the first and preliminary report on the GWAS on AYB nutritional traits. Research is still on-going to validate the QTLs and determine if the QTLs exhibit genetic stability.

Conflict of interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgements

The authors would like to appreciate and thank very much the Genetic Resources Center, IITA and Covenant University for funding the research and the publication of the article, respectively.

References

- Mohantya CS, Singha V, Chapman MA. Winged bean: An underutilized tropical legume on the path of improvement, to help mitigate food and nutrition security. *Sci Horticult* 2020; 206:1087897.
- Kuyper EM, Engle-Stone R, Arsenault JE, Arimond M, Adams KP, Dewey KG. Dietary gap assessment: an approach for evaluating whether a country's food supply can support healthy diets at the population level. *Public Health Nutr*. 2017; 20: 2277-2288.
- Obayelu AE. A preventable tragedy: how close is Africa to end hunger and malnutrition in all its forms by 2030? *AJFAND*. 2019; 19:1-5.
- Pandey VL, Mahendra DS, Jayachandran U. Impact of agricultural interventions on the nutritional status in South Asia: A review. *Food Policy* 2016; 62:28-40.
- Welch RM and Graham RD. Breeding for micronutrients in staple food crops from a human nutrition perspective. *J Exp Bot*. 2004; 55:353-364.
- Black RE, Allen LH, Bhutta ZA, Caulfield LE, De Onis M, Ezzati M, Mathers C, Rivera J, Maternal and Child Under-nutrition Study Group. Maternal and child undernutrition: global and regional exposures and health consequences. *The Lancet* 2008; 371:243-260.
- Alake CO and Alake OO. Genetic diversity for agro-nutritional traits in African landraces of *Vigna subterranean* Germplasm. *J Crop Improv*. 2016; 30:378-398.
- White PJ and Broadley MR. Biofortifying crops with essential mineral elements. *Trends Plant Sci*. 2005; 10:586-593.
- Zhao FJ, Su YH, Dunham SJ, Rakszegi M, Bedo Z, McGrath SP, Shewry PR. Variation in mineral micronutrient concentrations in grain of wheat lines of diverse origin. *J Cereal Sci*. 2009; 49:290-295.
- Aremu C, Abberton M, Adebisi T, Asaleye AJ, Inegbedion H, Abolusoro S, Adekiya A, Aboyeji C, Dunsin O. Selecting yield and nutritional traits in *Sphenostylis stenocarpa* landraces for food improvement. *Recent Pat Food Nutr Agric*. 2020; 11:69-81.
- Bello OA, Ayanda OI, Aworunse OS, Olukanmi BI, Soladoye MO, Esan EB, Obembe OO. *Solanecio bialfrae*: An underutilized nutraceutically-important African indigenous vegetable. *Pharmacogn Rev*. 2018; 12:128-132.
- Aworunse OS, Bello OA, Popoola JO, Obembe OO. Pharmacotherapeutic properties of *Telfairia occidentalis* Hook F.: A systematic review. *Pharmacogn Rev*. 2018; 12:238-249.
- Obembe OO, Aworunse OS, Bello OA, Ani AO. Multiple Shoots Induction from Indigenous Nigerian Pumpkin (*Cucurbita pepo* L.). *ARRB*. 2017; 17:1-10.
- Aremu CO, Ojuederie OB, Ayo-Vaughan F, Dahunsi O, Adekiya AO, Olayanju A, Adebisi OT, Sunday I, Inegbedion H, Asaleye AJ, Abolusoro S. Morphometric analysis and characterization of the nutritional quality in African yam bean accessions. *Plant Physiol Rep*. 2019; 24:446-459.
- Eromosele CO, Arogundade LA, Eromosele IC, Ademuyiwa O. Extractability of African yam bean (*Sphenostylis stenocarpa*) protein in acid, salt and alkaline aqueous media. *Food Hydrocoll*. 2008; 22:1622-1628.
- Adewale BD, Dumet DJ, Vroh-Bi I, Kehinde OB, Ojo DK, Adegbite AE, Franco J. Morphological diversity analysis of African yam bean (*Sphenostylis stenocarpa* Hochst. ex A. Rich.) Harms and prospects for utilization in germplasm conservation and breeding. *Genet. Resour. Crop Evol*. 2012; 59:927-936.
- Baiyeri SO, Uguru MI, Ogbonna PE, Samuel-Baiyeri CCA, Okechukwu R, Kumaga FK, Amoatey C. Evaluation of the nutritional composition of the seeds of some selected African yam bean (*Sphenostylis stenocarpa* Hochst Ex. A. Rich (Harms)) accessions. *Agro-Sci*. 2018; 17:37-44.
- Ameh IG. Proximate and mineral composition of seed and tuber of African yam bean, *Sphenostylis stenocarpa* (Hochst. ex. a. rich.) Harms. *ASSET: An Int J (Series B)* 2007; 6:1-10.
- Ojuederie OB, Balogun MO, Fawole I, Igwe DO, Olowolafe MO. Assessment of the genetic diversity of African yam bean (*Sphenostylis stenocarpa* Hochst ex. A. Rich. Harms) accessions using amplified fragment length polymorphism (AFLP) markers. *Afr J Biotechnol*. 2014; 13:1850-1858.
- Li D, Zhao X, Han Y, Li W, Xie F. Genome-wide association mapping for seed protein and oil contents using a large panel of soybean accessions. *Genom*. 2018; 111(1):90-95.
- Liu HJ and Yan J. Crop genome-wide association study: a harvest of biological relevance. *Plant J*. 2019; 97:8-18.
- Doyle JJ and Doyle JL. Isolation of plant DNA from fresh tissue. *Focus* 1990; 12:13-15.
- FOSS. Manual for Kjeltac System 8400 Distilling and Titration Unit; FOSS: Hilleroed, Denmark; Slangerupgade, Denmark. 2003.
- Onitilo MO, Sanni L, Daniel I, Maziya-Dixon B, Dixon A. Physicochemical and functional properties of native starches from cassava varieties in southwest Nigeria. *J Food Agric Environ*. 2007; 5:10.
- AOAC. Methods of the association of official analytical chemists. In *Official Methods of Analysis*, 15th ed.; Virginia Association of Official Analytical Chemistry: Rockville, MD, USA, 1990. 1141 p.
- Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. TASSEL: software for association mapping of complex traits in diverse samples. *Bioinform*. 2007; 23:2633-2635.
- Collard BCY, Jahufer MZZ, Brouwer JB, Pang ECK. An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: The Basic Conc Euphy 2005; 142:169-196.
- Yuan Y, Wang X, Wang L, Xing H, Wang Q, Saeed M, Tao J, Feng W, Zhang G, Song X-L, Sun X-Z. Genome-Wide Association Study Identifies Candidate Genes Related to Seed Oil Composition and Protein Content in *Gossypium hirsutum* L. *Front Plant Sci*. 2018; 9:1359.