

Copyright 1995

The CIAT Library would like to thank the publisher of ***Tropical Grasslands*** for giving us permission to include the full text of this article on our website.

Derechos de autor 1995

La Biblioteca del CIAT agradece al editor de ***Tropical Grasslands*** el permiso de incluir el texto completo de este artículo en nuestra pagina web.

Risk of serious anthracnose damage to stylo cultivars

S. CHAKRABORTY¹, D. CAMERON¹,
R. BOLAND¹, S. KELEMU²,
C.D. FERNANDES³ AND
M.J. D'A. CHARCHAR³

¹CSIRO Division of Tropical Crops and Pastures,
Brisbane, Queensland, Australia

²CIAT, Cali, Colombia

³EMBRAPA, Brazil

In Australia, the anthracnose fungus has adapted to attack previously resistant cultivars, resulting in the demise of once productive Townsville, Fitzroy, Endeavour, Graham and Cook stylos, among others. As a result, commercial seed of only 5 cultivars is now produced, although 12 have been released. Seca, resistant to all known races of anthracnose at its release, was affected by a new race within 5 years. These examples reinforce the view that the threat to our stylo-based pastures from new anthracnose races cannot be ignored.

This threat may come from within Australia, where 4 races within each of the 2 biotypes of the fungus may change to damage newly released cultivars. The recent discovery of a new biotype,

which arose through a genetic change in one of the races, is a case in point. We already know that, as cultivars such as Seca are grown over a period of time, frequency of the Seca race increases relative to other races. For example, at a site near Calliope, the frequency of the Seca race increased from 26% to 49% between 1991 and 1994. Whether this will lead to the selection of a more damaging race on Seca, as the area under Seca continues to expand, is not known.

The Australian population of anthracnose represents only a small part of the overall genetic diversity found in its centre of origin in South America. There is a potential risk of damage to our cultivars from races, which may be accidentally introduced from overseas. Research is needed to characterise the diversity that exists in global populations of the fungus, so that the full genetic potential of the fungus to change is well understood. Studying the evolution of the fungus in Australia, and field surveys to detect the appearance of new races, must be a part of ongoing research. As stylo technology relies mainly on a few selected cultivars such as Seca, there is an urgent need to establish the potential risk of serious damage from a new pathogenic race accidentally introduced into Australia and/or evolving locally.

Tropical Grasslands

The Journal of the Tropical Grassland Society of Australia Inc.

Editor

L. Winks

Associate Editors

J.C. Tothill

J.B. Hacker

L.R. Humphreys

R.M. Jones

D.P. Poppi

C.P. Miller

R.E. Hendricksen

R. McLean

K.F. Lowe

A.J. Ash

Editorial Advisory Board

Mr D.G. Cameron

MS 316

Goomboorian Qld 4570

Australia

Prof I. Goto

Kyushu University, 46-06

Fukuoka-shi 812

Japan

Prof L. 't Mannetje

Agricultural University

6709 RZ Wageningen

The Netherlands

Dr M.J. Fisher

CIAT

Apartado aereo 6713

Cali, Colombia

Dr M.M. de Carvalho

EMBRAPA/CNPGL

36155 Coronel Pacheco

MG, Brazil

Prof L.R. Humphreys

University of Queensland

St Lucia Qld 4072

Australia

Mr J.S. Pulsford

3 Forest Hill Drive

Morayfield Qld 4506

Australia

Dr J.R. Wilson

CSIRO

St Lucia Qld 4067

Australia

Dr B.H. Dzowela

SADC-ICRAF

Agroforestry Project

PO Box CY594

Causeway, Harare

Zimbabwe

Tropical Grasslands is an International Journal published quarterly by the Tropical Grassland Society of Australia Inc. and is open to authors from all countries.

The Journal publishes the results of research and development in the evaluation, management and utilisation of pastures and fodder crops in tropical and subtropical agriculture, descriptions of farming systems and review articles.

Authors should follow the style and layout in the current issue of the Journal. Detailed notes for contributors were published in Volume 30 No 1 (March 1996) or are available from the Editor. Authors may seek advice from members of the Editorial Board.

Submit papers directly to Mr L. Winks, Editor, Tropical Grasslands, McNeill's Road, MS 825, Peak Crossing, Qld 4306, Australia.

Subscription: A\$120.00 per annum by sea mail. Send subscription to The Office Manager, Tropical Grassland Society of Australia Inc., c/- CSIRO, St Lucia, Qld 4067, Australia.

The Tropical Grassland Society of Australia Inc. was formed in 1963 to improve the communication of research findings on pasture improvement and management in tropical and subtropical environments.

The activities of the Society include the holding of field meetings, awards to acknowledge innovation, publication through *Tropical Grasslands*, a *Newsletter* and occasional publications and as an agent for relevant books on grassland science and technology.

Membership of the Society is open to anyone interested in pasture improvement. *Tropical Grasslands* is issued free to those with full membership.

Applications for membership should be addressed to The Secretary, Tropical Grassland Society of Australia Inc., c/- CSIRO, St Lucia, Qld 4067, Australia.

Copyright: No part of this Journal may be reproduced or published in any form by print, photoprint, microfilm or any other means without the permission of the Tropical Grassland Society of Australia Inc.