

**MONILIOPHTHORA PERNICIOSA AND M. RORERI PERFORMANCE AT POD LEVEL ON CLONES WITH DIFFERENT DEGREE OF RESISTANCE TO THE DISEASES THEY CAUSE**

Carmen Suárez-Capello<sup>1</sup>; Karina Solís<sup>1</sup>; Sofia Peñaherrera<sup>1</sup>, Patricio del Pozo<sup>2</sup> and Bertus Eskes<sup>3</sup>

<sup>1</sup> Instituto Nacional Autónomo de Investigaciones Agropecuarias, Estación Experimental Tropical, Pichilingue, Departamento Nacional de Protección Vegetal. P. O. Box 24, Quevedo, Provincia de Los Ríos, Ecuador. E-mail: [protveg@hotmail.com](mailto:protveg@hotmail.com);

<sup>2</sup> Plant pathologists, E-mail: [pato70@gmail.com](mailto:pato70@gmail.com),

<sup>3</sup> CFC/ICCO/IPGRI, Parc Scientifique Agropolis II, 34397 Montpellier Cedex 5, France. E-mail : [b.eskes@cgiar.org](mailto:b.eskes@cgiar.org)

**SUMMARY**

Close relationship between the causal agents of Witches' Broom Disease (WBD) and Frosty Pod Rot (FPR) in cocoa have been demonstrated, leading to placed both agents within the same genus, *Moniliophthora* (*perniciosa* and *roreri* respectively) despite conspicuous differences between both species, where the high specialization of the latter to attack only pods is the most stricken character. A consequence of this pod specialization of *M. roreri* is that all tests for resistance has to be performed on adult trees and attempts to develop an early test have failed so far. On the other hand, there is some indirect evidence that clones that show resistance for FPR seems to be as well resistant for WBD on pods. The study herein reported pretends to show this relationship. Clones showing different levels of resistance against WBD and FPR from INIAP's collection at EET Pichilingue were used. Open pollinated pods of about three to four months and ~5cm in size, were inoculated with a cotton ball soaked on a spore suspension of each pathogen. Twenty pods were used for each of the following treatments: only *M. perniciosa* or *M. roreri*; each species in opposite sites at the widest circumference of the pods; each species on the base or the tip of the pod; and a mixed suspension of both organisms together in one single inoculation point; then all treatments were kept cover with a plastic bag for seven days. The study was repeated with two months old hand pollinated pods. In general, *M. roreri* seems to perform better on pods, with the disease progressing 2 to 5 days faster than *M. perniciosa*; each organism was recovered only from correspondingly infected pod tissue. Although there were certain differences in ranges, the latent period for each infection showed similar response for both diseases. A description of symptom development and severity of both diseases were part of this study.

# 16

**INTERNATIONAL COCOA RESEARCH CONFERENCE  
CONFERENCE INTERNATIONALE SUR LA RECHERCHE  
CACAUYERE**

**CONFERÊNCIA INTERNACIONAL DE INVESTIGAÇÃO  
SOBRE O CACAU**

**CONFERENCIA INTERNACIONAL DE INVESTIGACION  
EN CACAO**

---

**BALI - INDONESIA**

**COPAL**



**E**

## **SUMMARIES**

**16 - 21**

**ORGANISED BY THE COCOA PRODUCERS' ALLIANCE (COPAL) IN  
COLLABORATION WITH THE GOVERNMENT OF INDONESIA**

**11**

**ORGANISÉE PAR L'ALLIANCE DES PAYS PRODUCTEURS DE CACAO (COPAL) EN  
COLLABORATION AVEC LE GOUVERNEMENT D'INDONESIE**

**2009**

**ORGANIZADA PELA ALIANÇA DOS PAISES PRODUTORES DE CACAU (COPAL)  
EM COLABORAÇÃO COM O GOVERNO DA INDONÉSIA**

**ORGANIZADA POR LA ALIANZA DE PAISES PRODUCTORES DE CACAO (COPAL)  
EN COLABORACIÓN CON EL GOBIERNO DE INDONESIA**

**16<sup>TH</sup> INTERNATIONAL COCOA RESEARCH CONFERENCE**

**DENPASAR, BALI – INDONESIA**

**From 16 - 21 NOVEMBER 2009**

**ORGANISED BY THE COCOA PRODUCERS' ALLIANCE IN CONJUNCTION WITH  
THE GOVERNEMENT OF INDONESIA**

**THEME**

**"TOWARDS RATIONAL COCOA PRODUCTION AND EFFICIENT USE FOR A  
SUSTAINABLE WORLD COCOA ECONOMY"**

**SUMMARIES**