



**Figure 1.** Conceptual model of agroecological domains

ever, depending upon the geographic location (high-potential production systems or marginal growing environments), the size of each domain will vary. For instance, in marginal environments for rice, the extreme domain will be relatively larger as compared to other domains; whereas, in favorable environments, the middle domains will be relatively larger.

#### *Landraces/varieties distribution across domains*

Until the distribution of landraces/varieties across domains, the features of domains, and the traits of cultivars are analyzed, one cannot appreciate the complexity of farmers' strategies to manage plant genetic resources to meet their multiple needs. From the analysis, it is apparent that one landrace/variety is best suited or most competitive in only one domain, though farmers might grow the same cultivar in more than one domain. This implies that the cultivar competes with other cultivars from within the domain, and that there is less competition between cultivars across domains, except when there is an overlap of cultivars. Overlap signifies the presence of transitional zones between domains, which explains the presence of landraces/varieties in two different but adjacent domains. Within domains, the area and number of households growing different landraces/varieties is explained by market forces, farmers' socioeconomic status, cultural factors, preferences for specific traits, and other abiotic and biotic factors.

Although landraces/varieties may overlap in adjacent domains, no case was registered where a landrace/variety was found in more than two domains. This suggests that landraces/varieties have very specific adaptations. In other words, it reinforces the idea that a cultivar is most competitive in only one domain.

Landraces/varieties falling within the same domain are more likely to be similar in their genetic composition as compared to landraces/varieties from dissimilar domains. The logic behind is that they have been put under similar management conditions have been selected over time for adaptation. However, this hypothesis needs to be proved from laboratory analysis of some of the samples from each domain. If it proves true, then there is a strong case, from a conservation point of view, for disaggregating genetic materials across agroecological domains. Nevertheless, this process still holds true where diversity deployment is the prime objective of the project.

## **Implications of agroecological domains for PPB**

The distribution of landraces/varieties in different domains is the result of farmers' experimentation with those landraces/varieties over years. In other words, they are the "best fit" under farmers' management conditions. Therefore, researchers definitely need to know the characteristics of each domain, as well as the specific traits of the landraces/varieties in each domain and their distribution across domains in order to make any intervention in the present system. The analysis of agroecological domains is worth the money and time invested in collecting and analyzing the information.

### *Planning conservation strategies for landraces*

Identifying landraces that are grown in small areas by a limited number of farmers and devising ways and means of conserving them might seem to be a straightforward task for conserving endangered landraces. Sometimes, weighted diversity, as well, might be computed for facilitating the decision-making process in choosing which landraces to focus on for conservation when there are numerous landraces falling in the endangered category. However, all these processes and steps consider the diversity of landraces at the aggregated/landscape (community) level and thus ignore the influence of agroecological domains in determining the position of landraces in different domains.

The need for micro-level analysis emerges from the fact that landraces are conditioned over years by their continued growth and selection over time in specific domains. As a result, they have developed adaptive traits, which are unique to landraces falling in that domain. Therefore, analysis of landrace diversity at the aggregated level fails to appreciate the position of landraces in specific domains, which in fact might be harboring genes of important traits. Selecting landraces from an aggregated list might exclude certain strategically important landraces from conservation.

PPB has been used as one means to conserve useful genes in landraces through crossing with modern varieties. However, there could be number of landraces within a domain that might require some form of conservation (through breeding and nonbreeding means). Understanding the features of domains and the distribution of landraces in them will facilitate decision making about selecting landraces for conservation. Failing to do this could result in selecting landraces with similar genetic traits for conservation (via PPB) from just one or two domains. This would lead to the neglect of some and overrepresentation of others.

### *Strategies for diversity deployment*

Diversity deployment in simple term means "providing farmers with options of genetic materials to choose from." The introduction of new genetic material results in temporal disequilibrium because of competition between existing and new genetic material. The competition is for space in farmers' fields, for farm labor, for capital inputs, and so on. As time elapses, the new entrant finds its rightful

place in the given environment. This is the outcome of farmers constantly trying to maintain an equilibrium (meeting farmers' objectives) in terms of stabilizing yield and production over time.

The strategy for diversity deployment must begin by analyzing the distribution of landraces/varieties across agroecological domains. Once this is done, researchers would have a clear picture of each domain, along with the distribution of landraces/varieties, and the dominance of certain cultivars against others would become evident. Researchers would also come to know the reasons for this dominance. Only then could they develop their strategy for diversity deployment. In the absence of this information, new genetic materials might fit into domains where there is not much competition. It could also happen that new genetic materials compete with each other landraces/varieties in similar domains, resulting in limited impact of diversity deployment.

### *Justifying PPB*

The conflict between breeding varieties for wide adaptability or for niche environments will perhaps go on. (*Wide adaptability* means the domain for which the suitability of the landrace/variety is large. *Niche environment* means the domain for the given landrace/variety is limited.) In the truest sense, *wide adaptability* should encompass the ability of a cultivar to be grown in several different domains and vice versa for the niche environment. However, such is not the case.

Whatever the case, the proponents of PPB must bear in mind that the approach has to prove its worth in terms of churning out farmer-acceptable varieties efficiently on such a scale that the economic return on investment is positive. But this is possible only when researchers have a clear knowledge of the size and characteristics of the domains the new variety will fit into. In addition, they also need to know the likely existing cultivar to be replaced. Without this information, it would be rather difficult to estimate the potential adoption ceiling of PPB varieties, which implies that the estimation of economic returns at the household level is difficult. This will become an increasingly important issue in the future, when enough time has elapsed between the development and adoption/dissemination of PPB varieties and the evaluation of their impact.

Another important issue that can be addressed by analyzing agroecological domains is orienting PPB programs towards "poverty alleviation" and food security at the household level. Since resource-poor farmers mainly own marginal land, there is limited varietal choice. By conducting PPB programs using landraces from marginal environments, the chances of providing greater options in such environments is increased, which would contribute to food security, particularly in resource-poor households. Targeting PPB for equity of benefits for the resource-poor can also be justified along similar lines.

## **Conclusion**

Agroecological delineation using key informants/farmers from the given community can be reliably done. The identified domains and the associated varieties in each domain have to be verified through a transect walk with the key informants. This exercise helps prioritize landraces/varieties in each domain based on the number of households growing them and the area covered. Using this information, a selection of landraces/varieties for PPB work could be made. Diversity deployment and conservation of certain landraces/varieties could also be planned using this information. The arguments presented here clearly indicate the need to focus PPB initiatives on marginal environments for which there are no MVs, and where, at the same time, the majority of the resource-poor dwell. This exercise has to be conducted prior to initiating PPB work in a given area. Information

required to delineate agroecological domains and associated landraces/varieties can easily be gathered using key informants at the village level. It has been suggested that this exercise be incorporated as a component of PPB work.

## References

- Chaudhary, P. 2000. *Progress updates, Kachorwa, Bara, Nepal (March 1998–December 1999). Strengthening the scientific basis of in situ conservation of agricultural biodiversity on-farm*. A project jointly implemented by NARC/LI-BIRD/IPGRI.
- Collinson, M.P. 1980. *A farming systems contribution to improved relevancy in agricultural research: Concepts and procedures and their promotion by CIMMYT in Eastern Africa*. Nairobi: Centro Internacional de Mejoramiento de Maíz y Trigo, East African Economics Program.
- Cornick, T. and A. Alberti. 1985. Recommendation domain reconsidered. Paper presented at the 1985 Farming Systems Research and Methodology Symposium, Manhattan, Kansas. Manhattan, Kansas: Kansas State University.
- Franzel, S. 1985. Evaluating a method for defining recommendation domains: A case study from Kenya. Paper presented at the 1985 Farming Systems Research and Methodology Symposium, Manhattan, Kansas. Manhattan, Kansas: Kansas State University.
- Harrington, L.W. and R. Tripp. 1985. *Recommendation domains: A framework for on-farm research*. Working Paper 02/84. Mexico, DF: Centro Internacional de Mejoramiento de Maíz y Trigo.
- PRA. 2000. *Participatory Rural Appraisal on Intensive Data Plots Exercise at Begnas, Kaski*. Field Report. Pokhara, Nepal: Local Initiatives for Biodiversity Research and Development.
- Vaidya, A.K. and C.N. Floyd. 1997. *From recommendation domains to providing a basis for research prioritisation and locating representative sites for technology generation and verification in the hills of Nepal*. LARC Occasional Paper No. 97/3. Kaski, Nepal: Lumle Agricultural Research Centre.
- Wotowiec, P. Jr., S. Poats, and P.E. Hildebrand. 1986. Research, recommendation, and diffusion domains: A farming systems approach to targeting. Paper submitted to the 1986 conference on Gender Issues in Farming Systems Research and Extension.

# Listening to Farmers' Perceptions through Participatory Rice Varietal Selection: A Case Study in Villages in Eastern Uttar Pradesh, India

*Thelma R. Paris, Abha Singh, Joyce Luis with Hari Nath Singh, Omkar Nath Singh,  
Sanjay Singh, Ram Kathin Singh, and Surapong Sarkarung*

## Abstract

This paper presents a case study based on the findings in two villages in eastern Uttar Pradesh, India, part of a project started in 1997 to develop, test, and refine methodologies of participatory research and gender analysis as they apply to the development of new technologies in germplasm and natural resource management. The two villages occupy different agroecological areas and also differ in sociocultural characteristics. Both male and female farmers were included in the study, and details of their preferences for the rice varieties studied are presented in this paper.

## Introduction

Decisions about the adoption of technology are conditional to farmers' perceptions of the performance of a new technology relative to that of the technology currently being practiced. Farmers may assess a new technology, such as an improved variety, in terms of a range of attributes, such as grain quality, straw yield, and input requirements, in addition to grain yield (Traxler and Byerlee 1993). In Orissa, eastern India, farmers indicated preference not only for the visual appearance of rice grain, but also for attributes such as cooking quality, taste, keeping quality, and straw quality (Kshirsagar, Pandey, and Bellon 1997). If farmers perceive an improved variety to be inferior to traditional varieties in terms of one or more attributes, they are unlikely to adopt such a variety (Adesina and Zinnah 1993, as cited by Kshirsagar, Pandey, and Bellon 1997). Crop improvement could potentially benefit from farmers' assessments of the relative performance of different varieties under farmer management. Information on the traits desired by farmers and their knowledge of the production system could be invaluable in setting the goals of a breeding program, delineating the target environment, identifying the parents for breeding and defining the management treatment for breeding work (Sperling et al. 1996; Eyzaguirre and Iwanaga 1996).

Varietal preferences may differ, not only between socioeconomic groups but also by gender. In a farmer-participatory breeding (FPB) project on pearl millet in the Jodhpur district, Rajasthan, India, grain yield, early availability of grain, and the ease of harvesting by hand (lower panicle number and lower plant height) were the main considerations for making selections by women. For the men, yield and quality appeared to be a stronger concern (Weltzien, Whitaker, and Anders 1996). While women have traditionally been seed selectors and managers of germplasm in low-input farming systems, scientists have not given enough attention to their local knowledge, criteria for selection, and perceptions regarding new seeds until recently. For instance, the criteria for selecting seeds, practices of animal care and food processing, and the consequent preferences for different kinds of blending various food materials are useful starting points for building on women's

---

Thelma R. Paris and Joyce Luis are with the Social Sciences Division and Surapong Sarkarung is in Plant Breeding, Genetics & Biochemistry at the International Rice Research Institute (IRRI), Los Baños, Philippines. Abha Singh, Hari Nath Singh, Omkar Nath Singh, and Sanjay Singh are with Narendra Deva University and Agricultural Technology (NDUAT), Kumarganj, Faizabad, eastern Uttar Pradesh, India. Ram Kathin Singh is at IRRI's New Delhi Office.

perspectives in participatory research (Gupta et al. 1996). Another example is when high labor demands for manual threshing may create incentives for women to adopt varieties that are easier to thresh (Adesina and Forson 1995). Including women in the early evaluation of varieties ensures that new seeds can be adopted rapidly. Thus, men's and women's criteria and preferences for rice varieties should be well understood and considered in plant-breeding strategies.

In March 1997, a farmer-participatory plant-breeding program for rainfed rice was developed at the International Rice Research Institute (IRRI) in collaboration with the Indian Council of Agricultural Research (ICAR). This project includes six research sites representing different rice ecosystems in eastern India. The project is under the umbrella of the CGIAR's Systemwide Initiative on Participatory Research and Gender Analysis. The goal of this initiative is to develop, test, and refine methodologies of participatory research and gender analysis as they apply to the development of new technologies in germplasm and natural resource management. This FPB project aims to test the hypothesis that farmer participation in rainfed rice breeding can help develop suitable varieties more efficiently. It is also designed to identify the stages in a breeding program where farmer interfacing is optimal. The project has two components: the first is a plant-breeding component, which aims to develop and evaluate a methodology for participatory improvement of rice for heterogeneous environments, and to produce and improve adoption of material suiting farmers' needs. The second is a social-science component (including gender analysis) that aims (1) to characterize cropping systems, diversity of varieties grown, and the crop-management practices of rice farmers, (2) to analyze male and female farmers' selection criteria and their reactions to a range of cultivars and breeding lines, and (3) to enhance the capacities of national agricultural research systems (NARS) in participatory research and gender analysis in plant breeding and rice varietal selection (Courtois et al. 2000). This paper focuses on farmers' selection criteria and their reactions to a range of cultivars and breeding lines under participatory varietal selection conducted on farmers' fields.

## **Characteristics of the villages**

The results of the socioeconomic and gender analysis in the FPB project includes only two villages (table 1): Mungeshpur in the Faizabad district and Basalatpur in the Siddathnagar district, eastern Uttar Pradesh. These sites are among the research sites under the FPB project. A similar study was conducted in the other FPB research sites in Orissa and Madhya Pradesh.

Basalatpur represents favorable (but submergence prone) lowland, rainfed areas. Mungeshpur represents shallow, submergence-prone areas that are favorably rainfed during years of low rainfall. Basalatpur and Mungeshpur have a higher proportion of lowland fields (70% and 60%, respectively) with heavier soil and good water-holding capacity. The flow of natural resources like rainwater (field hydrological conditions) throughout the season has also had a major impact on varietal selection in these villages. Farmers in Mungeshpur have more access to supplementary irrigation, which enables them to diversify into other crops, particularly vegetables and fodder crops. Only one diesel pump exists in Basalatpur and this limits crop diversification. The importance of livestock between the two villages also differs. Livestock in Mungeshpur is more important than in Basalatpur. In Mungeshpur, bullocks continue to be used for land preparation, and threshing is done manually. In contrast, land preparation and threshing in Basalatpur is mechanized with the use of tractors. The degree of market orientation is higher in Basalatpur (nearer the city) where more rice is sold.

**Table 1. Village Characteristics, Basalatur (Siddathnagar District) and Mungeshpur (Faizabad District), India, 1997**

Agroecology	Basalatur, Siddathnagar	Mungeshpur, Faizabad
	Favorable lowland	Shallow, submergence-prone, favorable rainfed during years of low rainfall
Total no. of households	140	133
Sample size for surveys	50	50
No. of male farmers	30	30
No. of female farmers	20	20
<b>Land types (%)</b>		
Lowland	70%	60%
Medium land	0	20%
Upland	30%	20%
Irrigation source (private pump)	1%	10%
Importance of livestock	Low	High
Degree of market orientation	High	Low

The socioeconomic characteristics of the sample households are shown in table 2. Households are classified by official social categories of caste. Muslims dominate in Basalatur (55%), followed by scheduled and backward castes. In Mungeshpur, the backward and scheduled castes dominate (89%). The Yadavs, a subcaste of the backward caste in Mungeshpur, take care of milch animals. The majority of the farming households are owner-cultivators, and share cropping is of limited importance. Female labor participation in rice production is four times higher than that of males in Basalatur and three-fourths in Mungeshpur. There is wide disparity in terms of access to education between men and women. In general, females have lower literacy rates than men. The differences in resource endowments, socioeconomic status, importance of livestock, degree of market orientation, gender roles and responsibilities in rice production, and family size may determine the choice of rice varieties/cultivars and agronomic management practices.

## Cropping systems

Rice followed by wheat + mustard is the predominant cropping pattern in all villages. In Basalatur, wheat and oilseed are grown mainly for domestic use, but rice is grown for consumption as well as marketing. On the other hand, in Mungeshpur, rice is mainly grown for consumption because of low yields and low marketable surplus. Rice is followed by wheat + mustard, which are grown for both domestic consumption and sale. Land preparation for rice is started in June after the arrival of the monsoon. Transplanting and broadcasting are done in July; weeding, in August; and harvesting and threshing, in October to December. During the *rabi* (dry) season from November to April, crops such as wheat + mustard, peas, grams, lentils, *berseem* as green fodder, and vegetables are grown. A few farmers, who have their own irrigation sources, grow crops like mung, maize, vegetables, and green fodder during the *zaid* season (late April to June) in Mungeshpur. Growing crops during the *rabi* and *zaid* seasons is not common in Basalatur because of the lack of irrigation facilities.

**Table 2. Socioeconomic Characteristics of Sample Households, 1997**

Characteristics	Basalatpur, Siddathnagar	Mungeshpur, Faizabad
<b>Caste composition (% of households)</b>		
Upper caste	6%	9%
Backward caste	18%	49%
Scheduled caste	21%	42%
Minority	55%	0
<b>Area by tenure (% of households)</b>		
Share-in	3%	0
Share-out	0	1%
Owner-cultivated	97%	99%
<b>Labor inputs in rice (days/ha)</b>		
Male farmers	25 days/ha (19)	45 days/ha (25)
Female farmers	105 days/ha (81)	130 days/ha (75)
<b>Categories of farmers (%)</b>		
Marginal (<1 ha)	68%	80%
Small (1–2 ha)	24%	16%
Large (>2 ha)	8%	4%
Ave. operational size	1.00 ha	0.70 ha
<b>Literacy rates (%)</b>		
Male head	72%	51%
Female head	40%	14%
Average family size	7	7

*Note:* Figures in parentheses are percentages of total male and female labor inputs in rice production.

## **The gender division of labor in rice production**

The majority of the respondents belong to the lower social class, with small-sized landholdings. Females are younger and have lower literacy rates, compared to males, and have over 20 years of farming experience. The extent of female participation in rice production is high in both villages. Some tasks in rice production and postharvest operations are gender specific. Land preparation and the application of chemicals are men's responsibilities in both villages (10% of fertilizer application is done by women in Basalatpur). In Mungeshpur, women from the lower social status dominate in the work of pulling seedlings (100%), transplanting (70%), weeding (80%), applying farmyard manure (60%), harvesting (82%), and threshing (82%). In Basalatpur, more men than women participate in pulling seedlings and harvesting. Women do the transplanting of seedlings (100%) and most of the weeding (75%), with men doing most of the spraying (90%). Women are also mainly responsible for postharvest activities such as cleaning and selecting the seeds for the next season, storage, and processing rice into other food products for home consumption and for sale. They are the primary end-users of rice byproducts and biomass for livestock and other farm use. A village study in eastern India revealed that women from the lower castes provided 60% to 80% of the total labor input in rice production (Paris et al. 1996). Aside from their significant contributions in rice production, women also provide labor in non-rice crops, collect green animal

fodder, and feed and tend livestock. Thus, men's and women's preferences for specific traits in rice varieties may differ, based on gender-specific roles and responsibilities. With increasing male migration to cities, women are taking on more responsibilities as farm managers, aside from their normal household and childcare responsibilities (Paris et al. 1996).

## Rice varieties

### *Varieties grown by farmers*

The rice varieties currently grown by farmers are shown in table 3. Traditional varieties are more common in Basalatpur than in Mungeshpur. Although modern varieties (MVs) show higher adoption rates in Mungeshpur, these varieties often suffer from submergence, drought, and stress at reproductive and ripening phases when the crop is planted late. Most farmers felt that traditional varieties are more tolerant to drought, submergence, pests, and diseases, while MVs performed well under irrigated conditions. The majority of the farmers indicated that they felt that MVs needed better management than traditional varieties. Modern varieties need more labor, higher levels of fertilizer, and more irrigation, but more farmers prefer to grow MVs because of their higher yields.

**Table 3. Popular Rice Varieties Grown by Farmers According to Land Type**

Land type	Variety	Basalatpur	Mungeshpur
Upland/midland	Traditional	Bengalia, Sarya, Kuwari Mashuri, Oriswa, Malwa	Ari, Bagri, Balbagra, Chaini
	Improved	NDR-97, Sarju-52, PNR-381	Saket-4, NDR-80, 97, 118 NDR-359, Pant-4, Pant-10, Pant-12, Sarju-52
Shallow lowland/lowland	Traditional	Kalamanak, Motibaddam, Malwa, Malasia	Bilaspuri, Indrasan
	Improved	Mashuri, Rajshree, Sambha Mashuri	Mashuri, Madhu, BKP-246, Dwarf Mashuri

### *Topographical adaptations*

Farmers generally match varieties with their environment. For rainfed rice, this means an adaptation to the hydrological conditions of their fields. Each field position in the topo-sequence corresponds to a risk of drought or submergence. The drought risk increases from the bottom to the top of the topo-sequence, while submergence risk decreases along the same path, associated with progressively lower water depths and earlier recession of the water. This translates into different ideotypes for the different situations. Table 4 shows varietal diversity according to land type/topography. In Basalatpur, varieties such as *Bengalia*, *Sarya*, *Oriswa*, *Kuwari Mashuri*, *Malwa*, and *Ghanbhanan* are the major traditional rice cultivars grown in the uplands, and *Kalamanak*, *Malasia*, *Motibaddam*, and *Malwa* are the major varieties grown in the lowlands. Improved varieties, such as NDR-97, PNR-381, and Sarju 52 are grown in the uplands by a few farmers, but the improved variety, Mashuri, occupied more area in the lowlands. In Mungeshpur, the common local varieties grown on upland fields are *Ari*, *Bagri*, 90 days, *Sonia*, *Lalmati*, *Punjab*, *Labbagra*, *Ashwani*, *Indrasan*, and *Bilaspuri*. The improved varieties are Saket-4, NDR-80, and NDR-118 in upland and medium fields and Sarju 52, Mashuri, and dwarf Mashuri mostly in lowland fields.

**Table 4. Farmers' Perceptions of Useful Traits in Selecting Rice Varieties According to Land Type**

Traits	Mungeshpur							
	Upland		Lowland		Upland		Lowland	
	Male	Female	Male	Female	Male	Female	Male	Female
Grain yield	36.67	39.50	48.67	49.67	41.67	35.96	42.06	40.45
Duration	25.83	34.50	0.67	1.00	20.56	25.84	20.56	15.00
Grain price	0.00	0.00	15.67	16.00	1.67	2.81	2.97	1.82
Resistance to abiotic stress	8.33	6.70	0.67	0.33	6.10	6.18	5.10	5.00
Biomass quality	3.33	2.50	5.33	4.67	5.00	2.25	5.52	8.64
Taste	1.67	0.50	10.33	12.33	2.78	2.81	2.12	3.18
Bold and pure grain	7.67	1.50	1.67	0.00	4.44	4.49	3.40	5.00
Adaptation to specific soil type	3.33	3.00	2.33	0.67	5.00	4.49	5.52	6.36
Postharvest quality	0.83	3.00	6.67	7.67	0.00	5.06	0.00	2.27
Resistance to biotic stress	4.17	2.50	1.00	1.33	3.89	1.69	4.25	3.18
Cooking characteristics	0.83	1.00	1.67	2.00	3.89	3.92	3.40	5.00
Response to fertilizer	2.50	1.00	2.67	1.33	5.00	2.25	4.25	1.82
Competitiveness with weeds	0.00	0.00	0.00	2.33	0.00	2.25	0.00	2.27
Resistance to lodging	1.67	0.00	2.65	0.67	0.00	0.00	0.85	0.00
Adaptation to several preparations	2.34	4.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

*Note:* Traits are listed in order of importance. Grain yield includes tillering, panicle length, and number of grains. Resistance to biotic stress includes resistance to pests and blast. Resistance to abiotic stress includes resistance to zinc deficiency and drought. Biomass quality includes height and quality and quantity of straw. Postharvest quality includes ease of hulling and milling recovery. Cooking characteristics include cooking time, elongation ability, aspect after cooking, and impression in the stomach.

Medium-duration fields are grown mostly in medium land. Varieties such as Sarju-52, Ashwani, NDR-359, Pant-4, -10, and -12, and Indrasan are grown on the fields that are located in between upper and lower levels of land type. Farmers of Mungeshpur prefer to grow these varieties on the these land types on the belief that they need optimum moisture during the growth period. Fields differ in agrohydrological characteristics in Basalapur; therefore, some farmers prefer to grow medium varieties on upland fields also.

## Farmers' perceptions of useful traits in varietal adoption

To determine whether there are gender differences in perceptions of useful traits in varietal adoption, we used graphic illustrations of traits. We first showed cards that illustrate useful traits in selecting rice varieties. We then asked each farmer what traits he or she consider in selecting rice varieties for specific land types—upland and lowland fields. To assess how farmers valued each trait, we asked the question, "If you had 100 *paisa*, how much would you pay for each trait? The value in *paisa* allocated to a particular trait corresponded to the importance given by the farmer. Because many traits are interrelated, we reclassified them in consultation with a plant breeder. For example, we grouped traits such as ease in hulling and milling recovery under postharvest quality. Table 2 shows the selection criteria of male and female farmers for different land types and villages.

### ***Favorable rainfed lowlands (Basalatpur, Siddathnagar district)***

In the lowland areas in Basalatpur, yield and duration are the most important traits male and female farmers consider in selection rice varieties.

In this village, the popular traditional varieties are *Bengalia*, *Oriswa*, and *Kuwari mashuri*. These are short-duration (90–110 days), medium-height varieties. The average yields are 2.5 tons per hectare. Farmers prefer short-duration rice varieties in the uplands because of the importance of growing early winter crops such as oilseed, linseed, pulses, peas, and potatoes. They prefer to parboil *Bengalia*; otherwise, its grains break easily. Women in Basalatpur use traditional rice varieties for making puffed rice and *churra*, beaten rice like cornflakes. For women who continue to use the traditional method of hand-pounding rice, postharvest qualities such as ease of hulling and high milling recovery are additional useful traits. The men did not mention these. The finding that women are more concerned than men with postharvest traits and milling recovery are similar to the findings in a participatory breeding project in the high altitudes in Nepal. Sthapit, Joshi, and Witcombe (1996) also observed that women farmers are particularly skillful in assessing postharvest traits, such as milling recovery, and the cooking and eating quality of rice. They found that the evaluation scores between male and female farmers in Chhomrong village showed significant agreement. Women farmers reported that they would like to decide on variety selection after the postharvest evaluation. Consumers preferred white-grained rice to red-pericarped rice because it saves women time in milling.

In Basalatpur, both male and female farmers agreed upon the important traits for lowland rice varieties. Grain price is an important consideration for farmers here because they sell traditional varieties in the market. These, like *Kalamanak*, command a higher price because of their good taste and aroma. *Kalamanak* gives low yields of 1.5 to 2 tons per hectare. In contrast, grain price is not an important consideration in Mungeshpur because rice is mainly used for home consumption and is seldom sold in the market.

### ***Shallow, submergence-prone uplands (Mungeshpur, Faizabad district)***

In Mungeshpur, both male and female farmers agreed upon important traits in selecting varieties for the uplands. Women gave more importance to postharvest qualities and grain quality such as bold and pure grains. For the lowlands, both males and females cited better grain yield, medium duration (125–135 days), biomass, and resistance to abiotic stress as their selection criteria for lowland rice varieties. Women gave greater weight to better adaptation to specific soil types and to grain quality. Women mentioned additional useful traits for varieties in the uplands and lowlands that were not mentioned by men: competitiveness with weeds and postharvest quality. Weeds are the major problem in the uplands, particularly when rice is direct-seeded. In the lowlands, weeds are more prevalent during drought. These additional traits are related to the roles and responsibilities of female family members (e.g., hand weeding and feeding rice straw to livestock).

## **Farmers' evaluation of new rice genotypes grown in farmers' fields**

During the 1999 monsoon season, two farmers from each of the villages of Mungeshpur and Sariyawan (rainfed neighboring village) of the Faizabad district and from Basalatpur were selected to check the performance of rice genotypes in their fields. The genotypes were (1) advanced lines from a shuttle breeding project from Uttar Pradesh, (2) released varieties, and (3) the most common local varieties. Of the 14 genotypes screened in Basalatpur, two are scented varieties (*Kamini*,