

Coverage of Nutrition and Health Interventions in GUJARAT *Insights from the National Family Health Survey-4*


ABOUT THIS DATA NOTE

Nutrition-specific interventions are aimed at improving the proximal food, health, and care environment for women and children during the first 1000 days. They can help improve maternal and child health, improve diets and other nutrition practices, and reduce infections. These interventions span pregnancy, postnatal, and early childhood periods and include food and micronutrient supplementation, nutrition education and/or counselling, growth monitoring and promotion, as well as routine immunization, deworming, and care during illness. At 90% coverage, these interventions can contribute to 20% reduction in stunting and 61% reduction in severe wasting¹.

India's policy framework for health and nutrition is robust and includes most evidence-based nutrition and health interventions. Two large-scale national program platforms – the Integrated Child Development Services and the National Health Mission together provide the public sector delivery platforms with the mandate to deliver these interventions across the country. India's efforts at scaling up nutrition interventions are now also supported by the National Nutrition Mission.

This *Data Note* describes the coverage of a set of key nutrition and health interventions. The findings here are based on data from the National Family Health Survey 2015-2016. Indicators to capture the coverage of the interventions here are all calculated for women (15-49 years) with a child under five years of age. All analyses are based on the last-born child for each woman and indicator definitions are provided in Annex 1 of this Note. For some indicators, age categories may vary.


FIGURE 1 Coverage of interventions across the continuum of care, in 2016



Note: ANC = Antenatal care; ARI = Acute respiratory infection; FP = Family planning; IFA = Iron and folic acid; MCP = Mother and child protection; ORS = Oral rehydration salts; The bars represent state average, the dots represent India average.
Source: NFHS-4.

¹Bhutta, Z.A., J.K. Das, A. Rizvi, M.F. Gaffey, N. Walker, S. Horton, P. Webb, A. Lartey, R.E. Black, Lancet Nutrition Interventions Review G, et al. 2013. "Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost?" *The Lancet* 382(9890):452-477.

MAP 1 Percentage of women who received 4 or more ANC visits, by district, 2016




Top 5 districts, %	
Ahmadabad	95.8
Navsari	91.9
Gandhinagar	88.6
Rajkot	86.8
Bharuch	86.6

Bottom 5 districts, %	
Dohad	39.5
Amreli	41.0
The Dangs	47.5
Bhavnagar	47.5
Kheda	49.9

Source: NFHS-4.

MAP 2 Percentage of women who were weighed during pregnancy, by district, 2016




Top 5 districts, %	
Ahmadabad	96.9
Gandhinagar	94.0
Navsari	93.5
Bharuch	92.7
Rajkot	91.4

Bottom 5 districts, %	
Panchmahal	67.7
Dohad	69.2
Kheda	71.1
Banaskantha	71.2
The Dangs	71.8

Source: NFHS-4.

MAP 3 Percentage of women who received or bought any IFA during pregnancy, by district, 2016




Top 5 districts, %	
Ahmadabad	88.4
Narmada	86.4
Navsari	85.8
The Dangs	85.7
Anand	84.1

Bottom 5 districts, %	
Panchmahal	57.7
Banaskantha	62.8
Surendranagar	65.3
Sabarkantha	68.0
Dohad	68.5

Source: NFHS-4.

MAP 4 Percentage of women who consumed IFA for 100+ days during pregnancy, by district, 2016




Top 5 districts, %	
Navsari	59.7
Ahmadabad	57.4
Bharuch	52.0
Gandhinagar	45.8
Narmada	45.7

Bottom 5 districts, %	
Dohad	14.2
Panchmahal	20.8
Banaskantha	23.6
Patan	24.4
Sabarkantha	28.1

Source: NFHS-4.

MAP 5 Percentage of women who received breastfeeding counselling during pregnancy, by district, 2016




Top 5 districts, %	
Navsari	67.4
The Dangs	59.6
Anand	58.5
Tapi	57.0
Gandhinagar	56.5

Bottom 5 districts, %	
Surat	21.6
Banaskantha	23.9
Kheda	27.6
Rajkot	28.2
Valsad	28.9

Source: NFHS-4.

MAP 6 Percentage of women who received food supplements during pregnancy, by district, 2016




Top 5 districts, %	
The Dangs	85.2
Narmada	83.9
Tapi	80.0
Navsari	77.0
Anand	74.8

Bottom 5 districts, %	
Surat	26.1
Ahmadabad	44.6
Banaskantha	45.3
Kheda	49.2
Dohad	49.8

Source: NFHS-4.

MAP 7 Percentage of women who gave birth in a health facility, by district, 2016



Top 5 districts, %


Jamnagar	97.4
Porbandar	97.1
Mahesana	97.0
Navsari	95.4
Ahmadabad	95.2

Bottom 5 districts, %

The Dangs	57.0
Narmada	76.2
Panchmahal	79.9
Valsad	81.4
Bharuch	82.6

Source: NFHS-4.

MAP 8 Percentage of women who received food supplements during lactation, by district, 2016



Top 5 districts, %


The Dangs	79.6
Tapi	76.3
Narmada	70.5
Mahesana	65.6
Navsari	64.8

Bottom 5 districts, %

Surat	22.6
Ahmadabad	33.1
Banaskantha	41.1
Kheda	41.8
Dohad	41.8

Source: NFHS-4.

MAP 9 Percentage of mothers who received postnatal care within two days of childbirth, by district, 2016




Top 5 districts, %	
Anand	81.8
Navsari	75.5
Gandhinagar	72.9
Surat	69.4
Maheana	69.2

Bottom 5 districts, %	
The Dangs	38.2
Panchmahal	53.6
Banaskantha	54.0
Surendranagar	54.6
Jamnagar	56.2

Source: NFHS-4.

MAP 10 Percentage of children (0-59 months) who received postnatal care within two days of their birth, by district, 2016




Top 5 districts, %	
Gandhinagar	31.5
Anand	25.1
Navsari	24.5
Ahmadabad	23.9
Bharuch	22.0

Bottom 5 districts, %	
Dohad	5.1
Surendranagar	8.2
Sabarkantha	9.2
The Dangs	9.8
Rajkot	10.3

Source: NFHS-4.

MAP 11 Percentage of children (12-23 months) who were fully immunized, by district, 2016




Top 5 districts, %	
Navsari	77.7
Tapi	72.1
Jamnagar	71.9
Narmada	68.3
Porbandar	67.4

Bottom 5 districts, %	
Panchmahal	33.7
Banaskantha	35.5
Patan	36.2
Dohad	36.5
Kheda	38.3

Source: NFHS-4.

MAP 12 Percentage of children (6-35 months) who received food supplements, by district, 2016




Top 5 districts, %	
Tapi	88.2
The Dangs	84.5
Narmada	79.3
Junagarh	78.6
Navsari	78.5

Bottom 5 districts, %	
Ahmadabad	45.0
Surat	46.1
Banaskantha	48.2
Vadodara	55.2
Rajkot	56.7

Source: NFHS-4.

MAP 13 Percentage of children (6-59 months) who received vitamin A supplements, by district, 2016



Top 5 districts, %


Porbandar	91.9
Bhavnagar	90.3
Amreli	87.4
Tapi	83.3
Junagarh	82.7

Bottom 5 districts, %

Banaskantha	58.6
Ahmadabad	60.6
Gandhinagar	62.4
Sabarkantha	63.4
Mahesana	64.2

Source: NFHS-4.

MAP 14 Percentage of children (0-59 months) with diarrhea who received ORS, by district, 2016



Top 5 districts, %


Junagarh	84.8
Jamnagar	66.6
Surendranagar	66.3
Dohad	65.1
Ahmadabad	61.0

Bottom 5 districts, %

Banaskantha	21.0
Amreli	21.0
Panchmahal	21.5
Valsad	24.9
Kachchh	27.1

Source: NFHS-4.

MAP 15 Percentage of children (0-59 months) who were weighed in the last 12 months, by district, 2016




Top 5 districts, %	
Tapi	85.3
The Dangs	80.4
Navsari	77.4
Narmada	75.9
Junagarh	71.8

Bottom 5 districts, %	
Surat	35.0
Kheda	45.1
Banaskantha	46.8
Vadodara	49.0
Ahmadabad	52.0

Source: NFHS-4.

MAP 16 Percentage of mothers with children (0-59 months) who were counselled about child growth after their child was weighed, by district, 2016



Top 5 districts, %	
The Dangs	68.1
Tapi	65.9
Porbandar	61.6
Navsari	59.8
Narmada	57.5

Bottom 5 districts, %	
Surat	25.9
Kheda	26.4
Banaskantha	30.0
Vadodara	30.8
Rajkot	35.1

Source: NFHS-4.

DASHBOARD: Coverage of nutrition-specific interventions, by district, 2016

<25% 25-50% 50-75% ≥75%

State/District		GUJARAT	Ahmadabad	Amreli	Anand	Banaskantha	Bharuch	Bhavnagar	Dohad	Gandhinagar	Jamnagar	Junagarh	Kachchh	Kheda	Mahesana	Narmada	Navsari	Panchmahal	Patan	Porbandar	Rajkot
Pre-pregnancy	Demand for FP satisfied	70.5	61.6	71.2	67.3	67.0	73.2	64.5	65.2	63.7	69.7	76.0	71.8	60.7	71.5	81.1	80.9	59.9	69.0	65.5	66.4
	Iodized salt	94.2	92.8	93.6	84.5	95.9	94.1	86.9	93.1	94.8	97.2	96.2	95.5	96.2	94.6	97.1	92.5	93.9	96.6	97.7	97.7
Pregnancy	Any ANC visits	86.1	97.5	75.9	91.2	79.0	93.8	77.8	77.7	95.6	88.7	90.3	78.2	74.7	91.5	85.4	94.3	75.6	86.1	79.9	94.3
	≥4ANC	70.5	95.8	41.0	78.5	59.0	86.6	47.5	39.5	88.6	80.2	76.2	62.1	49.9	81.2	70.4	91.9	50.9	67.5	56.9	86.8
	Received MCP card	76.0	80.0	88.2	85.1	55.4	89.7	86.9	63.2	85.5	80.7	86.1	75.3	69.0	82.5	88.3	81.9	61.3	77.8	88.1	78.0
	Received IFA tab/syrup	76.7	88.4	80.3	84.1	62.8	79.9	69.1	68.5	77.9	73.8	76.8	74.2	79.8	80.0	86.4	85.8	57.7	81.0	81.1	74.5
	Consumed IFA 100+ days	37.7	57.4	32.1	40.4	23.6	52.0	32.4	14.2	45.8	44.8	28.3	38.3	33.1	34.6	45.7	59.7	20.8	24.4	43.9	42.1
	Deworming	19.7	18.5	15.0	11.2	16.7	10.9	24.4	20.3	13.9	8.9	30.1	14.9	13.3	15.3	28.4	21.6	9.1	27.7	19.3	22.3
	Weighing	83.8	96.9	74.0	90.5	71.2	92.7	77.2	69.2	94.0	88.7	90.3	77.5	71.1	90.6	82.1	93.5	67.7	83.0	79.9	91.4
	Breastfeeding counselling	37.8	35.5	37.1	58.5	23.9	50.7	40.2	42.7	56.5	46.4	43.2	50.2	27.6	40.3	54.0	67.4	36.5	47.1	45.4	28.2
	Food supplementation	55.2	44.6	61.2	74.8	45.3	68.6	57.2	49.8	71.1	54.5	68.5	64.1	49.2	69.0	83.9	77.0	52.1	60.8	69.1	57.9
	Health & nutrition education	47.1	41.1	57.6	65.2	37.4	56.0	49.7	33.9	64.0	51.1	56.2	51.7	36.0	59.3	75.8	66.2	36.1	53.7	63.3	53.3
	Tetanus injection	85.9	93.3	94.3	81.1	77.4	84.1	86.6	74.4	93.3	95.4	87.1	95.0	66.7	88.0	80.3	88.7	68.2	90.0	94.6	91.5
Delivery & postnatal	Institutional birth	90.2	95.2	86.1	94.1	92.0	82.6	93.2	86.8	95.0	97.4	91.2	84.8	89.8	97.0	76.2	95.4	79.9	93.0	97.1	93.9
	Skilled birth attendant	88.5	95.2	86.5	94.7	81.0	83.7	94.1	84.6	96.2	92.5	95.3	85.7	86.7	88.7	74.4	95.8	78.3	94.8	96.7	81.6
	Postnatal care for mothers	63.3	67.4	60.6	81.8	54.0	69.2	57.6	57.6	72.9	56.2	59.6	64.1	58.4	69.2	64.4	75.5	53.6	62.7	66.2	56.3
	Postnatal care for babies	15.7	23.9	14.5	25.1	10.8	22.0	18.5	5.1	31.5	15.9	17.6	17.7	12.3	12.2	18.8	24.5	11.4	17.7	15.1	10.3
	Food supplementation	48.2	33.1	55.0	62.0	41.1	56.1	50.6	41.8	56.6	53.1	61.6	62.1	41.8	65.6	70.5	64.8	48.1	53.0	61.8	52.6
	Health & nutrition education	40.5	30.5	51.1	52.3	35.4	39.1	48.5	29.2	46.6	49.1	46.4	46.9	30.7	56.6	61.6	55.6	33.7	51.1	55.3	49.2
Early childhood	Full immunization	51.0	50.8	59.2	64.7	35.5	57.8	53.9	36.5	64.6	71.9	58.0	44.0	38.3	55.3	68.3	77.7	33.7	36.2	67.4	50.3
	Vitamin A	71.4	60.6	87.4	75.0	58.6	69.0	90.3	72.9	62.4	68.0	82.7	66.9	71.8	64.2	66.1	76.1	66.8	75.3	91.9	71.7
	Paediatric IFA	33.3	29.0	40.0	26.8	28.3	22.1	48.5	37.4	17.5	23.2	46.4	24.5	22.5	31.7	27.2	30.5	33.8	46.5	37.0	32.6
	Deworming	29.8	22.7	27.9	21.9	22.1	11.9	42.7	30.5	17.9	16.6	48.3	25.9	21.5	28.7	23.7	36.7	21.2	44.4	33.7	26.2
	Care seeking for ARI	70.3	63.1	49.8	69	65.2	79.6	70.1	71.1	85.1	83.6	87.4	83.7	79.1	29.0	54.6	59.0	70.5	53.0	79.5	73.1
	ORS during diarrhea	45.6	61.0	21.0	29.0	21.0	33.0	34.0	65.1	41.3	66.6	84.8	27.1	36.7	53.4	44.8	51.0	21.5	51.8	44.3	44.8
	Zinc during diarrhea	19.2	15.5	15.7	12.1	0.0	2.9	22.1	25.5	20.7	45.9	59.4	29.2	12.3	6.3	16.2	5.5	18.6	22.1	0.0	20.6
	Food supplementation	59.9	45.0	61.9	77.1	48.2	63.8	65.2	64.2	69.3	57.3	78.6	72.8	56.8	62.1	79.3	78.5	65.7	66.4	75.1	56.7
	Weighing	56.8	52.0	62.4	70.4	46.8	66.0	59.1	56.2	68.7	65.1	71.8	68.9	45.1	63.4	75.9	77.4	59.4	63.4	71.7	53.6
	Counselling on child growth	40.6	39.5	41.3	52.5	30.0	42.6	48.6	35.8	55.8	53.0	49.5	54.1	26.4	46.2	57.5	59.8	37.2	39.3	61.6	35.1

Source: NFHS-4

DASHBOARD: Coverage of nutrition-specific interventions, by district, 2016

<25% 25-<50% 50-<75% ≥75%

State/District		Sabarkantha	Surat	Surendranagar	Tapi	The Dangs	Vadodara	Valsad	Sabarkantha	Surat	Surendranagar	Tapi	The Dangs	Vadodara	Valsad
Pre-pregnancy	Demand for FP satisfied	69.2	72.1	74.8	82.9	74.1	73.4	69.9	69.2	72.1	74.8	82.9	74.1	73.4	69.9
	Iodized salt	91.2	97.5	93.4	95.3	95.2	95.3	91.0	91.2	97.5	93.4	95.3	95.2	95.3	91.0
Pregnancy	Any ANC visits	81.8	86.3	81.2	87.0	77.9	86.4	78.9	81.8	86.3	81.2	87.0	77.9	86.4	78.9
	≥4ANC	66.4	71.7	52.5	74.3	47.5	71.2	55.0	66.4	71.7	52.5	74.3	47.5	71.2	55.0
	Received MCP card	75.9	68.6	73.2	89.0	83.0	75.1	74.8	75.9	68.6	73.2	89.0	83.0	75.1	74.8
	Received IFA tab/syrup	68.0	81.2	65.3	82.2	85.7	79.2	73.9	68.0	81.2	65.3	82.2	85.7	79.2	73.9
	Consumed IFA 100+ days	28.1	37.7	28.2	43.6	39.6	38.2	41.1	28.1	37.7	28.2	43.6	39.6	38.2	41.1
	Deworming	22.2	29.3	25.0	12.5	9.0	19.3	30.1	22.2	29.3	25.0	12.5	9.0	19.3	30.1
	Weighing	78.6	85.7	78.7	86.5	71.8	83.7	76.9	78.6	85.7	78.7	86.5	71.8	83.7	76.9
	Breastfeeding counselling	40.0	21.6	36.3	57.0	59.6	35.4	28.9	40.0	21.6	36.3	57.0	59.6	35.4	28.9
	Food supplementation	69.9	26.1	60.4	80.0	85.2	56.4	60.8	69.9	26.1	60.4	80.0	85.2	56.4	60.8
	Health & nutrition education	60.3	24.6	49.2	68.8	68.1	44.6	53.9	60.3	24.6	49.2	68.8	68.1	44.6	53.9
	Tetanus injection	88.1	89.4	88.2	91.4	89.5	79.7	78.4	88.1	89.4	88.2	91.4	89.5	79.7	78.4
Delivery & postnatal	Institutional birth	88.4	93.4	82.9	83.0	57.0	83.0	81.4	88.4	93.4	82.9	83.0	57.0	83.0	81.4
	Skilled birth attendant	81.9	95.6	87.9	85.5	57.3	79.5	83.9	81.9	95.6	87.9	85.5	57.3	79.5	83.9
	Postnatal care for mothers	60.5	69.4	54.6	60.1	38.2	66.2	58.9	60.5	69.4	54.6	60.1	38.2	66.2	58.9
	Postnatal care for babies	9.2	12.7	8.2	19.8	9.8	14.8	12.2	9.2	12.7	8.2	19.8	9.8	14.8	12.2
	Food supplementation	60.6	22.6	49.9	76.3	79.6	50.5	53.4	60.6	22.6	49.9	76.3	79.6	50.5	53.4
	Health & nutrition education	53.0	19.1	41.1	64.2	61.6	39.7	47.1	53.0	19.1	41.1	64.2	61.6	39.7	47.1
Early childhood	Full immunization	49.6	44.0	39.8	72.1	44.3	65.2	50.1	49.6	44.0	39.8	72.1	44.3	65.2	50.1
	Vitamin A	63.4	77.4	81.1	83.3	71.7	67.8	81.6	63.4	77.4	81.1	83.3	71.7	67.8	81.6
	Paediatric IFA	34.8	43.0	38.1	35.8	25.9	26.6	43.6	34.8	43.0	38.1	35.8	25.9	26.6	43.6
	Deworming	29.2	43.3	42.1	30.8	20.8	25.8	41.8	29.2	43.3	42.1	30.8	20.8	25.8	41.8
	Care seeking for ARI	56.5	81.1	55.4	61.2	53.1	72.3	79.0	56.5	81.1	55.4	61.2	53.1	72.3	79.0
	ORS during diarrhea	44.4	54.4	66.3	54.1	43.2	41.2	24.9	44.4	54.4	66.3	54.1	43.2	41.2	24.9
	Zinc during diarrhea	13.4	32.0	31.7	33.4	19.1	8.6	8.1	13.4	32.0	31.7	33.4	19.1	8.6	8.1
	Food supplementation	65.1	46.1	61.4	88.2	84.5	55.2	68.1	65.1	46.1	61.4	88.2	84.5	55.2	68.1
	Weighing	61.5	35.0	62.1	85.3	80.4	49.0	62.7	61.5	35.0	62.1	85.3	80.4	49.0	62.7
	Counselling on child growth	42.7	25.9	37.4	65.9	68.1	30.8	51.5	42.7	25.9	37.4	65.9	68.1	30.8	51.5

Source: NFHS-4

Summary of findings

- **High coverage (≥75%):** Sustained efforts are required to maintain the high coverage for the following interventions:
 - *Before & during pregnancy:* iodized salt, any ANC visits, MCP card, IFA provision, weighing, tetanus toxoid injection.
 - *Delivery & postnatal:* institutional birth, skilled birth attendant.
 - *Early childhood:* no interventions achieved high coverage.
- **Low coverage (<50%):** The state should focus efforts on increasing coverage of the following interventions:
 - *Before & during pregnancy:* consumption of 100+ IFA, deworming, breastfeeding counselling, health & nutrition education.
 - *Delivery & postnatal:* postnatal care for babies, food supplementation, health & nutrition education.
 - *Early childhood:* IFA supplementation, deworming, ORS and zinc during diarrhea, counselling on child growth.
- **Missed opportunities:** Coverage of postnatal care is higher for mothers than for babies. Many women receive some IFA but fewer are consuming IFA for at least 100 days during pregnancy, and few women receive deworming pills. Weight monitoring during pregnancy is high but during early childhood is low.

ANNEX 1 Definition of indicators used in the analyses

Indicators	Definition
Pre-pregnancy	
Demand for FP satisfied	Percentage of women (15-49 years) who have their need for family planning satisfied with modern methods.
Iodized salt	Percentage of households with children under 5 years of age using iodized salt.
Pregnancy	
Any ANC visits	Percentage of women (15-49 years) with children under 5 years of age who were attended by any trained provider ever or at least once, when they were pregnant with their youngest child.
≥ 4ANC	Percentage of women (15-49 years) with children under 5 years of age who were attended by any trained provider 4 or more times, when they were pregnant with their youngest child.
Received MCP card	Percentage of women (15-49 years) with children under 5 years of age who received the Mother and Child Protection card after pregnancy registration, when they were pregnant with their youngest child.
Received IFA	Percentage of women (15-49 years) with children under 5 years of age who received or bought any IFA tablets/syrup, when they were pregnant with their youngest child.
Consumed IFA for 100+ days	Percentage of women (15-49 years) with children under 5 years of age who consumed IFA tablets/syrup for 100 days or more, when they were pregnant with their youngest child.
Deworming	Percentage of women (15-49 years) with children under 5 years of age who received any deworming drug, when they were pregnant with their youngest child.
Weighing	Percentage of women (15-49 years) with children under 5 years of age who were weighed when they were pregnant with their youngest child.
Breastfeeding counselling	Percentage of women (15-49 years) with children under 5 years of age who received advice on breastfeeding from any provider, when they were pregnant with their youngest child.
Food supplementation	Percentage of women (15-49 years) with children under 5 years of age who received food supplements from the <i>anganwadi center</i> (AWC), when they were pregnant with their youngest child.
Health & nutrition education	Percentage of women (15-49 years) with children under 5 years of age who received health and nutrition education from the AWC, when they were pregnant with their youngest child.
Tetanus injection	Percentage of women (15-49 years) with children under 5 years of age who received at least 2 TT injections when they were pregnant with their youngest child.
Delivery and postnatal care	
Institutional birth	Percentage of women (15-49 years) with children under 5 years of age who delivered their youngest child in a health facility.
Skilled birth attendant	Percentage of women (15-49 years) with children under 5 years of age who were attended by skilled health personnel when they delivered their youngest child.
Postnatal care for mothers	Percentage of women (15-49 years) with children under 5 years of age who received postnatal care, while in facility or at home, two days after they delivered their youngest child.
Postnatal care for babies	Percentage of last-born children (0-59 months) in the last 5 years, who received postnatal care within two days after they were born.
Food supplementation	Percentage of women (15-49 years) with children under 5 years of age who received food supplements from the AWC, when they were breastfeeding their youngest child.
Health & nutrition education	Percentage of women (15-49 years) with children under 5 years of age who received health and nutrition education from the AWC, when they were breastfeeding their youngest child.
Early childhood	
Full immunization	Percentage of last-born children (12-23 months) in the last 5 years who received one dose of BCG vaccine, three doses of polio vaccine, three doses of DTP3 vaccine, and one dose of measles vaccine.
Vitamin A	Percentage of last-born children (6-59 months) in the last 5 years, who received vitamin A supplements in the six months preceding the survey.
Pediatric IFA	Percentage of last-born children (6-59 months) in the last 5 years, who received iron supplements in the last 7 days prior to the survey.
Deworming	Percentage of last-born children (12-59 months) in the last 5 years, who received albendazole or any other deworming drug in the last 6 months prior to the survey.
Care seeking for ARI	Percentage of last-born children (0-59 months) in the last 5 years, with suspected ARI symptoms in the last two weeks, who were taken to a health care provider.
ORS during diarrhea	Percentage of last-born children (0-59 months) in the last 5 years, with diarrhea in the last two weeks, who received ORS.
Zinc during diarrhea	Percentage of last-born children (2-59 months) in the last 5 years, with diarrhea in the last two weeks who received zinc.
Food supplementation	Percentage of last-born children (6-35 months) in the last 5 years, who received food supplements from the AWC in the last 12 months prior to the survey.
Weighing	Percentage of last-born children (0-59 months) in the last 5 years, who were ever weighed in the last 12 months.
Counselling on child growth	Percentage of women (15-49 years) with children under 5 years of age, who were counselled about their youngest child's growth after they were weighed in the last 12 months prior to the survey.

Led by IFPRI 

AUTHORS

Priyanjana Pramanik, Research Analyst, IFPRI

Rasmi Avula, Research Fellow, IFPRI

Phuong Hong Nguyen, Research Fellow, IFPRI

Samuel Scott, Associate Research Fellow, IFPRI

Purnima Menon, Senior Research Fellow, IFPRI

SUGGESTED CITATION

Pramanik, P., R. Avula, P.H. Nguyen, S. Scott, and P. Menon. 2018. *Coverage of Nutrition and Health Interventions in Gujarat: Insights from the National Family Health Survey-4*. POSHAN Data Note 12. New Delhi, India: International Food Policy Research Institute.

ACKNOWLEDGEMENTS

Lan Mai Tran, Independent Researcher, for support with district-level data extraction from National Family Health Survey factsheets.

ABOUT POSHAN

Partnerships and Opportunities to Strengthen and Harmonize Actions for Nutrition in India (POSHAN) is a multi-year initiative that aims to support the use of data and evidence in decision-making for nutrition in India. It is supported by the Bill & Melinda Gates Foundation and led by IFPRI in India.

ABOUT DATA NOTES

POSHAN Data Notes focus on data visualization to highlight geographic and/or thematic issues related to nutrition in India. They draw on multiple sources of publically available data.

CONTACT US

Email: IFPRI.POSHAN@cgiar.org

IFPRI-NEW DELHI INTERNATIONAL FOOD POLICY RESEARCH INSTITUTE

NASC Complex, CG Block,
Dev Prakash Shastri Road,
Pusa, New Delhi 110012, India
T+91.11.66166565
F+91.11.66781699

IFPRI-HEADQUARTERS INTERNATIONAL FOOD POLICY RESEARCH INSTITUTE

1201 Eye Street, NW,
Washington, DC 20005 USA
T. +1.202.862.5600
F. +1.202.467.4439
Skype: IFPRIhomeoffice
ifpri@cgiar.org
www.ifpri.org

This publication has been prepared by POSHAN. It has not been peer reviewed. Any opinions stated herein are those of the author(s) and do not necessarily reflect the policies of the International Food Policy Research Institute.

Copyright © 2018 International Food Policy Research Institute. All rights reserved. For permission to republish, contact ifpri-copyright@cgiar.org.