

State Nutrition Profile: West Bengal

ABOUT THIS DATA NOTE

This *Data Note* describes the trends for a set of key nutrition and health outcomes, determinants, and coverage of interventions. The findings are based on data from multiple rounds of the National Family Health Surveys using estimates from reports, factsheets, or unit level data. In addition to standard prevalence-based analyses, this *Data Note* includes headcount-based analyses aligned to the POSHAN Abhiyaan monitoring framework to provide evidence that helps identify priority districts and the number of districts where public health concerns exist as per the WHO guidelines¹. This *Data Note* also includes a color-coded dashboard to compare the coverage of nutrition interventions by district. It concludes with key takeaways for children, women, and men, and identifies areas for improvement.





Figure 1. Trends in undernutrition outcomes 2005-2006, 2015-2016, 2019-2021




Source: NFHS-3 (2005-2006) national report and data [IFPRI estimates], NFHS-4 (2015-2016) national report, and NFHS-5 (2019-2021) national and state factsheets. Anemia among non-pregnant and pregnant women for 2005-2006 are IFPRI estimates using woman dataset. ¹WHO. Nutrition Landscape Information System (NLIS). Help Topic: Malnutrition in children. Stunting, wasting, overweight and underweight. (<https://apps.who.int/nutrition/landscape/help.aspx?menu=0&helpid=391&lang=EN>).

²In NFHS-3, NFHS-4, and NFHS-5, 49.3%, 13.9%, and 3.7% of data were missing, respectively.

Map 1 & 2. Number of stunted & anemic children <5y, 2019-2021

Map 1. Stunting




Note: Number in '000s in the above figure

Highest burden districts

1	Murshidabad	267,553
2	South 24 Parganas	250,741
3	North 24 Parganas	205,594
4	Barddhaman	199,000
5	Maldah	160,360

No. of districts with public health concern²: 20 of 20

Map 2. Anemia



Note: Number in '000s in the above figure


Highest burden districts

1	Murshidabad	435,397
2	South 24 Parganas	432,072
3	Barddhaman	360,982
4	North 24 Parganas	330,040
5	Pashchim Medinipur	267,639

No. of districts with public health concern²: 20 of 20

Map 3 & 4. Number of wasted children <5y, 2019-2021

Map 3. Wasting




Note: Number in '000s in the above figure

Highest burden districts

1	South 24 Parganas	144,842
2	Pashchim Medinipur	136,781
3	Barddhaman	128,086
4	Murshidabad	109,576
5	North 24 Parganas	84,395

No. of districts with public health concern²: 20 of 20

Map 4. Severe Wasting



Note: Number in '000s in the above figure

Highest burden districts

1	South 24 Parganas	80,620
2	Pashchim Medinipur	43,336
3	Kolkata	37,194
4	Barddhaman	36,832
5	Haora	36,488

No. of districts with public health concern²: 20 of 20


Source: IFPRI estimates - The headcount was calculated as the product of the undernutrition prevalence, and the total eligible projected population for each district in 2019. Prevalence estimates were obtained from NFHS-5 (2019-2021) district factsheet, and the projected population for 2019 was estimated using Census 2011. Note: The newly formed districts, for which no spatial boundaries were available, were not depicted on the maps.

¹The total number of children <5 years is 6,977,232.

²Public health concern is defined as $\geq 20\%$ for stunting, $\geq 40\%$ for anemia, $\geq 10\%$ for wasting, and $\geq 2\%$ for severe wasting (WHO 2011).

Map 5 & 6. Number of underweight children (<5y) & women (15-49y), 2019-2021

Map 5. Underweight children




Note: Number in '000s in the above figure

Highest burden districts

1	South 24 Parganas	219,996
2	Murshidabad	217,807
3	Bardhaman	199,550
4	Pashchim Medinipur	180,568
5	North 24 Parganas	149,753

No. of districts with public health concern²: 20 of 20

Map 6. Underweight women



Note: Number in '000s in the above figure


Highest burden districts

1	Bardhaman	426,647
2	Murshidabad	415,279
3	Pashchim Medinipur	350,745
4	Bankura	304,791
5	Puruliya	283,857

No. of districts with public health concern²: 16 of 20

Map 7 & 8. Number of anemic women (15-49y), 2019-2021

Map 7. Anemia among non-pregnant women




Note: Number in '000s in the above figure

Highest burden districts

1	North 24 Parganas	2,106,259
2	Bardhaman	1,781,590
3	Murshidabad	1,627,725
4	South 24 Parganas	1,571,494
5	Pashchim Medinipur	1,521,752

No. of districts with public health concern²: 20 of 20

Map 8. Anemia among pregnant women



Note: Number in '000s in the above figure

Highest burden districts

1	Murshidabad	114,422
2	North 24 Parganas	93,405
3	South 24 Parganas	73,385
4	Maldah	70,157
5	Bardhaman	69,254

No. of districts with public health concern²: 16 of 16

Source: IFPRI estimates - The headcount was calculated as the product of the undernutrition prevalence, and the total eligible projected population for each district in 2019. Prevalence estimates were obtained from NFHS-5 (2019-2021) district factsheet, and the projected population for 2019 was estimated using Census 2011. Note: The newly formed districts, for which no spatial boundaries were available, were not depicted on the maps. ¹The total number of children <5 years is 6,977,232, pregnant women 15-49 years is 1,524,627, and non-pregnant women 15-49 years is 26,751,293. ²Public health concern is defined as $\geq 20\%$ for underweight (children), $\geq 10\%$ for underweight (women), $\geq 40\%$ for anemia among non-pregnant women, and $\geq 40\%$ for anemia among pregnant women (WHO 2011).

Figure 2. Trends in overweight/obesity & NCDs¹
2005-2006, 2015-2016, 2019-2021




Table 1. Overweight/obesity & NCDs¹ at district-level
2015-2016, 2019-2021

Category	Outcomes	Worst performing districts (pp)	Best performing districts (pp)	Highest burden districts (thousands) ⁴	No of districts with public health concern ⁵ (total=20)
		<i>Difference between (2019-2021) & (2015-2016)³</i>	<i>Difference between (2019-2021) & (2015-2016)³</i>	2019-2021	2019-2021
Children <5 years	Overweight/obesity	S 24 Parg ⁶ : +6.4 Utt Dinaj ⁶ : +5.2	Dak Dinaj ⁶ : -0.5 Pa Medi ⁶ : -0.3	S 24 Parg ⁶ : 48 N 24 Parg ⁶ : 41	0
	Overweight/obesity	Murshidabad: +10.4 Maldah: +10.0	Kolkata: -11.6 Haora: -3.1	N 24 Parg ⁶ : 851 Hugli: 616	13
Women (15-49 years)	Hypertension	Darjiling: +11.4 S 24 Parg ⁶ : +10.9	Dak Dinaj ⁶ : -1.9 Utt Dinaj ⁶ : -0.8	N 24 Parg ⁶ : 611 S 24 Parg ⁶ : 544	10
	Diabetes	<i>Not Available²</i>		N 24 Parg ⁶ : 605 S 24 Parg ⁶ : 417	3
	Overweight/obesity	<i>Not Available²</i>			
Men (15-54 years)	Hypertension	Pu Medi ⁶ : +10.3 Hugli: +6.5	Dak Dinaj ⁶ : -39.1 Kolkata: -25.8	N 24 Parg ⁶ : 565 S 24 Parg ⁶ : 496	11
	Diabetes	<i>Not Available²</i>		N 24 Parg ⁶ : 769 S 24 Parg ⁶ : 552	15

Source: NFHS-3 (2005-2006) national report, NFHS-4 (2015-16) national report and data [IFPRI estimates], and NFHS-5 (2019-2021) national and state factsheets. pp: percentage points.
¹NCDs: non-communicable diseases. ²Indicates unavailability of data for a particular indicator in the specified NFHS round. Diabetes data for NFHS-4 are not included in the NFHS-5 factsheet because definition of diabetes is not comparable between NFHS-4 and 5.
³The difference is calculated only between districts that are comparable between 2015-2016 and 2019-2021. All districts except Purba Bardhaman and Paschim Bardhaman are comparable between the time periods. ⁴Burden: The headcount was calculated as the product of the overweight/obesity prevalence, and the total eligible projected population for each district in 2019. Prevalence estimates were obtained from NFHS-5 (2019-2021), and the projected population for 2019 was estimated using Census 2011. ⁵Public health concern is defined as prevalence $\geq 15\%$ for overweight/obesity (children), $\geq 20\%$ for overweight/obesity (women and men), $\geq 20\%$ hypertension (women and men), and $\geq 20\%$ diabetes (women and men) (WHO 2011). ⁶District codes: S 24 Parg- South 24 Parganas; N 24 Parg- North 24 Parganas; Dak Dinaj: Dakshin Dinajpur; Utt Dinaj: Uttar Dinajpur, Pu Medi: Purba Medinipur, Pa Medi: Paschim Medinipur.

Figure 3. Trends in immediate determinants (%)
2005-2006, 2015-2016, 2019-2021




Table 2. Immediate determinants at district-level
2015-2016, 2019-2021

Category	Immediate determinants	Worst performing districts (pp)	Best performing districts (pp)	Top coverage districts (%) ²
		<i>Difference between (2019-2021) & (2015-2016)¹</i>	<i>Difference between (2019-2021) & (2015-2016)¹</i>	<i>2019-2021</i>
Maternal determinants	Underweight women	<i>Not Applicable⁴</i>	Puruliya: -13.7 Pa Medi ³ : -11.0	N 24 Parg ³ : 6.5 Kolkata: 6.6
	Consumed IFA 100+ days	<i>Not Applicable⁴</i>	Utt Dinaj ³ : +53.2 N 24 Parg ³ : +51.8	Dak Dinaj ³ : 76.2 N 24 Parg ³ : 73.3
IYCF practices	Early initiation of breastfeeding	S 24 Parg: -20.3 Kolkata: -8.8	Pu Medi ³ : +35.3 N 24 Parg: +34.5	Nadia: 72.4 Hugli: 72.1
	Exclusive breastfeeding	Koch Bihar: -11.7 Maldah: -4.0	Puruliya: +10.2	S 24 Parg ³ : 73.5 Utt Dinaj ³ : 66.4
	Timely introduction of complementary foods	<i>Not Available⁵</i>		
	Minimum adequate diet	Nadia: -15.1 Pa Medi ³ : -10.5	Darjiling: +19.2 Kolkata: +18.1	Pu Bard ³ : 34.2 Dak Dinaj ³ : 32.6
Diseases	Diarrhea in the last 2 weeks	Bankura: +7.6 Birbhum: +3.9	Jalpaiguri: -6.0 Dak Dinaj ³ : -5.7	N 24 Parg ³ : 2.0 Dak Dinaj ³ : 2.7
	ARI in the last 2 weeks	Birbhum: +4.3 Nadia: +2.5	Koch Bihar: -5.1 N 24 Parg ³ : -3.5	Hugli: 0.8 Pa Medi ³ : 1.4

pp: percentage points.

Source: NFHS-3 (2005-2006) national and state reports and data [IFPRI estimates], NFHS-4 (2015-2016) state report and data [IFPRI estimates], and NFHS-5 (2019-21) national and state reports, factsheets, and data [IFPRI estimates].

Note: Definitions of IYCF indicators are based on WHO guidelines.

¹The difference is calculated only between districts that are comparable between 2015-2016 and 2019-2021. All districts except Purba Barddhaman and Paschim Barddhaman are comparable between the time periods.

²For all indicators, top coverage districts refer to the districts with the highest prevalence in immediate determinants, except for underweight women, diarrhea in the last 2 weeks, and ARI in the last 2 weeks, for which it refers to the districts with the lowest prevalence in coverage.

³District codes: Pu Medi- Purba Medinipur, Pa Medi: Paschim Medinipur; Pu Bard: Purba Barddhaman; S 24 Parg- South 24 Parganas; N 24 Parg- North 24 Parganas; Dak Dinaj: Dakshin Dinajpur; Utt Dinaj: Uttar Dinajpur.

⁴Prevalence did not increase or decrease in any of the districts.

⁵Indicates unavailability of data for a particular indicator in the specified NFHS round.

**Figure 4. Trends in underlying determinants (%)
2005-2006, 2015-2016, 2019-2021**



**Table 3. Underlying determinants at district-level
2015-2016, 2019-2021**

Category	Underlying determinants	Worst performing districts (pp)	Best performing districts (pp)	Top coverage districts (%) ²
		<i>Difference between (2019-2021) & (2015-2016)¹</i>	<i>Difference between (2019-2021) & (2015-2016)¹</i>	2019-2021
Maternal determinants	Women with ≥10 years education	<i>Not Applicable³</i>	Utt Dinaj ⁴ : +12.3 S 24 Parg ⁴ : +12.1	Kolkata: 55.4 Darjiling: 41.3
	Women 20-24 years married before age of 18 years	Pu Medi ⁴ : +13.6 Hugli: +8.9	Jalpaiguri: -15.8 Utt Dinaj ⁴ : -9.3	Kolkata: 16.7 Jalpaiguri: 18.7
	Women 15-19 years with child or pregnant	Pa Medi ⁴ : +9.4 Koch Bihar: +3.7	Puruliya: -9.7 Murshidabad: -8.8	Kolkata: 4.9 Darjiling: 6.3
Household determinants	HHs with improved drinking water source	Pu Medi ⁴ : -12.0 Hugli: -1.9	Darjiling: +19.0 Jalpaiguri: +10.0	S 24 Parg ⁴ : 100 Utt Dinaj ⁴ : 99.8
	HHs using improved sanitation facility	Pu Medi ⁴ : -1.4	Dak Dinaj ⁴ : +31.5 Utt Dinaj ⁴ : +30.5	Darjiling: 83.0 N 24 Parg ⁴ : 80.4
	HHs with electricity	Pa Medi ⁴ : -1.1 Kolkata: -0.3	S 24 Parg ⁴ : +11.0 Koch Bihar: +7.5	Kolkata: 99.6 N 24 Parg ⁴ : 99.5

pp: percentage points.

Source: NFHS-3 (2005-2006) national and state reports and data [IFPRI estimates], NFHS-4 (2015-2016) national and state reports and data [IFPRI estimates], and NFHS-5 (2019-2021) state and district factsheets, state report, and data [IFPRI estimates].


¹The difference is calculated only between districts that are comparable between 2015-2016 and 2019-2021. All districts except Purba Bardhaman and Paschim Bardhaman are comparable between the time periods.

²For all indicators, top coverage districts refer to the districts with the highest prevalence in underlying determinants, except for women 20-24 years married before age of 18 years and women 15-19 years with child or pregnant for which it refers to the districts with the lowest prevalence in coverage.

³Prevalence did not decrease in any of the districts.

⁴District codes: Pu Medi- Purba Medinipur, Pa Medi: Paschim Medinipur; Pu Bard: Purba Bardhaman; S 24 Parg- South 24 Parganas; N 24 Parg- North 24 Parganas; Dak Dinaj: Dakshin Dinajpur; Utt Dinaj: Uttar Dinajpur.

Figure 5. Trends in interventions across the first 1000 days (%)
2005-2006, 2015-2016, 2019-2021



Source: NFHS-3 (2005-2006) national and state reports and data [IFPRI estimates], NFHS-4 (2015-2016) national and state reports and data [IFPRI estimates] and NFHS-5 (2019-2021) state factsheet, state and national report, and data [IFPRI estimates].

Note 1: Refer to district dashboard for the inter-district variability in the coverage of interventions.

Intervention coverage at district-level, 2019-2021

<20%	20-<40%	40-<60%	60-<80%	>=80%	Not Available
------	---------	---------	---------	-------	---------------

District name	Pre-pregnancy		Pregnancy										Delivery & postnatal										Early childhood													
	Demand for FP satisfied	Iodized salt	ANC first trimester	≥4 ANC	Received MCP card	Received IFA tab/syrup	Tetanus injection	DeWorming	Weighing	Birth preparedness counselling	Breastfeeding counselling	Counselling on keeping baby warm	Cord care counselling	Food supplementation	Health & nutrition education	Malaria prevention- use of bed nets	Institutional birth	Financial assistance (ISY)	Skilled birth attendant	Postnatal care for mothers	Postnatal care for babies	Food supplementation	Health & nutrition education	Health checkup	Full immunization	Vitamin A	Paediatric IFA	DeWorming	Care seeking for ARI	ORS during diarrhea	Zinc during diarrhea	Food supplementation (6-35 months)	Weighing	Counselling on child growth	Preschool at AWC	Health checkup
West Bengal	74.6	94.5	72.6	75.8	98.4	96.1	94.6	25.3	99.4	1.5	92.1	91.3	87.6	79.1	67.2	95.0	91.7	35.2	94.1	68.0	76.8	74.9	63.5	62.9	87.8	68.4	20.4	20.0	71.3	75.3	36.0	81.7	72.6	73.4	70.7	66.5
Bankura	75.6	86.8	74.7	75.7	99.5	95.5	89.5	24.2	99.5	1.6	89.6	88.4	82.0	92.3	76.0	83.4	93.3	36.9	93.8	67.2	76.5	85.4	74.9	67.6	85.0	73.4	20.7	19.3	71.1	87.3	44.5	88.8	74.2	69.6	81.3	65.8
Birbhum	84.8	93.3	81.9	78.4	98.7	94.8	98.4	29.8	99.6	1.0	83.9	81.6	81.8	92.6	68.3	95.2	90.3	46.8	92.2	67.5	81.7	83.1	54.4	54.2	81.6	58.4	13.4	18.8	78.9	68.1	31.5	89.2	72.1	75.0	79.9	66.6
Dakshin Dinajpur	69.6	97.3	70.9	76.4	99.6	97.9	95.3	19.0	100.0	0.7	99.6	99.1	96.0	95.7	90.5	98.9	91.4	50.3	94.0	83.2	89.1	89.3	85.8	83.2	92.6	67.2	28.4	19.8	75.5	92.8	84.9	77.7	92.8	77.7	92.8	82.1
Darjiling	76.1	97.7	85.0	79.2	96.8	96.3	96.6	32.3	99.5	0.4	83.7	85.0	81.1	68.8	50.5	78.1	98.4	44.7	99.2	73.9	84.7	66.0	43.9	44.4	94.2	76.2	24.4	26.7	78.6	66.6	59.2	72.2	58.6	55.3	55.3	
Haora	78.0	97.4	84.8	82.7	96.5	97.8	35.5	99.0	0.5	88.5	89.6	85.1	69.1	69.1	56.2	96.5	95.5	40.7	96.9	70.7	79.8	70.1	60.3	61.9	87.8	70.1	25.4	24.8	53.8	72.4	74.8	70.2	76.9	69.4	69.4	
Hugli	74.8	90.5	76.0	72.2	98.6	96.6	92.2	19.1	99.5	3.7	95.6	91.1	86.7	79.9	74.9	94.8	97.0	20.1	97.4	56.7	76.7	76.6	70.7	73.3	95.4	75.0	25.4	21.7	62.8	81.0	77.4	63.5	78.9	77.8	77.8	
Jalpaiguri	80.8	94.8	77.5	88.4	100.0	97.3	96.6	29.8	100.0	1.3	96.5	96.1	94.9	75.7	66.3	99.0	95.8	43.5	97.7	74.8	82.5	74.0	63.8	61.1	87.9	65.5	23.3	21.6	80.9	76.3	68.1	74.9	65.7	64.8	64.8	
Koch Bihar	78.5	97.6	71.3	77.3	99.6	98.3	96.6	27.8	100.0	1.1	94.3	93.4	90.9	84.1	73.1	100.0	93.0	46.8	92.5	71.9	79.7	79.9	67.8	66.7	94.2	61.4	23.4	21.1	59.8	84.9	78.0	67.4	80.1	71.5	71.5	
Kolkata	82.6	98.6	74.2	71.7	94.4	91.4	100.0	20.7	98.7	0.7	93.1	91.3	80.6	38.5	32.1	80.5	97.5	18.6	99.3	72.8	84.6	38.2	29.7	31.5	80.2	58.0	18.7	16.9	39.8	36.3	81.0	39.1	33.5	33.5	33.5	
Maldah	64.5	97.8	77.8	82.8	99.7	96.3	93.1	19.0	99.2	0.7	93.4	91.7	88.6	79.8	60.0	98.1	87.7	38.4	88.9	61.6	67.9	78.6	62.0	56.7	87.2	57.7	11.1	14.5	65.0	80.7	69.7	71.4	72.2	66.4	66.4	
Murshidabad	73.0	92.5	73.9	66.9	97.3	91.6	91.1	26.1	99.0	1.0	87.3	87.0	84.5	75.1	70.8	99.3	87.0	30.6	92.3	49.8	67.6	69.3	65.8	66.2	90.0	67.7	20.3	25.6	83.0	71.5	35.9	80.4	72.6	72.5	73.1	70.2
Nadia	67.7	96.0	61.1	71.2	98.4	95.8	95.7	20.9	100.0	0.9	99.0	99.0	98.4	89.4	83.9	99.5	97.7	23.8	98.6	89.4	89.9	82.8	79.8	79.9	89.9	74.0	26.3	18.5	69.2	93.3	80.0	74.3	86.3	74.4	74.4	
North Twenty Four Pargana	73.3	98.9	72.7	89.9	98.7	97.7	97.7	20.7	98.8	1.2	99.5	98.9	98.9	80.7	80.3	99.1	97.5	32.0	98.3	87.8	92.4	74.0	73.2	68.0	92.8	78.1	23.1	22.8	77.3	74.1	90.7	79.1	72.0	72.0		
Paschim Barddhaman	69.9	89.8	70.3	70.4	95.5	93.6	96.6	22.5	99.3	0.4	74.8	70.1	63.5	55.0	29.9	72.7	89.8	31.4	91.3	64.1	70.5	49.2	24.3	27.2	62.4	60.6	12.8	18.5	86.3	81.2	48.9	57.5	36.5	69.2	43.2	33.1
Paschim Medinipur	76.9	93.8	52.6	63.1	99.1	97.8	96.2	31.7	100.0	2.1	95.8	97.5	89.8	85.0	79.2	93.8	87.6	35.3	95.1	79.7	80.9	76.1	73.1	73.9	76.9	78.3	20.6	17.6	72.1	92.3	75.8	72.4	82.7	68.8	68.8	
Purba Barddhaman	76.5	95.2	72.5	80.5	100.0	97.2	90.8	18.4	100.0	0.9	89.8	89.2	82.1	85.6	68.5	97.0	96.0	34.6	94.3	65.2	75.0	87.1	72.3	64.8	97.2	65.2	10.5	9.3	56.6	88.9	83.3	72.0	82.4	76.6	76.6	
Purba Medinipur	67.2	90.0	69.3	56.8	98.8	95.6	87.1	25.6	99.7	1.7	87.7	87.3	81.4	79.3	72.0	99.1	91.5	20.3	94.6	48.5	58.1	71.2	64.4	68.5	88.4	83.7	28.3	26.3	72.5	88.5	79.2	68.6	74.0	72.4	72.4	
Puruliya	61.5	89.7	59.1	57.8	98.0	96.2	91.2	30.0	99.5	3.6	84.6	78.2	67.1	76.0	52.6	73.8	89.8	39.7	91.4	51.5	68.4	61.8	41.3	43.5	84.7	68.0	15.2	18.1	66.7	72.8	35.6	79.3	60.1	53.5	61.1	
South Twenty Four Pargana	78.4	94.1	77.1	87.5	98.6	98.5	96.4	29.1	99.3	2.7	95.4	96.0	91.9	86.5	63.8	98.0	91.0	46.9	93.2	67.0	70.8	87.5	64.7	66.7	91.7	58.8	17.3	12.1	60.9	87.4	78.5	76.6	75.0	75.9	75.9	
Uttar Dinajpur	71.4	93.7	76.6	70.1	97.8	94.9	96.7	25.9	98.2	1.6	85.3	83.1	80.7	76.5	63.7	99.1	76.9	39.3	84.7	60.9	69.4	76.7	60.5	59.0	84.4	67.4	26.0	29.3	84.1	64.9	50.6	84.7	73.3	68.6	71.6	67.5

Source: NFHS-5 district factsheets and state reports (2019-21) and data [IFPRI estimates].

Note 1: In NFHS-5 district factsheets, estimate for indicators with sample size <25 is not reported.

Table 4. Intervention coverage at district level 2015-2016, 2019-2021

Category	Interventions	Worst performing districts (pp)	Best performing districts (pp)	Top coverage districts (%)
		<i>Difference between (2019-2021) & (2015-2016)²</i>	<i>Difference between (2019-2021) & (2015-2016)²</i>	<i>2019-2021</i>
Pregnancy	ANC first trimester	Nadia: -7.2 Kolkata: -1.5	Utt Dinaj ³ : +39.6 Maldah: +35.3	Darjeeling: 85.0 Haora: 84.8
	≥4 ANC visits	Pu Medi ³ : -22.5 Pa Medi ³ : -20.9	Maldah: +30.2 Utt Dinaj ³ : +27.0	N 24 Parg ³ : 89.9 Jalpaiguri: 88.4
	Received MCP Card	Puruliya: -2.0 Murshidabad: -1.6	Darjiling: +3.6 Maldah: +3.4	Pu Bard ³ : 100 Jalpaiguri: 100
	Tetanus injection	Pu Medi ³ : -10.9 Murshidabad: -8.2	Utt Dinaj ³ : +8.5 Kolkata: +7.8	Kolkata: 100 Birbhum: 98.4
Delivery and post-natal	Institutional birth	<i>Not Applicable¹</i>	S 24 Parg ³ : +38.9 Maldah: +32.7	Darjiling: 98.4 Nadia: 97.7
	Skilled birth attendant	<i>Not Applicable¹</i>	Maldah: +27.4 Utt Dinaj ³ : +26.3	Kolkata: 99.3 Darjiling: 99.2
	Postnatal care for mothers	<i>Not Applicable⁴</i>	<i>Not Applicable⁴</i>	Nadia: 89.4 N 24 Parg ³ : 87.8
	Postnatal care for babies	<i>Not Applicable⁴</i>	<i>Not Applicable⁴</i>	N 24 Parg ³ : 92.4 Nadia: 89.9
Early childhood	Full immunization	Pa Medi ³ : -15.2 Bankura: -11.2	Utt Dinaj ³ : +18.4 Maldah: +17.7	Pu Bard ³ : 97.2 Hugli: 95.4
	Vitamin A supplementation	Koch Bihar: -22.5 S 24 Parg ³ : -21.4	Darjiling ³ : +20.7 N 24 Parg ³ : +14.1	Pu Medi ³ : 83.7 Pa Medi ³ : 78.3
	Care seeking for ARI	Haora: -27.1 Koch Bihar: -21.8	Utt Dinaj ³ : +23.0 Pu Medi ³ : +20.2	Pa Bard ³ : 86.3 Utt Dinaj ³ : 84.1
	ORS treatment during diarrhea	Utt Dinaj ³ : -2.8	Murshidabad: +11.5	Bankura: 87.3 Pa Bard ³ : 81.2
	Zinc treatment during diarrhea	<i>Not Applicable¹</i>	Murshidabad: +28.9 Utt Dinaj ³ : +28.2	Utt Dinaj ³ : 50.6 Pa Bard ³ : 48.9

Key takeaways

- Children:** Stunting prevalence declined by 12 percentage points (pp) between 2006 and 2016 and increased by 1pp between 2016 and 2021. Underweight declined by 7pp between 2006 and 2016 and no further improvements between 2016 and 2021. Anemia declined by 7pp between 2006 and 2016 but increased by 15pp between 2016 and 2021.
- Women:** Underweight declined by 18pp between 2006 and 2016 and further declined by 6pp between 2016 and 2021. Anemia among non-pregnant women and pregnant women increased by 9pp and 8pp between 2016 and 2021, respectively. Overweight/obesity increased by 9pp between 2006 and 2016 and by 3pp between 2016 and 2021.
- Men:** Overweight/obesity increased by 8pp between 2006 and 2016 and by 2pp between 2016 and 2021.
- Attention is needed to improve** (%s in 2021):
- **Outcomes:** Stunting (34%); underweight (32%); wasting (20%) and anemia among children (69%); anemia among women (62-72%)
 - **Immediate determinants:** Consumed IFA 180+ days (31%); exclusive breastfeeding (53%); minimum adequate diet for children (23%); early initiation of breastfeeding (59%)
 - **Underlying determinants:** Women with ≥10 years education (33%); early marriage (42%); improved sanitation (68%)
 - **Coverage of interventions:** Deworming during pregnancy (25%); financial assistance (35%); zinc during diarrhea (36%)

pp: percentage points. Source: NFHS-3 state and national reports and data [IFPRI estimates], NFHS-4 (2015-2016) state and national reports and data [IFPRI estimates], and NFHS-5 (2019-2021) state and national reports/factsheets and data [IFPRI estimates]. Note: Interventions' coverage are based on the last child data. ¹Prevalence did not decrease in any of the districts. ²The difference is calculated only between districts that are comparable between 2015-2016 and 2019-2021. All districts except Purba Bardhaman and Paschim Bardhaman are comparable between the time periods. ³District codes: Pu Medi- Purba Medinipur, Pa Medi: Paschim Medinipur; Pu Bard: Purba Bardhaman; S 24 Parg- South 24 Parganas; N 24 Parg- North 24 Parganas; Dak Dinaj: Dakshin Dinajpur; Utt Dinaj: Uttar Dinajpur. ⁴Indicators postnatal care for women and babies are not comparable between NFHS-4 and 5.

Indicator definition

Nutrition outcomes	Definition
Low birth weight^{3,5}	% live births in 5 years (y) preceding the survey with a reported birth weight <2.5 kilograms (kg), based on either a written record or the mother's recall
Stunting	% children 0-59 months (m) whose height-for-age z score < -2 standard deviations (SD)
Wasting	% children 0-59 m whose weight-for-height z score < -2SD
Severe wasting	% children 0-59 m whose weight-for-height z score < -3SD
Underweight children	% children 0-59 m whose weight-for-age z score < -2SD
Anemia among children	% children 6-59 m whose haemoglobin (Hb) <11.0 g/dl
Underweight women	% women 15-49 y whose Body Mass Index (BMI) <18.5 kg/m ² ; sample excluded pregnant women and women with a birth in the preceding 2 m
Anemia among non-pregnant women¹	% non-pregnant women 15-49 y whose Hb <12.0 g/dl
Anemia among pregnant women¹	% pregnant women 15-49 y whose Hb <11.0 g/dl
Overweight/obesity – children	% children 0-59 m whose weight-for-height z score > 2SD
Overweight/obesity – women	% women 15-49 y whose BMI ≥25.0 kg/m ² ; sample excluded pregnant women and women with a birth in the preceding 2 m
Overweight/obesity – men	% men 15-49 y whose BMI ≥25.0 kg/m ²
Hypertension among women^{2,3,10}	% women 15-49 y with elevated blood pressure (systolic ≥140 mm Hg or diastolic ≥90 mm Hg) or are currently taking medication to control blood pressure
Hypertension among men^{2,3,10}	% men 15-54 y with elevated blood pressure (systolic ≥140 mm Hg or diastolic ≥90 mm Hg) or are currently taking medication to control blood pressure
Diabetes among women¹⁰	% women 15-49 y with high (>140 mg/dl) or very high (>160 mg/dl) blood sugar or are currently taking medication to control blood sugar
Diabetes among men¹⁰	% men 15-54 y with high (>140 mg/dl) or very high (>160 mg/dl) blood sugar or are currently taking medication to control blood sugar
Immediate determinants	Definition
Underweight women	% women 15-49 y whose BMI <18.5 kg/m ² ; sample excluded pregnant women and women with a birth in the preceding 2 m
Consumed IFA 100+ days¹	% women 15-49 y who consumed iron folic acid (IFA) for 100 days or more during pregnancy for the most recent live birth in the 5 y preceding the survey
Consumed IFA 180+ days¹	% women 15-49 y who consumed IFA for 180 days or more during pregnancy for the most recent live birth in the 5 y preceding the survey
Early initiation of breastfeeding^{1,2,3,8,9}	% youngest children <2 y who started breastfeeding <1 hour of birth
Exclusive breastfeeding	% youngest children 0-5 m who were fed exclusively with breastmilk the previous day
Timely introduction of complementary foods	% youngest children 6-8 m living with their mother who received solid or semi-solid food and breastmilk
Continued breastfeeding at 2 years^{1,3,4,5,6}	% youngest children 12-23 m living with their mother who were fed breastmilk during the previous day
Minimum adequate diet^{2,3,4,5,9,10}	% youngest children 6-23 m who consumed a minimum acceptable diet during the previous day
Minimum dietary diversity^{1,2,3,4,5,9}	% youngest children 6-23 m who consumed foods and beverages from at least 5 out of 8 defined food groups during the previous day
Minimum meal frequency^{2,3,4,5,9,10}	% youngest children 6-23 m who consumed solid, semi-solid or soft foods (but also including milk feeds for non-breastfed children) at least the minimum number of times during the previous day
Eggs and/or flesh foods consumption^{1,2,3,4,5,9}	% youngest children 6-23 m who consumed egg and/or flesh food during the previous day
Sweet beverage consumption^{1,2,3,4,5,9}	% youngest children 6-23 m who consumed a sweet beverage during the previous day
Bottle feeding for infants^{1,2,3,4,5,9}	% youngest children 0-23 m who were fed from a bottle with a nipple during the previous day
Diarrhea in the last 2 weeks	% children <5 y who had diarrhea in the 2 weeks preceding the survey
ARI in the last 2 weeks	% children <5 y who had symptoms of acute respiratory infection (ARI) in the 2 weeks preceding the survey
Underlying determinants	Definition
Women with ≥10 years of education	% women 15-49 y with ≥10 y of schooling
Women 20-24 years married before age of 18 years¹	% women 20-24 y who were married when <18 y
Women 15-19 years with child or pregnant¹⁰	% women 15-19 y who were mothers or pregnant at the time of the survey
HHS with improved drinking water source	% population living in households whose source of drinking water is piped into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, tanker truck, cart with small tank, bottled water, or community RO plant
HHS using improved sanitation facility	% population living in households using flush to pipe sewer system, flush to septic tank, flush to pit latrine, ventilated improved pit (VIP)/biogas latrine, pit latrine with slab or twin pit/composting toilet, which is not shared with any other household
HHS with hand washing facility^{2,4,10}	% households where a place for washing hands with water and soap was present
Open defecation^{6,7}	% households that have no toilet facility/defecate in open space or field
Safe disposal of feces^{3,4,5}	% youngest children <5 y living with their mother whose stools were disposed of safely
HHS with BPL card^{3,5,6,7}	% households possessing a BPL card
HHS with electricity	% population living in households with electricity

Note: Unless specified, state and district estimates taken from NFHS-3 national/state report for NFHS-3, NFHS-4 national/state report or NFHS-5 factsheet for NFHS-4, and NFHS-5 national/state report or factsheet for NFHS-5.

¹Indicator estimated using NFHS-3 unit-level data (state level)

²Indicator estimated using NFHS-4 unit-level data (state level)

³Indicator estimated using NFHS-4 unit-level data (district level)

⁴Indicator estimated using NFHS-5 unit-level data state level

⁵Indicator estimated using NFHS-5 unit-level data (district level)

⁶Indicator estimated using NFHS-4 unit-level data for union territories as their summary reports are not available (state level)

⁷Indicator estimated using NFHS-5 unit-level data for union territories as their summary reports are not available (state level)

⁸Indicator estimated using NFHS-5 unit-level data for union territories as their summary reports are not available (district level)

⁹Indicator constructed following WHO guidelines

¹⁰Indicator not available in NFHS-3 and/or 4 tool or not comparable between NFHS-3/4 and 5

Indicator definition

Interventions	Definition
Demand for FP satisfied ^{1,3,5}	% currently married women 15-49 years (y) with demand for family planning (FP) satisfied by modern methods
Iodized salt	% households using iodized salt
ANC first trimester	% women 15-49 y who received antenatal care (ANC) during the first trimester for the most recent live birth in the 5 y preceding the survey
≥ 4ANC¹	% women 15-49 y who received ≥4 ANC visits for the most recent live birth in the 5 y preceding the survey
Received MCP card¹⁰	% women 15-49 y who received a mother and child protection (MCP) card during pregnancy for the most recent live birth in the 5 y preceding the survey
Received IFA tab/syrup^{3,8}	% women 15-49 y who received IFA (given or purchased) during pregnancy for the most recent live birth in the 5 y preceding the survey
Tetanus injection¹	% women 15-49 y whose most recent live birth in the 5 y preceding the survey was protected against neonatal tetanus
Deworming- pregnancy^{3,8}	% women 15-49 y who took deworming medication during pregnancy for the most recent live birth in the 5 y preceding the survey
Weighing- pregnancy^{3,5,6,7}	% women 15-49 y who were weighed during ANC for the most recent live birth in the 5 y preceding the survey
Birth preparedness counselling^{1,2,3,4,5}	% women 15-49 y with a child 0-60 months (m) who had ≥1 contacts with a community health worker in the 3 m preceding the survey and were advised on birth preparedness
Breastfeeding counselling^{1,3,5,6,7}	% women 15-49 y who met with a community health worker in the last 3 m of pregnancy for the most recent live birth in the 5 y preceding the survey and were advised on breastfeeding
Counselling on keeping baby warm^{1,5,6,7}	% women 15-49 y who met with a community health worker in the last 3 m of pregnancy for the most recent live birth in the 5 y preceding the survey and were advised on keeping the baby warm
Cord care counselling^{5,6,7,10}	% women 15-49 y who met with a community health worker in the last 3 m of pregnancy for the most recent live birth in the 5 y preceding the survey and were advised on cord care
Food supplementation – pregnancy^{3,5}	% women 15-49 y with a child <6 y who received food supplements from ICDS during pregnancy
Health & nutrition education – pregnancy⁵	% women 15-49 y with a child <6 y who received health and nutrition education from ICDS during pregnancy
Malaria prevention- use of bed nets^{2,4,5,10}	% women 15-49 y who slept under a treated bed net during pregnancy of the most recent live birth in the 5 y preceding the survey
Institutional birth	% live births in the 5 y preceding the survey to women 15-49 y delivered in a health facility
Financial assistance (JSY)^{6,7,8,10}	% women 15-49 y who received financial assistance under Janani Suraksha Yojana (JSY) for the most recent live birth that took place in a health facility in the 5 y preceding the survey
Skilled birth attendant	% live births in the 5 y preceding the survey to women 15-49 y that were assisted by a skilled provider
Postnatal care for mothers¹⁰	% women 15-49 y who received postnatal care from a doctor/nurse/LHV/auxiliary nurse midwife (ANM)/midwife/other health personnel <2 days of delivery for the most recent live birth in the 5 y preceding the survey
Postnatal care for babies¹⁰	% most recent live births in the 5 y preceding the survey to women 15-49 y with a postnatal check for the newborn from a doctor/nurse/LHV/ANM/midwife/other health personnel <2 days of delivery
Food supplementation – postnatal^{3,5}	% women 15-49 y with a child <6 y who received food supplements from ICDS during breastfeeding
Health & nutrition education – postnatal^{3,5}	% women 15-49 y with a child <6 y who received health and nutrition education from ICDS during breastfeeding
Health checkup (ICDS)^{3,5}	% women 15-49 y with a child <6 y who received a health check-up from ICDS during breastfeeding
Full immunization	% children 12-23 m fully vaccinated based on information from either vaccination card or mother's recall
Vitamin A – early childhood¹	% children 9-35 m who received a vitamin A dose in the 6 m preceding the survey
Pediatric IFA^{3,4,5,10}	¹¹ % children 6-59 m who were given iron supplements in the 7 days preceding the survey ¹² % children 6-36 m who were given iron supplements in the 7 days preceding the survey
Deworming – early childhood^{3,4,5,10}	¹¹ % children 6-59 m who received deworming medication in the 6 m preceding the survey ¹² % children 6-36 m who received deworming medication in the 6 m preceding the survey
Care seeking for ARI	% children <5 y with fever or symptoms of ARI in the 2 weeks preceding the survey taken to a health facility or health provider
ORS during diarrhea	% children <5 y with diarrhea in the 2 weeks preceding the survey who received Oral Rehydration Solutions (ORS)
Zinc during diarrhea	% children <5 y with diarrhea in the 2 weeks preceding the survey who received zinc
Food supplementation (children 6-35 months)^{1,2,3,4,5}	% youngest children 6-35 m who received food supplements from ICDS in the 12 m preceding the survey
Weighing – early childhood^{3,5}	% children <5 y who were weighed at an anganwadi centre (AWC) or ICDS centre in the 12 m preceding the survey
Counselling on child growth^{3,5,10}	% mothers with a child <5 y who received counselling from an ICDS/anganwadi worker or ANM after they were weighed at an AWC in the 12 m preceding the survey
Preschool (AWC)^{3,5}	% children 36-71 m who went for early childhood care/preschool at an AWC in the 12 m preceding the survey
Health checkup (AWC)^{3,5}	% children <6 y who received health checkups from an AWC in the 12 m preceding the survey

Note: Unless specified, state and districts estimates taken from NFHS-3 national/state report for NFHS-3, NFHS-4 national/state report or NFHS-5 factsheet for NFHS-4, and NFHS-5 national/state report or factsheet for NFHS-5.

¹Indicator estimated using NFHS-3 unit-level data (state level)

²Indicator estimated using NFHS-4 unit-level data (state level)

³Indicator estimated using NFHS-4 unit-level data (district level)

⁴Indicator estimated using NFHS-5 unit-level data state level

⁵Indicator estimated using NFHS-5 unit-level data (district level)

⁶Indicator estimated using NFHS-4 unit-level data for union territories as their summary reports are not available (state level)

⁷Indicator estimated using NFHS-5 unit-level data for union territories as their summary reports are not available (state level)

⁸Indicator estimated using NFHS-5 unit-level data for union territories as their summary reports are not available (district level)

⁹Indicator constructed following WHO guidelines

¹⁰Indicator not available in NFHS-3 and/or 4 tool or not comparable between NFHS-3/4 and 5

¹¹Definition as per NFHS-3 and 4 construction

¹²Definition as per NFHS-5 construction

Led by IFPRI 

AUTHORS

Sattvika Ashok, Research Analyst, IFPRI
Phuong Hong Nguyen, Senior Research Fellow, IFPRI
S.K. Singh, Professor, IIPS
Rakesh Sarwal, Additional Secretary, NITI Aayog
Neena Bhatia, Senior Specialist, NITI Aayog
Robert Johnston, Nutrition Specialist UNICEF
William Joe, Assistant Professor, IEG
Esha Sarswat, Communications Specialist, IFPRI
Purnima Menon, Senior Research Fellow, IFPRI

SUGGESTED CITATION

Ashok, S., P.H. Nguyen, S.K. Singh, R. Sarwal, N. Bhatia, R. Johnston, W. Joe, E. Sarswat, and P. Menon. 2022. *State Nutrition Profile: West Bengal*. POSHAN Data Note 89. New Delhi, India: International Food Policy Research Institute.

ACKNOWLEDGEMENTS

Financial support for this Data Note was provided by the Bill & Melinda Gates Foundation through POSHAN, led by the International Food Policy Research Institute. The funder played no role in decisions about the scope of the analysis or the contents of the Note.

We thank Long Quynh Khuong (Independent Researcher) for creating the maps, Nishmeet Singh (IFPRI) and Anjali Pant (IFPRI) for working with the dataset and Julie Ghostlaw (IFPRI) & Abhilasha Vaid (Consultant) for editing and reviewing the Note.

PARTNERS

Institute of Economic Growth (IEG)
International Institute for Population Science (IIPS)
NITI Aayog
UNICEF



Disclaimer: The maps used in this Data Note are based on the districts in NFHS-5 factsheets/reports. The boundaries shown do not imply any official endorsement or acceptance by IFPRI.

ABOUT POSHAN

Partnerships and Opportunities to Strengthen and Harmonize Actions for Nutrition in India (POSHAN) is a multi-year initiative that aims to support the use of data and evidence in decision-making for nutrition in India. It is supported by the Bill & Melinda Gates Foundation and led by IFPRI in India.
<http://poshan.ifpri.info/>

ABOUT DATA NOTES

POSHAN Data Notes focus on data visualization to highlight geographic and/or thematic issues related to nutrition in India. They draw on multiple sources of publically available data.

CONTACT US

Email: IFPRI-POSHAN@cgiar.org

IFPRI-NEW DELHI INTERNATIONAL FOOD POLICY RESEARCH INSTITUTE

NASC Complex, CG Block,
Dev Prakash Shastri Road,
Pusa, New Delhi 110012, India
T+91.11.66166565
F+91.11.66781699

IFPRI-HEADQUARTERS INTERNATIONAL FOOD POLICY RESEARCH INSTITUTE

1201 Eye Street, NW,
Washington, DC 20005 USA
T. +1.202.862.5600
F. +1.202.467.4439
Skype: IFPRIhomeoffice
ifpri@cgiar.org
www.ifpri.org

This publication has been prepared by POSHAN. It has not been peer reviewed. Any opinions stated herein are those of the author(s) and do not necessarily reflect the policies of the International Food Policy Research Institute.

Copyright © 2022 International Food Policy Research Institute. All rights reserved. For permission to republish, contact ifpri-copyright@cgiar.org.