

What are people eating in Rajshahi, Bangladesh?

Food consumption, food security, food sources, and food perceptions

Data Note 3

December 2023

ABOUT THIS DATA NOTE |

The Transforming Agrifood Systems in South Asia (TAFSSA) district agrifood systems assessment aims to provide a reliable, accessible, and integrated evidence base that links farm production, market access, dietary patterns, climate risk responses, and natural resource management with gender as a cross-cutting issue in rural areas of Bangladesh, India, and Nepal. It is designed to be a district-level multi-year assessment. Using data collected in February–March 2023, this data note describes what people are eating, where they get their food, household food insecurity, and perceptions about food. This is one of a set of data notes that, together, provide a holistic picture of the agrifood system in the district.

Figure 1. District location in Bangladesh





Figure 2. Highlights from this data note



OVERVIEW OF CONTENTS

TAFSSA's district agrifood systems assessment aimed to interview three respondents per household: a female adult (aged 20+ years), a male adult (aged 20+ years), and an adolescent (aged 10-19 years). Information on the household and respondent sampling strategy is provided at the end of this data note.

In this data note, you will first find information on background characteristics of the households and individuals who were interviewed. This is followed by information on *what* people are eating, which was captured using several measurement methods. Respondents were asked about the foods they ate the day before the interview (24-hour recall) and about how often they ate certain foods in the past week (food frequency questionnaire). The 24-hour recall was conducted using the Global Diet Quality Score (GDQS) application, which also captured when (at what eating occasion such as breakfast, a snack between lunch and dinner, etc.) people ate each food item.

In addition to what people eat, you will find information in this data note on *where* they get their food and, if they buy it, what types of markets or shops they buy it from.

Finally, you will learn *why* people choose to eat certain healthy and unhealthy foods. Respondents were asked about availability, accessibility, taste, and other factors that may influence their decisions to consume certain foods. More details about the measurement methods are found on the following pages.

List of figures and tables

Figure 1. District location in Bangladesh

Figure 2. Highlights from this data note

Table 1. Household and individual characteristics

Figure 3A. Level of household food insecurity

Figure 3B. Individual FIES items

Figure 4A. Minimum dietary diversity

Figure 4B. Global diet quality positive (healthy) and negative (unhealthy) scores

Figure 4C. Diet related NCD risk

Figure 5. Consumption of food groups on previous day

Figure 6. Consumption quantity by GDQS food group

Figure 7. Most commonly consumed foods in each food group

Figure 8. Number of eating occasions per day

Figure 9. Who eats at different eating occasions

Figure 10. Frequency of consumption of sentinel foods in previous 7 days

Figure 11. Food sources

Figure 12. Where food is purchased

Figure 13. Food perceptions (% who agree with each statement)

Table 1. Household and individual characteristics

Household characteristics		Individual characteristics			
Number	1000				
Female-headed, %	8				
Education of head, years	5				
Average household size, members	4				
Involved in agriculture, %	91				
Has improved toilet, %	98				
Drinking water source					
Tube well or borehole, %	89				
Piped to neighbour, %	4				
Main source of income					
Crop cultivation, %	41				
Business, %	25				
Wages, %	22				
Type of fuel used for cooking					
Wood, %	88				
Straw/grass, %	78				
Dung cake, %	45				
		Number	968	449	1000
		Age, yrs. mean (range)	37 (20-71)	44 (20-72)	15 (10-19)
		Education, yrs. mean (range)	5 (0-17.5)	5 (0-17.5)	7 (0-11.5)
		Married, %	96	98	15
		Employed, %	9	99	10
		Primary occupation			
		Unpaid household work, %	92	0	14
		Farming, %	2	46	1
		Own enterprise or business, %	1	15	1
		Student, %	0	0	78

Measuring household food insecurity

Household food insecurity was measured using the Food and Agriculture Organization's Food Insecurity Experience Scale (FIES). Respondents were asked 8 questions about their household's experiences with food in the past 30 days. Categories of food security (none, mild, moderate, severe) were generated based on the number of questions with a positive response.

Fig 3A. Level of household food insecurity

■ None (0) ■ Mild (1-2) ■ Moderate (3-5) ■ Severe (5-8)



Fig 3B. Individual FIES items



FINDINGS

- ✓ Nearly one third of households reported experiencing moderate or severe food insecurity.

MEASURING WHAT PEOPLE EAT

Diets were measured by asking people about everything they ate or drank on the previous day, from the time they woke up until the time they went to bed and didn't eat or drink anything more. This includes all snacks and foods and drinks consumed at home and outside the home.

To capture this information, we used the Global Diet Quality Score (GDQS) application (Bromage et al. 2021). The GDQS allows us to understand diet quality, which is associated with the risk of disease. We report the percentage of individuals with at least minimum dietary diversity (FAO and FHI 360, 2016) (**Figure 4A**), that is those who consume at least 5 of the following 10 food groups daily: 1) grains, white roots and tubers, and plantains, 2) pulses (beans, peas, and lentils), 3) nuts and seeds, 4) dairy, 5) meat, poultry, and fish, 6) eggs, 7) dark green leafy vegetables, 8) other vitamin A-rich fruits and vegetables, 9) other vegetables, and 10) other fruits.

We also computed metrics that indicate how healthy or unhealthy diets are (**Figure 4B**). Higher GDQS+ and GDQS- scores indicate better diet quality. We then grouped GDQS scores into 3 categories to indicate diet related noncommunicable disease (NCD) risk (**Figure 4C**).

On the following pages, we show the percentage of individuals who consume various food groups (**Figure 5**), the consumption quantity by food group (**Figure 6**), the most commonly consumed foods (**Figure 7**), how many times per day people eat (**Figure 8**), who eats at various eating occasions (**Figure 9**).

Diet quality scores

Fig 4A. Minimum dietary diversity






Fig 4B. Global diet quality positive (healthy) and negative (unhealthy) scores




Note: The GDQS+ and GDQS- can be summed to give the total GDQS score, with a higher total GDQS score indicating better diet quality.

Fig 4C. Diet related NCD risk



FINDINGS

- ✓ Only about one-third of individuals consumed at least 5 of 10 food groups, but diet diversity was higher among adolescents compared to adults.
- ✓ Adult females were at higher diet related NCD risk compared to adult males and adolescent.

Figure 5. Consumption of food groups on previous day**FINDINGS**


- ✓ Consumption of fruits was <40% for all respondent types, and consumption of dark green leafy vegetables was <20% for all respondent types.
- ✓ Fish was the most commonly consumed animal-source food, and around 84% of individuals consumed some animal-source foods on the previous day.
- ✓ Adult males were the most likely to consume sweets & ice-cream

Note: High-fat dairy and red meat are considered unhealthy when consumed in high quantities.



Animal-source foods

23% of adult women and 25% of adult men consumed eggs on the day before the survey

Figure 6. Consumption quantity¹ by GDQS food group**FINDINGS**

- ✓ Few individuals consumed “high” quantities of healthy foods and “low/none” quantities of unhealthy foods.
- ✓ To reduce the risk of developing noncommunicable diseases, consumption of healthy foods such as fruits and whole grains should increase and consumption of unhealthy foods such as refined grains/baked goods, sweets & ice-cream, and white roots & tubers should decrease.

¹ “Low”, “medium”, and “high” describe consumption levels that predict noncommunicable disease risk in studies from Africa, Asia, and the Americas (Bromage et al. 2021). Disease risk is minimized when healthy foods are consumed in “high” quantities and unhealthy foods are consumed in “low” quantities.

Figure 7. Most commonly consumed foods in each food group (adult male, adult female, and adolescent combined)

	Top 3 foods consumed in each food group				
	% who consumed food group	1 st most common	2 nd most common	3 rd most common	
Healthy	Citrus fruits	1	N/A ¹	N/A ¹	N/A ¹
	Deep orange fruits	1	N/A ¹	N/A ¹	N/A ¹
	Other fruits	34	Holud boroi (yellow plum)	Banana	Guava
	Dark green leafy vegetables	15	Pui shak	Lal shak	NA ¹
	Cruciferous vegetables	6	Cabbage	Shak	NA ¹
	Deep orange vegetables	4	Pumpkin	NA ¹	NA ¹
	Other vegetables	99	Onion	Chili pepper	Brinjal
	Legumes	45	Masoor dal	Dal	Pulses
	Deep orange tubers	2	Carrots	NA ¹	NA ¹
	Nuts & seeds	5	Almonds	Nuts	NA ¹
	Whole grains	33	Brown rice	Wheat flour	Roti
	Liquid oils	100	Soybean oil	Mustard oil	Oil
	Fish & shellfish	55	Silver carp	Rui	Bata
	Poultry & game meat	8	Chicken	NA ¹	NA ¹
	Low fat dairy	2	N/A ¹	N/A ¹	N/A ¹
	Eggs	27	Eggs	Chicken eggs	Duck eggs
	Unhealthy	High fat dairy	7	Milk	N/A ¹
Red meat		5	Beef	N/A ¹	N/A ¹
Processed meat		17	Chicken	Beef	N/A ¹
Refined grains/baked goods		99	Rice	Roti	Puffed rice
Sweets & ice-cream		55	Sugar	Biscuit	Mishti
Sugar-sweetened beverages		3	N/A ¹	N/A ¹	N/A ¹
Juice		1	N/A ¹	N/A ¹	N/A ¹
White roots and tubers		90	Potatoes	Aloo bhartha	Cooking banana
Purchased deep fried foods		19	Onion	Wheat flour	Jalebi
Deep fried foods at home		2	N/A ¹	N/A ¹	N/A ¹

Figure 8. Number of eating occasions per day (mean)





Figure 9. Who eats at different eating occasions



FINDINGS

- ✓ Adult women ate fewer times per day than adult men or adolescents; this was due to fewer adult women eating food during pre-breakfast, mid-morning or afternoon eating occasions.
- ✓ More than 90% of respondents ate at the 3 main eating occasions (breakfast, lunch, dinner), but foods were often consumed outside of these 3 eating occasions.


¹Not applicable (N/A): Foods consumed by less than 2% of the respondents have not been displayed.

SENTINEL FOODS | In addition to the GDQS, which provided information about all foods consumed in the previous 24 hours, we selected a set of 25 “sentinel foods” to better understand how frequently these foods are consumed, food sources, where people buy food, and their perceptions about food.

Respondents were asked how frequently they consumed these foods in the past 7 days (**Figure 10**). They were also asked about where their household gets each food (purchased from outside, own production, received from others, received from government, gather/forage) (**Figure 11**). If they said their household purchases the food, we asked them where it was purchased (haat, retail shop, farm, government ration shop, or other market type) (**Figure 12**).


For a few foods, we dug deeper to understand people’s food perceptions, or what they think about the foods. This included whether they know of a vendor who sells the food, if the food is safe to eat, easy to acquire near where they spend most of their time, is not too expensive, is fast and easy to prepare, tastes good, fills their stomach, is nutritious, and if their family enjoys eating it (**Figure 13**).

Understanding these perceptions provides insights into drivers or barriers of consumption of healthy and unhealthy foods.




Sentinel food list


1. Rice
2. Wheat
3. Maize
4. Millets
5. Moong dal
6. Masoor dal
7. Chana dal
8. Chickpeas and beans
9. Potato
10. Poultry (chicken, ducks, pigeons, etc.)
11. Fish
12. Other meat (e.g., mutton)
13. Eggs
14. Milk (e.g., cow, buffalo, goat)
15. Orange vegetables (e.g., pumpkin, carrots)
16. Green leafy veg. (e.g., spinach, mustard, taro, pumpkin leaves, red amaranth leaves)
17. Onions
18. Tomatoes
19. Fruits (e.g., guava, banana, apple, mango)
20. Instant noodles (e.g., Maggi, Wai Wai)
21. Chips (e.g., Lays, Kurkure)
22. Biscuits and baked sweets (e.g., cakes and cookies, mithai)
23. Deep fried food (e.g., samosa, pakora)
24. Soda/soft drinks and packaged juices (e.g., Coke, Sprite, Fanta, Maaza)
25. Tea/coffee with sugar

Figure 10. Frequency of consumption of sentinel foods in previous 7 days**FINDINGS**

- ✓ Rice, potatoes, and onions were consumed daily by nearly all respondents.
- ✓ Animal-source foods were typically consumed a few times a week.
- ✓ Adult women consumed tea/coffee with sugar less frequently than adult men.


Figure 11. Food sources**FINDINGS**

- ✓ Most households purchased their food rather than producing it themselves.
- ✓ About one-third of households consumed rice from their own production.
- ✓ Receiving food from the government or gathering/foraging food was not a common food source.

Figure 12. Where food is purchased**FINDINGS**

- ✓ Food was predominantly purchased from haats.
- ✓ Milk was often directly purchased from the farmer/producer rather than through a market.
- ✓ Unhealthy foods were purchased from both haats and retail shops.

Note: Data shown are for the subset of households who purchase the food from outside (Figure 11). "Other" includes weekly markets, city markets, roadside markets, wholesale market, mobile vendors, and any other purchase source. Haats are wholesale markets where foods are sold in bulk directly by manufacturers/ farmers/ artisans at a fair price, in permanent or semi-permanent infrastructure. Retail shops means fixed or mobile individual shops where foods are sold directly to the consumers, including local grocery stores, specialized shops, vegetable/fruit shops, restaurants, and tea stalls.

Figure 13. Food perceptions (% who agree with each statement)**FINDINGS**

- ✓ Nearly all adult men knew a shop that sold the different foods; adult women and adolescents were less knowledgeable; knowledge of a shop selling biscuits was the highest.
- ✓ Eggs were perceived as a more affordable animal-source food than fish.
- ✓ Few respondents considered deep fried foods safe to eat or nutritious.

Note: GLVs = green leafy vegetables.

KEY TAKEAWAYS

1. There is room for improvement in diet quality.
 - Consumption of healthy food groups such as whole grains, nuts & seeds, fruits, and deep orange vegetables is low.
 - Consumption of unhealthy food groups such as refined grains/baked goods, sweets, and sugary tea/coffee is high.
 - Consumption of starchy foods with low nutrient density (rice, potatoes) is high.
 - Compared to adult male and adolescents, adult females are more likely to be at “high” diet related NCD risk.
2. Most households purchase food from markets rather than producing it themselves.
3. Food purchases are typically from haats, followed by retail shops.

KEY QUESTIONS FOR ACTION

1. What are the key barriers to improving diet quality in the district?
2. What are a few potential solutions to overcome these barriers? What is needed from decision-makers and from program teams to implement these solutions?
3. How can understanding eating patterns throughout the day and perceptions about healthy and unhealthy foods help inform strategies to influence consumption of these foods?
4. What additional information is needed to facilitate actions to improve diets in the district?



SURVEY METHODOLOGY

Village and household sampling

We selected 25 villages in the district with a probability proportional to the number of households that reside in each village. Within each village, we conducted a household listing to identify eligible households, that is, those with adolescents (10-19 years old). From the households with adolescents, we randomly invited 20 households to participate in the survey. If a household refused, we replaced that household with another randomly selected eligible household, to retain a total of 1,000 households in the district. Thus, the findings reported in this data note are representative of rural households from this district that include an adolescent.

Respondent selection

Within households, one adult female aged 20+ years, one adult male aged 20+ years, and one adolescent aged 10-19 years were selected as the respondents for the survey. When multiple adolescents were living in a household, the oldest adolescent was selected. In some households, an adult male was not available (often due to migration for work). In such households, the female was the only adult respondent (see Table 1 for respondent sample sizes). At the beginning of the interview, the adult in the household primarily involved in agriculture (either male or female) and the adult primarily responsible for food purchasing (either male or female) were identified as the primary respondents.

INDICATOR DEFINITIONS

Indicator	Definition
GDQS+	The Global Diet Quality Score positive sub-metric is composed of the summed score of the 16 healthy food groups and ranges from 0-32 points, with a higher score indicating higher diet quality.
GDQS-	The Global Diet Quality Score negative sub-metric is composed of the summed score of the 7 unhealthy food groups and the 2 food groups that are unhealthy when consumed in excess amounts (high-fat dairy and red meat), with a higher score indicating lower diet quality. The GDQS negative score has a possible range of 0-17.

REFERENCES

Bromage, S., C. Batis, S.N. Bhupathiraju, et al. 2021. "Development and Validation of a Novel Food-Based Global Diet Quality Score (GDQS)." *Journal of Nutrition* 151,(10S), Supp. 2.

FAO and FHI 360. 2016. *Minimum Dietary Diversity for Women: A Guide for Measurement*. Rome: FAO.



AUTHORS

Samuel Scott, Research Fellow, IFPRI
Sharvari Patwardhan, Research Analyst, IFPRI
Suman Chakrabarti, Associate Research Fellow, IFPRI
Alka Chauhan, Research Analyst, IFPRI
Archis Banerjee, Research Analyst, IFPRI
Sumanta Neupane, Program Manager, IFPRI
Aklima Parvin, Program Manager, IFPRI
Sabrina Rasheed, Scientist, icddr,b
Kaosar Afsana, Professor, James P Grant School of Public Health, BRAC University
Nazma Shaheen, Professor, University of Dhaka
Christine Blake, Associate Professor, University of South Carolina
Purnima Menon, Senior Director for Food and Nutrition Policy, CGIAR & IFPRI
Avinash Kishore, Senior Research Fellow, IFPRI

REFERENCES

Bromage, S., C. Batis, S.N. Bhupathiraju, et al. 2021. "Development and Validation of a Novel Food-Based Global Diet Quality Score (GDQS)." *Journal of Nutrition* 151,(10S), Supp. 2.

FAO and FHI 360. 2016. *Minimum Dietary Diversity for Women: A Guide for Measurement*. Rome: FAO.

SUGGESTED CITATION

Scott S, Patwardhan S, Chakrabarti S, Chauhan A, Banerjee A, Neupane S, Parvin A, Rasheed S, Afsana K, Shaheen N, Blake C, Menon P, Kishore A. 2023. What are people eating in Rajshahi, Bangladesh? TAFSSA Data Note 3. New Delhi, India: Transforming Agrifood Systems in South Asia (TAFSSA).

FUNDING ACKNOWLEDGEMENT

We would like to thank all funders who supported this research through their contributions to the CGIAR Trust Fund:

<https://www.cgiar.org/funders/>

To learn more, contact: a.kishore@cgiar.org

To learn more about TAFSSA, contact: t.krupnik@cgiar.org; p.menon@cgiar.org

ABOUT TAFSSA

TAFSSA (Transforming Agrifood Systems in South Asia) is a CGIAR Regional Integrated Initiative that supports actions improving equitable access to sustainable healthy diets, that boosts farmers' livelihoods and resilience, and that conserves land, air, and water resources in a climate crisis.

ABOUT CGIAR

CGIAR is a global research partnership for a food secure future. Visit <https://www.cgiar.org/research/cgiar-portfolio> to learn more about the initiatives in the CGIAR research portfolio

DISCLAIMER

The views and opinions expressed in this publication are those of the author(s) and are not necessarily representative of or endorsed by CGIAR, centers, our partner institutions, or donors. This data note has not been peer reviewed.