

TOPICAL REVIEW • OPEN ACCESS


Assessing the impact of nature-based solutions on soil health in sub-Saharan Africa through farmer-centred methods

To cite this article: Dominik Bittner *et al* 2026 *Environ. Res. Lett.* **21** 043004

View the [article online](#) for updates and enhancements.

You may also like

- [Temporally clustered streamflow events control focused groundwater recharge in drylands](#)
Gabriel C Rau, José Bastías Espejo, R Ian Acworth et al.
- [Nature-based solutions for freshwater conservation and equitable adaptation in the Amazon basin: a case study review](#)
Paola Carolina Stuparich Nouel, Sophia Picarelli, Fernanda Silva et al.
- [Arctic warming delays the Afro-Asian monsoon retreat amplifying autumn rainfall](#)
Jie Zhang, Haishan Chen, Weiqiang Ma et al.

ENVIRONMENTAL RESEARCH
LETTERS

TOPICAL REVIEW

OPEN ACCESS

RECEIVED

24 October 2025

REVISED

13 January 2026

ACCEPTED FOR PUBLICATION

16 January 2026

PUBLISHED

26 February 2026

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.

Assessing the impact of nature-based solutions on soil health in
sub-Saharan Africa through farmer-centred methods

Dominik Bittner^{1,2,*} , Jo Smith¹ , Georgios Leontidis² , Grant A Campbell¹ , Jeanne Biegel³ ,
Pete Smith¹, Matthias Kuhnert¹, Rastislav Skalský⁴, Lícida M Giuliani¹ , Abdul Walid Salik¹ ,
Paul Hallett¹, David F R P Burslem^{1,2}, Getahun Yakob⁵, Wolde Mekuria⁶, Euan Phimister^{7,8},
Amare Hailelassie⁶, Desalegn Tegegne⁶, Sarem Norouzi⁹ , Heyu Chen¹, Cecile Gubry-Rangin¹
and Adnan Khan¹⁰

¹ Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, United Kingdom

² Interdisciplinary Institute, University of Aberdeen, Aberdeen, United Kingdom

³ University of Burgundy, Institut Agro, Dijon, France

⁴ Biodiversity and Natural Resources Program, International Institute for Applied Systems Analysis, Laxenburg, Austria

⁵ Central Ethiopia Agricultural Research Institute, Durame, Ethiopia

⁶ International Water Management Institute (IWMI), East Africa and Nile Basin Office, PO Box 5689, Addis Ababa, Ethiopia

⁷ Business School, University of Aberdeen, Aberdeen, United Kingdom

⁸ Stellenbosch Business School, Stellenbosch University, South Africa

⁹ Department of Agroecology, Aarhus University, Tjele, Denmark

¹⁰ Information and Computational Sciences James Hutton Institute, Dundee, United Kingdom

* Author to whom any correspondence should be addressed.

E-mail: d.bittner.24@abdn.ac.uk

Keywords: artificial intelligence, decision-support system, Ethiopia, low to middle income countries, machine learning, process-based models, systems models

Supplementary material for this article is available [online](#)

Abstract

Soils underpin many ecosystem services, including food production, through functions such as organic matter decomposition. These functions are increasingly threatened by soil degradation, especially in climate-vulnerable regions, such as sub-Saharan Africa, where unstable soils are prone to severe erosion. As soils continue to degrade, farmers face multiple challenges; they cannot afford accurate tests to assess soil, their livelihoods are constrained by demand for food, fuel and water, and competition for valuable resources hampers farming. Hence, there is a pressing need for accessible tools to assess soil health and methods to provide tailored advice on resilient, climate-smart agricultural management and optimal use of resources. This narrative review offers a comprehensive overview of key issues and potential solutions. We highlight tools and approaches that can support farmers to improve soil and secure livelihoods. Practical indicators and field-ready tests are evaluated, with examples from Ethiopia, but tailored to support farmers and advisors across sub-Saharan Africa and other developing countries. A wide range of tests are reviewed, including physical, biological, chemical, function and service-related tests, drawing on scientific and farmers knowledge. Science-based tests require expertise, equipment and incur costs, while locally-derived tests are affordable and seamlessly applicable. We also review Nature-based Solutions for improving soil quality, and assess them against factors such as labour, costs, and crop production. There is no single universally applicable practice; suitability depends on farmers' priorities and circumstances. Therefore, we explore predictive methods—mechanistic, process-based soil models, data- and knowledge-driven Artificial Intelligence and systems models—to simulate the impact of practices on soil and farm dynamics. Promising approaches include hybrid approaches assimilating data, physics and knowledge through digital soil mapping.

Overall, this review emphasizes the need to empower farmers with accessible tools and methods to harness Nature-based Solutions, build climate resilience and secure sustainable futures for generations ahead.

1. Introduction

Humanity faces many global challenges that are closely interlinked and threaten human livelihoods, including environmental degradation, food insecurity and a changing climate. One specific component of many terrestrial ecosystems plays a significant role in all of these challenges; the soil [1–3]. More than 95% of the food produced depends directly or indirectly on soils [4, 5]. Despite this strong dependence of the human population on soil, this complex ecosystem is still relatively poorly understood [5, 6] and poorly managed in many regions [7]. Approximately 33% of the earth's soil are moderately to highly degraded as a result of different threats, many of which are influenced by human activities, including agricultural practices [2, 3, 8–10]. A common phenomenon worldwide is that topsoil (depth of 0–30 cm) has been almost entirely lost across substantial portions of currently or formerly intensively farmed land [8]. Future soil erosion rates are difficult to predict accurately and estimates can vary significantly. However, researchers strongly agree that rates will increase, particularly in the least developed economies, such as in many countries in sub-Saharan Africa (SSA), South America and Southeast Asia [8, 11–13]. While Africa's population is the fastest growing in the world, hunger and soil loss are simultaneously limiting the ability of African farmers to achieve long-term, sustainable food production [14, 15]. This phenomenon is referred to as “*the cycle of land degradation and social deprivation*“. It is characterised by a downward spiral in which severe land degradation, low farm productivity, poverty, low levels of education, inadequate infrastructure and social exclusion are inextricably linked [16–19]. In particular, many countries in SSA face considerable environmental and social challenges, and so find themselves at the core of this cycle. They are threatened by extreme soil degradation due to their climatically vulnerable regions and unstable soils [20]. For example, according to Nkonya *et al* [15], 85% of Ethiopia's soils are moderately to very severely degraded, with approximately 75% of the country being affected by desertification. Current rates of soil erosion on arable land in Ethiopia are estimated to vary between 42 and 300 t ha⁻¹ y⁻¹. This wide range is attributable to complex spatial and temporal variations, and methodological difficulties in estimating rates. The most common causes of land degradation in Ethiopia are erosion by water, wind and tillage [21–23]. With approximately 80% of the Ethiopian population employed in rain-fed agriculture, the economic cost of land degradation,

estimated at around \$4.3 billion y⁻¹, is a serious concern [15, 24]. If these issues are not addressed, degradation of land, and therefore, social deprivation, are estimated to become more severe across all of Ethiopia [25, 26]. Similar issues exist in many other countries in SSA. Breaking and potentially reversing this cycle requires a holistic approach based on established concepts related to soil [16].

1.1. Soil-related concepts

As outlined by Bünemann *et al* [27], the provision of services by the soil is entirely contingent on its underlying functions. These functions depend on the quality, health and fertility of the soil, which in turn are affected by the various soil threats.

1.1.1. Soil-based ecosystem services

Soil is not only vital for food, fuel and fibre production. It also provides other (soil-based) ecosystem services, such as biodiversity conservation, carbon (C) sequestration and erosion control [10]. In this context, ecosystem services are defined as ‘*the benefits which humans derive from ecosystems*’ [28]. These can be grouped into provisioning (biomass production, water supply), regulating (climate regulation, pest, disease and erosion control, water quality, and biodiversity conservation), and cultural services [27, 29–31]. Given the significant challenges of food security across Africa, especially in low to middle income countries (LMICs), many with rapidly growing populations, the sustainable production of biomass is critical [32, 33]. However, services, such as biodiversity conservation, should not be neglected, as their feedback on soil functions affect the current state of the soil as well as biomass production [29, 32, 34].

1.1.2. Soil functions

Ecosystem services are underpinned by specific soil functions, including habitat provision, element cycling (including nutrient cycling), decomposition, soil structure maintenance, biological population regulation, water cycling (retention, percolation/infiltration) and organic matter cycling (humus formation and C sequestration) [27]. According to Glenk *et al* [35], these functions are composed of ‘*(bundles of) soil processes that support the provision of ecosystem services*’ and are sometimes categorised as supporting ecosystem services.

1.1.3. Soil threats

The current state of soil, and consequently the provision of ecosystem services, is directly influenced by soil threats. These include erosion (e.g. through

water or wind), soil organic matter decline, contamination (e.g. with heavy metals, such as copper or iron in Ethiopian soils [36]), sealing, compaction, biodiversity loss, salinisation, landslides and floods [27]. According to Baritz et al [37], ‘soil threats are indicative circumstances which can damage or reduce a soil’s capacity to provide ecosystem services’.

1.1.4. Soil quality and soil health

In the literature, the terms ‘soil quality’ and ‘soil health’ are used more or less synonymously [27]. According to Doran et al [38, 39], soil quality is defined broadly as ‘the capacity of a soil to function within ecosystem and land-use boundaries to sustain biological productivity, maintain environmental quality, and promote plant and animal health’; here, human health is included in animal health. In contrast to this, Hannam et al [40] define soil health as ‘the ability of soil to perform its functions and to deliver ecosystem goods and services. The range of functions and ecosystem services provided should reflect the different capabilities of different soils—a “healthy” soil is therefore one in which ecosystem services are provided at an acceptable level given inherent underlying constraints and the purpose of the land use’. In essence, soil quality is more concerned with the soil’s ability to meet specific human needs. Soil health, on the other hand, is focused on the long-term ability of the soil to support plant growth and maintain its functions [27]. In this paper, the term ‘soil health’ is adopted because it more clearly reflects soil as a living organism [41, 42].

1.1.5. Soil fertility

The concept of ‘soil fertility’ focuses mainly on crop yields. It is defined as ‘the ability of the soil to supply essential plant nutrients and soil water in adequate amounts and proportions for plant growth and reproduction in the absence of toxic substances which may inhibit plant growth’ [27]. It is imperative to prioritise the interplay between soil fertility and soil health, because fertility is important for food production as a soil-based ecosystem service for the sustainable nutrition of the populations of countries in SSA [7, 16, 43].

1.2. Soil governance

In recent decades, considerable efforts have been made to address the numerous challenges facing humanity. These include the introduction of the Sustainable Development Goals and a range of citizen science initiatives.

1.2.1. Sustainable Development Goals and policies

As part of the United Nations’ multilateral blueprint for the Sustainable Development Goals (SDGs), a total of 17 different goals are being targeted which are either directly or indirectly related to soil. For example, goals of ending hunger (SDG 1) or clean water and sanitation (SDG 6) are highly dependent on soil-based ecosystem services, such as

biomass production [44–48]. In addition to these global initiatives, continental and national policies also play an important role as a driving force for sustainable development. These policies are largely in line with the SDGs, with specific adaptations to local circumstances [49–51]. In the context of SSA, national and local policies support the implementation of sustainable development by integrating them into political strategies, allocating relevant budgets, promoting investment, and facilitating collaborative and community-led efforts, as evidenced by initiatives in Ghana [52, 53]. However, while these measures aim to facilitate implementation, in some cases they can hinder progress. This may be due to inconsistent financing strategies, and lack of funding across sectors, leading to siloed approaches [54–56]. Without co-operation between different regulatory bodies and across multiple levels, from regulatory bodies to farmers, addressing human and environmental challenges appears to be less achievable.

1.2.2. Citizen science approaches

Alongside the interest and desire of high-level decision-makers, the commitment, participation, ownership and responsibility of citizens, specifically farmers as ground-level decision-makers, are becoming increasingly important [57, 58]. Regardless of the location, most farmers are interested in ‘looking after’ their soil. This applies not only to current and near future crop production and therefore income, but also, if they own individual property rights, for passing on fertile land to their descendants. From day-to-day business, working on and with the land all year round and knowledge being passed down from ancestors, farmers generally hold extensive, localised knowledge, as well as personal memories of their land [45]. While their experience and knowledge can vary greatly, it is worth noting that these types of emotional relationships with the environment can be of similar value in tackling human challenges as international high-level initiatives [45, 59]. This arises because it is usually the responsibility of farmers not only to follow the guidelines ‘handed down from above’ and adapt their farming practices, but also to maintain an acceptable income and feed their families. Therefore, the inclusion of farmers in community-based programmes (e.g. environmental monitoring, project design or implementation) [45] could be of great potential benefit. This can include large-scale initiatives, for example, where farmers can take in-field soil measurements themselves and contribute them to a database. Not only do farmers learn about their land from a different perspective and feel the health of their own soil, but they also provide data to projects, which rely on a large amount of data for critical scientific analysis and decision-making. Moreover, the involvement of farmers, as important stakeholders throughout a project enables tailored project development and the incorporation of local

and indigenous knowledge into the decision-making process. This is particularly useful through participatory workshops and the creation of fuzzy cognitive maps [45, 59, 60]. Furthermore, the needs of farmers tend to outlast high-level initiatives and projects due to the nature of specific funding periods and their commitments, while farmers remain responsible for managing the land, producing food and generating income over the long-term [61]. The fight against poverty and hunger, the need to support the health of people and nature and the achievement of many SDG targets depend partly (directly and indirectly) on how the threats to soil health are addressed. Only with joined forces and integrative initiatives can soil threats be mitigated and soil health improved; this requires appropriate indicators and measures to be selected.

1.3. Improving food production and soil health

Reducing hunger and breaking the '*cycle of land degradation and social deprivation*' requires an understanding of the soil, its threats and services, and also the possible strategies to mitigate them. A deep understanding is extremely valuable for farmers when it comes to making appropriate management decisions. Soil health can be of benefit, not only to food production, but also to the wider environment. Here, food production is at the core, although sustainability aspects are no less important, as they aim to increase resilience to extreme weather events and halt degradation while ensuring a long-lasting sustainable biomass production.

1.3.1. Agricultural intensification

Short-term crop productivity could be improved through agricultural intensification (on a smaller area), characterised by widespread tillage and the use of external inputs. However, this approach can lead to significant loss of soil organic matter, depletion of soil nutrients, negative effects on other soil services and a catastrophic deterioration of soil health with lasting environmental pollution [62–65]. In addition, smallholder farmers from LMICs often have only limited access to external inputs due to financial and market-related restrictions. To maintain long-term food production without further depleting resources, and compromising soil health, quality and fertility, a different solution must be found to maintain productivity. A more accessible yet sustainable agricultural approach is needed for LMICs that does not compromise food production, particularly in countries already under pressure due to soil degradation or where sections of the population are suffering from malnutrition.

1.3.2. Sustainable agriculture

The concept of sustainable agriculture and the approaches and practices it entails appear to be a

feasible way forward [66, 67]. It includes biophysical aspects, such as '*building healthy soil and preventing erosion, managing water wisely, minimising air and water pollution, storing C on farms, increasing resilience to extreme weather, and promoting biodiversity*', as well as socio-economic aspects, such as '*maintaining profitability to support farming families while contributing to their local economies*' [65, 68]. Improving soil health is a crucial part of many aspects of sustainable agriculture. A healthy soil is more resilient to soil threats and is better able to provide soil functions that contribute to ecosystem services, which in turn help to achieve the SDGs. There is compelling evidence in favour of sustainable practices, demonstrating that improving soil health yields substantial benefits, including increased crop production, while enhancing resilience to soil hazards [69–71]. Sustainable agriculture includes approaches, such as agroecology, nature-inclusive agriculture, organic farming, regenerative agriculture and climate-smart agriculture [66, 67]. These include several sustainable practices, such as crop rotation, crop diversification, cover cropping and intercropping. All of these have significant effects on the quantity and quality of organic inputs to the soil [72, 73]. These approaches and practices are in line with the three central pillars of sustainable development, i.e. environmental, social and economic sustainability [66].

1.3.3. Nature-based solutions

Valuable tools for sustainable development to reduce biodiversity loss and the threats to soil are Nature-based Solutions (NbSs) [67]. These approaches work by enhancing soil structure, improving nutrient cycling, increasing habitat diversity and restoring degraded lands, thereby reinforcing the ecological processes that support resilient and productive farming systems. NbSs are defined as '*actions to protect, sustainably manage and restore natural or modified ecosystems, which address societal challenges (e.g. climate change, food and water security or natural disasters) effectively and adaptively, while simultaneously providing human well-being and biodiversity benefits*' [59, 66, 74]. They are increasingly recognised in global sustainability agendas, including the United Nations SDGs and the United Nations Decade on Ecosystem Restoration, as key approaches to tackling interconnected environmental and societal challenges. The application of NbS in agriculture is relatively diverse. They can be used in different ecosystem conditions (natural, managed, modified or artificial) to simultaneously deliver co-benefits that address diverse societal challenges, such as climate regulation, biodiversity conservation, food production, recreation and human health [75]. However, the adoption of NbS can also involve trade-offs, including potential short-term yield reductions, increased labour or management requirements, and competition for land, which must be carefully evaluated in

each context. Furthermore, as with other management practices, long-term economic viability cannot be estimated precisely and must be assessed on a case-by-case basis using cost-benefit analysis. However, it is assumed that NbS tend to improve production as they are more affordable and sustainable than conventional approaches, which should benefit the viability and overall uptake of NbS in SSA [76].

The overarching concept of NbS comprises a variety of practices, including crop rotation, cover cropping, afforestation, agroforestry and watershed restoration. In many instances these practices have been used for decades, and are grounded in indigenous knowledge or were recognised under different frameworks of sustainable agriculture [59, 75, 77]. Although an extensive list of NbS for improving soil health and food production exists, there is no universally applicable solution. Every farmer requires tailor-made solutions for each individual field, depending on the climate, ecosystem type, regional context, geographical features and the farmer's socio-cultural conditions [45].

1.4. Research gaps, questions, objectives and contributions

In view of the pressing threats to soil, urgent measures are required to mitigate these challenges. To enhance soil health, it is first essential to determine the current condition of soil health and identify limiting factors. However, the poverty faced by farmers requires the availability of simple, low-cost tools to assess soil. Once tests are conducted, a multitude of NbS can be applied to restore and strengthen soil functions, but their specific benefits in relation to farmers' primary concerns, such as labour, costs, and crop yields, must be clearly communicated to encourage adoption. A further complication stems from the food–fuel–water nexus, where limited and valuable farm resources, such as livestock manures, cannot be used as soil amendments because they are used for other purposes, such as fuel for cooking [78]. To ensure the optimal use of resources and the adoption of the most appropriate NbS according to the farmers' circumstances, predictive models can be used to simulate soil processes and farm dynamics. These tools can provide helpful guidance to farmers. However, numerous models are available to assess the potential effects of NbS on soil health and livelihoods. Their relevance needs to be clarified, as models should explicitly address ecosystem services that are most critical to farmers, including crop production and the prevention of soil erosion.

Despite the existence of literature that analyses available soil health assessment methods (e.g. [5, 79–81]) and approaches for improving (e.g. [44, 67]) and modelling soil health (e.g. [82–85]), these aspects are often disconnected from each other and do not offer a single, comprehensive solution to the problem. Previous research on farmer-based soil health

assessment has often focused on conditions where farmers are not faced with such extreme soil degradation problems as seen in SSA and are therefore not so disadvantaged by poor soil health. Furthermore, these farmers often are not able to afford expensive and elaborate soil health test kits. In addition, the literature often fails to establish a clear link between the specific test for soil health, and the targeted soil functions and ecosystem services; this is essential to ensure the relevance and reliability of the test.

The present manuscript aims to address these knowledge gaps via a comprehensive, narrative literature review. The narrative (snowballing) approach offers the opportunity to cover a broad spectrum of relevant topics (see supplementary figure S1 for a systematic mapping of the literature across these topics), which would not be possible with other methods, such as the systematic review. The work combines a review of methodologies to assess and improve soil health and provide actionable advice to farmers and agricultural advisors in SSA. Therefore, the aim of this study is to answer the following research questions (RQs):

- **RQ1:** Which soil health assessment methods are available for use by farmers in SSA and how do these link to soil functions and ecosystem services?
- **RQ2:** What NbS can be used to improve soil health in farmlands in SSA?
- **RQ3:** What methods can be used to predict the impact of these NbS on soil health to enable actionable advice to farmers?

To address RQ1, relevant criteria for soil health metrics are discussed, highlighting their links to ecosystem services and pathways for interpreting and integrating soil health tests. Subsequently, agricultural approaches and practices for improving soil health are examined in order to answer RQ2. Thereafter, modelling methods available to assess the potential impact of agricultural practices on soil health are reviewed to answer RQ3. This study concludes with a discussion and a summary of the most suitable tests, approaches and models that could be used to assist farmers and agricultural advisors in making informed decisions.

1.5. Feasibility, constraints and challenges

The review includes tests and models that require the use of smartphones. These could offer significant advantages and potential solutions to the problem, now and increasingly in the future, as the availability of smartphones in SSA increases while their cost decreases [86]. The findings of this review are likely to remain valid in the coming years. However, at present, access to smartphone access is limited for smallholder farmers in SSA. The World Bank Group [87] reported that in the year 2018, approximately 16% of the rural population in Ghana, 14% in Kenya,

9% in Nigeria, 2% in Mozambique and only 1% in Rwanda owned a smartphone. In addition, network coverage across SSA is relatively sparse, with limited connectivity especially in remote areas [87–89]. Apart from the technological feasibility, other aspects, such as the financial resources and the level and type of education of farmers vary greatly. According to reports by the World Bank Group and the Food and Agriculture Organization, in 2024 around 67% of the people in SSA were living in extreme poverty on less than 2.15 USD per person per day, while in 2017, around 60% of the rural household members in SSA earned less than 1 USD per day [90, 91]. This constrains the ability of rural households in many aspects of economic activity, beyond buying food, fuel or medicine. Household financial circumstances not only play a central role with respect to access to smartphones and material for soil tests, but also indirectly through education. This means that wealthier farmers are likely to be better able to benefit from the functions potentially made available through smartphones [87]. In addition to technological, financial and educational constraints, there are also political challenges that pose major obstacles to the transition from current agricultural practices to more sustainable approaches. This applies in particular to land ownership in many countries in SSA. For example in Ethiopia, farmers do not own the land, nor do they pass it on to their descendants. Instead, they lease it from the state for life, returning it after their death [92]. The state may then redistribute or lease the land to someone else. Consequently, the same land is not cultivated by the same family across generations. This can lead to the current owners placing less emphasis on soil health and longevity during their tenure than if they were passing it on to descendants who would also have to live off the land [92]. Furthermore, socio-cultural factors such as gender roles within predominantly patriarchal and hierarchical systems, which govern and influence power relationships within societies, traditional practices and mistrust towards new technologies could pose obstacles to the transition [93, 94]. These facts need to be considered when designing and deploying solutions.

2. Soil health assessment in the farmlands of sub-Saharan Africa

As Doran and Safley [42] in Pankhurst *et al* [41] stated in 1997, the ‘assessment of [...] soil health is invaluable in determining the sustainability of soil and land management systems and in evaluating their long-term effectiveness.’ Because of the complexity of soil, its’ health cannot be measured easily, let alone directly; a singular metric or value is not sufficient to describe the level of soil health adequately. However, it is also not possible (or necessary) to measure all soil properties; this means that an appropriate balance in the number of soil properties used to assess soil

health is required [95]. Soil health is usually assessed in relation to its functions, ecosystem services and threats. This means that measuring the health of the soil requires it to be examined from multiple angles. At the same time it is also important to repeat individual soil health measurements at sufficient locations across an entire field to characterise variability. This is particularly true when external factors, such as topography, are not homogeneous.

2.1. Static and dynamic soil properties, and extrinsic factors

All physical, chemical and biological properties of soil are influenced by extrinsic factors, e.g. parent material, climate, topography, hydrology, organisms, time and management. These properties are classified into relatively static (inherent) and dynamic (manageable) properties, although this distinction is not absolute [27, 96]. The classification also depends on the sensitivity of the soil property to external influences, such as weather, natural disturbances or agricultural management. For example, texture and stone content are relatively static soil properties, whereas organic C content, aggregate stability and nutrient availability are relatively dynamic properties, which change more quickly in response to changes in land-use or management practices [97]. Water content, temperature and nutrients change even more rapidly with time, affecting how soil health influences ecosystem services within a growing season. The differentiation in types of properties is important, as static properties provide a baseline for the potential of soils in terms of their functions and services. In contrast, dynamic soil properties vary more significantly on a small scale and are used to evaluate the actual state of the soil (i.e. soil health, relative to its potential) and to monitor temporal changes [98].

2.2. Criteria for choosing soil health indicators

Different dynamic soil properties could serve as reliable indicators for soil health assessment, depending on the extrinsic factors. Pankhurst *et al* [41] define the term ‘indicator’ in this respect as a ‘measurable surrogate for ecosystem services or soil functions’. Such surrogates, usually physical, chemical or biological, provide information on the soil’s ability to function. However, entire ecosystem services or soil functions can also serve as indicators, such as the crop yield (ecosystem service ‘biomass production’) or the decomposition of standard organic amendments on the forest floor (soil function ‘decomposition’), which reflects the diverse fauna that would otherwise be challenging to measure [99]. One of the most commonly measured dynamic soil properties across all the categories is soil organic matter, often expressed as soil organic C (SOC). With respect to farmers in SSA, SOC is seen as the main indicator of soil health and fertility, and is crucial for crop growth and productivity, especially in

Table 1. T-shaped matrix showing the relationship between soil-based ecosystem services and soil functions as well as soil threats and soil functions adapted from Bünemann *et al* [27], CC BY 4.0, whereas a cross (X) marks a consistent, unweighted relationship between the corresponding row and the specific soil function in this column [95]. The green shaded columns ‘biomass production’ and ‘erosion control’ indicate the most urgent services to be targeted [32, 33] and the red shaded column ‘erosion’ indicates the greatest threat to soil in farmlands in SSA [21, 22]; SOM = soil organic matter; C = carbon; N = nitrogen.

	Soil threats								Soil functions		Soil-based ecosystem services				
											Provision	Regulation			
Erosion	SOM decline	Contamination	Sealing	Compaction	Biodiversity loss	Salinization	Landslides & floods				Biomass production	Water quality & supply	Biodiversity conservation	Erosion control	Pest & disease control
	X	X	X	X	X	X	X	X	Habitat provision	X			X	X	X
	X	X	X		X				Element cycling (e.g. C, N cycling)	X	X				X
	X	X		X	X		X		Soil structure maintenance	X			X	X	
		X			X				Biological population regulation	X					X
	X			X	X		X	X	Water cycling	X	X	X	X	X	X
		X	X	X			X	X	Organic matter decomposition	X	X		X		X

low-input, resource-limited agroecosystems of small-holder farmers [72, 100]. According to Pankhurst *et al* [41], the following criteria should be considered for selection of the most appropriate indicators:

- **Link:** An indicator should be linked via soil functions to relevant ecosystem services and soil threats, as shown in table 1;
- **Time:** An indicator should be time-effective to measure *in-situ* with respect to sampling, hardware, analysis and labour [101, 102];
- **Cost:** The indicator should be cost-effective to measure [101, 102];
- **Ease of use:** The indicator should be easy to measure [101, 102];
- **Sensitivity:** An indicator should be responsive to variations in management on a timescale of 1–3 years [101, 102], and not strongly influenced by short-term weather patterns [27].

In addition, Bünemann *et al* [27] report on some other requirements for soil health indicators/tests:

- **Correlation:** An indicator should be correlated to long-term responses of the soil;
- **Reliability:** An indicator should allow accurate and reliable measurements, with limited subjectivity [101, 102];
- **Interpretability:** A test should be clear (absolute) and easy to interpret [101, 102];
- **Redundancy:** An indicator should not overlap with other indicators.

Hughes *et al* [101] highlight the necessity for soil health indicators and tests to be practical for farmers, particularly considering the skills required and the associated costs. In addition, particular attention

should be given to the applicability of the assessment methods to the present soil conditions and the destructiveness of the tests:

- **Applicability:** A test should be applicable in terms of the local pedological conditions.
- **Destructiveness:** A test should be as non-destructive as possible.

2.3. Assessment of soil health indicators

Any assessment of soil health should include the measurement of several indicators, including at least one indicator from each of the three categories: biological, physical and chemical properties [27]. Five different types of soil health indicator assessments exist, although the boundaries between them are not strict and assessments can sometimes be mapped to several types: (1) Laboratory assessment, often referred to as analytical assessment; (2) In-field analytical assessments that require specialist apparatus; (3) In-field visual or rapid assessments, e.g. field-based test kits for profile or spade methods; (4) Sensor-based assessment, including e.g. proximal and remote sensing; (5) Modelling, e.g. through pedotransfer functions. Furthermore, Artificial Intelligence-based approaches represent a state-of-the-art alternative to the conventional methods mentioned above. Laboratory, sensor-based, modelling and Artificial Intelligence based assessments usually yield quantitative (continuous) data, while in-field visual approaches mainly produce qualitative (categorical) data [81, 103].

2.3.1. Laboratory assessment

Quantitative methods that can be used to assess ecosystem services require analytical solutions, and these tend to be laboratory-based, such as the assessment

of SOC through a standard method (for example loss on ignition) [104]. The results from the laboratory are usually more accurate than other approaches but tend to be costly and require more expert knowledge and time for analysis. They are accessible to farmers mostly at a relatively high cost, either directly or indirectly as part of an extension service.

2.3.2. In-field analytical assessment

Analytical field-based approaches that provide quantitative data generally require specialist equipment, such as for measuring water transport (infiltrometer), erodibility (rainfall simulator) or root growth mechanical restriction (penetrometer). Simpler versions of some of these tests exist, which are directly accessible to farmers for lower costs, but these are usually not as reliable.

2.3.3. In-field visual assessment

By contrast, the most accessible approaches for farmers to assess soil health are visual in-field methods. These use empirical indicators that are easy to learn and apply. Such visual approaches emphasise the educational aspect of soil health assessment, provide immediate results and can foster better communication between farmers, advisors and scientists [27]. Approaches could be as simple as counting earthworms, assessing soil colour, visually characterising soil structure, or estimating aggregate stability from dispersion upon immersion in water (e.g. Emerson test) [105]. Combined soil health tests can also be used, such as the Muenchberg Soil Quality Rating [106], or others [107–112], which tend to be more complex. As Hussain *et al* state [5], low-cost equipment, intuitive interpretation and immediate results are clear strengths of these field assessment methods. However, their scope is sometimes limited. For instance, they are unable to assess the status of biologically and chemically mediated soil ecosystem services, and the quality of the final results is highly influenced by the experience of the sampler [27, 113]. The assessments also do not provide quantitative data, such as on water transport/retention or nutrient status, so they are not directly transferable to assess ecosystems services.

2.3.4. Sensor-based assessment

Proximal sensors, such as hand-held devices, can be used in the field to estimate soil properties with greater accuracy than a visual assessment [114]. Examples of such sensors are spectroradiometers which measure the near-infrared reflection of the soil to assess factors, such as particle size distribution, water retention or soil organic matter content [115, 116]. Some sensors can also be placed directly in the soil, for example, to measure moisture or solar radiation [117, 118]. The use of such sensors by farmers has expanded considerably, but the costs make

them inaccessible to most in SSA. Other technologies, such as satellites or drones, can perform similar measurements remotely (from a long distance), such as estimating soil moisture from satellite images [119]. The results of these measurements could be made available to farmers through smartphones.

2.3.5. Model-based assessment

In instances where the assessment of a particular indicator does not fulfil the requirements for a useful indicator, for example the ease and cost-effectiveness of the measurement, an attempt can be made to model a relationship to the indicator by measuring other properties. This can be done using pedotransfer functions, to provide a proxy value for a secondary soil property, as described by Greiner *et al* [27, 120, 121]. In this context, pedotransfer functions are ‘simple to complex knowledge rules that relate available soil information to soil properties and variables’ [122].

A prominent use of pedotransfer functions is to infer the difficult-to-measure water retention curve from readily measured attributes, such as particle-size and bulk density [116, 123]. Most of the mathematical-statistical frameworks used as a basis for the pedotransfer functions are entirely empirical and purely data-driven, based on data from databases or look-up tables [122]. Consequently, the performance of a pedotransfer function is only as good as the data used for parameterisation, so for data-poor regions, such as SSA, their reliability can be compromised.

2.3.6. Artificial intelligence-based assessment

Artificial Intelligence (AI)-based models are increasingly being adopted to estimate soil health indicators and avoid the limitations of pedotransfer functions in terms of their accuracy and reliability [27]. The discipline of AI, which is defined as ‘the science of making machines do things that would require intelligence if done by men’ [124–126], entails various methodologies of interest for assessing soil health. Solutions using AI can improve the efficiency of estimates while reducing the complexity of the approach, whether used in soil health assessment or in modelling the impact of management on soil health. This is achieved by establishing data-based relationships without the requirement for detailed knowledge about soils [127]. Based on the AI categories generally accepted in the literature [128–130], Wadoux [124] proposed a taxonomy of AI in soil science, encompassing the following three categories; reasoning and decision-making, learning and prediction, and sensing and interaction. The AI-based assessments of soil health indicators are mainly located in learning and predicting, as well as sensing and interacting. While the domain of learning and prediction involves applications, such as Machine Learning, data mining and

Table 2. A summary of selection methods, which are categorised according to their type, to reduce the number of soil health indicators and derive at a minimum data set.

Type	Method	Source(s)
Qualitative	Expert judgement	[38, 145]
	Participatory approach	[146]
	Logical-sieve approach	[146, 147]
Semi-quantitative	Decision trees	[142, 148]
	Bayesian Belief Networks	[149, 150]
Quantitative	Correlation analysis	[81]
	Analysis of Variance	[151]
	Partial least squares	[152]
	Principal component analysis	[153, 154]
	Redundancy analysis	[155]
	Regression analysis	[156]

pattern recognition, applications in sensing and interaction aim to establish a link between AI and the physical world, including robotics, computer vision and image processing.

An example of this is where the time-consuming and cost-intensive procedures of laboratory analyses can be streamlined through the application of AI models to analyse large data sets and potentially identify patterns more effectively than using manual analysis [131, 132]. Pedotransfer functions can also be supplemented with AI to bridge the gap between available and required data for predicting inaccessible indicators, utilising methods, such as neural networks [116, 119, 122]. A common computer vision-based approach used in soil science is the identification of patterns to predict soil properties from space through satellites, whereby environmental covariates from products (e.g. Normalized Difference Vegetation Index) or variables (e.g. topography or slope profile) are sourced to predict soil properties (e.g. pH) [127, 133, 134]. Another example of a sensor-based assessment, that greatly benefits from computer vision or other AI approaches, is proximal sensing, particularly when assessing soil properties in the field using a smartphone [135–137]. Here, models (e.g. neural networks) are used to extract information from images of the topsoil layer to estimate properties (e.g. SOC content), as demonstrated through the SOCit app [124, 127, 138–140].

2.3.7. Combined assessment

There are advantages and disadvantages to different assessment types, so combining them could be of great benefit [27, 141, 142]. Given the challenging circumstances of farmers in SSA, the options for assessing soil health are limited. Based on the above soil health indicator criteria, certain types of assessments are avoided, such as those that require laboratory analysis, as they are unlikely to provide the cost and time-efficient assessment that farmers in SSA require.

For farmers, in-field assessments are favoured because they are fast and cost-effective. In cases

where technology is available, such as a smartphone, proximal sensors through phone cameras, data from satellite imagery and modelling can be integrated. This could significantly improve the accuracy of the in-field assessment in a relatively quick and inexpensive way [139]. Smartphones could be used to communicate educational aspects of soil health, as well as providing guidance about subsequent soil management.

2.3.8. Selection of assessment methods

An appropriate balance in the number of indicators and assessment methods is key. More measurements generally provides more information. However, the increase in the number leads to a greater chance of collinearity between the indicators. Too many indicators also increase the cost, the workload and the expertise required to assess soil health [27, 143]. Bünemann *et al* [6] and Rinot *et al* [81] identified different approaches to derive a minimum data set (table 2). The final number of assessment methods selected typically ranges from six to eight [27]. A number of studies [79, 144, 145] demonstrate that a small set of tests, carefully chosen based on expert opinion and participatory approaches, can provide adequate information for land management decisions while enabling a streamlined process [101].

2.3.9. Evaluation of assessment methods

This section highlights potential soil health indicators and their assessment methods, which are evaluated according to the criteria for suitability (section 2.2) for the example of Ethiopian farmlands. Although this assessment refers to suitability in the Ethiopian context, it is assumed that the soil health indicators and tests are equally applicable and effective in other SSA and LMIC contexts. This assumption is also supported by the fact that the list is compiled from resources focusing on science-based [27, 29, 45, 79, 80, 101, 113, 157] and farmer knowledge-based methods from across SSA [144, 158–166]. It does not include methods that are based solely on remote

sensing techniques (see section 2.4.3) as this is outside the scope of this paper. However, it does include smartphone-based tests, which could partly be based on Earth Observation data, such as the SOCit app. Each method is linked to the corresponding soil functions to indicate its relevance for assessing an indicator of soil health. The legend for the heatmap evaluation can be found in table 3. The heat map, presented in the format of a traffic light system, shows favourable ratings in green (e.g. quick test), through to amber for less favourable ratings and then into red for those which are not favourable at all (e.g. time-consuming or unreliable test). In this way, the indicators and tests listed in table 4 can be compared effectively based on their advantages and disadvantages, using expert judgement, and then selected based on the frequency of the respective colours. Additional criteria mentioned in section 2.2, such as the redundancy with other tests for similar indicators, influence the final selection. The final selection of tests should be refined by involving farmers in the decision-making process through participatory approaches. To summarise the assessment given in table 4, based on the heat map and considering minimal redundancy of tests within each indicator category, a number of indicators offer particularly high value to Ethiopian farmers (RQ1). These are:

- Visual evaluation of soil structure (VESS),
- Colour,
- Penetration resistance,
- Beerkan test,
- Slakes app (Emerson test for farmers without smartphone access),
- pH test,
- SOCit app (colour, feel and smell test for farmers without smartphone access),
- Simplified tropical soil biology and fertility (TSBF) test
- Root development,
- Teabag index,
- Crop yield and
- Gully retreat.

Most in-field tests are associated with physical indicators, such as the VESS test. This test is widely used to assess soil structure, and can be adapted to Ethiopian conditions. A common test derived from farmers' knowledge is the analysis of the colour of the soil. This test can be carried out relatively quickly and reliably without any costs or expert knowledge required. To determine the degree of soil compaction caused by cultivation and the depth of the soil (static property), farmers can use a fencing wire to penetrate the soil. Another widely used method is the Beerkan test, which, using minimal equipment, helps farmers to assess the soil's ability to infiltrate water. The stability of soil aggregates could be determined using a smartphone and the Slakes app. If farmers do not

have access to a smartphone, the Emerson test offers a suitable alternative with a similar approach, whereby water is added to soil aggregates and their dispersion is assessed. Evaluating chemical and biological indicators in-field appears to be limited. Some measurements, such as soil pH, require specialist equipment, but it is feasible to measure pH with relatively low-cost strips that could be provided to farmers. The assessment of pH is critical, due to its significant contribution to multiple soil functions and the importance of pH in regulating microbial activity and population size. The importance of SOC for biomass production and erosion control makes it another property that is crucial to assess. With limited reliability, farmers can get a sense of SOC from the colour, feel and smell of their soil without using a smartphone. Farmers can compare their agricultural land to parts of their farm under different land use, such as a home garden, which in Ethiopia tends to receive more organic inputs [226], or perennial planting, such as enset. If smartphones are available, moving to sensor-based solutions, such as the SOCit app, is highly recommended, as these provide a more reliable and interpretable SOC assessment at no extra cost or need for additional equipment.

In terms of biological indicators, mainly visual tests which account for soil fauna, e.g. termites or roots, are found to be applicable in Ethiopian farmlands, mostly because of the cost of other tests (see Head *et al* [79]). However, both tests listed in table 2 (the simplified TSBF test and assessment of root development) come with limitations, including time to conduct the test, reliability and the expertise required to interpret the results. Entire soil functions can alternatively be represented by indicators, such as the rate of decomposition, and assessed with tests such as the 'teabag index'. Although other soil function tests, such as dung removal, toilet paper decomposition or the predation of seeds, also assess the biodiversity, they mainly target above-ground fauna [99].

Beyond this, wider ecosystem service assessments have also been noted, such as assessing the extent of weed infestation, relative crop yield or the rate of retreat of gullies. These can be derived from farmers' knowledge, and prove valuable for farmer-based soil health assessment. For example, the assessment to identify the retreat of gullies using pegs is relatively cost- and time-efficient as well as straightforward to interpret [224]. Once specific tests to assess soil on a one-off basis are selected, each test result must be contextualised and interpreted individually to fully understand changes in soil health.

2.4. Contextual interpretation of soil health indicators

To identify problems, monitor changes or make realistic predictions for agricultural management in relation to the soil, it is essential that the measured

Table 3. Heatmap legend using the traffic light system colours green (G), amber (A) and red (R) as adopted by [79] CC BY 4.0, indicating the suitability of the solution from high (G) to very low (R). The categories relate to the criteria outlined in section 2.2 with a short description including placeholders A and B (in italic). These placeholders are to be replaced by corresponding values appropriate for each classification, e.g. under disturbance, the test requires either no soil sample (G) or some soil (A). The threshold of USD3.0 is based on the poverty threshold highlighted by the World Bank in June 2025 [167]. Some criteria assume a training day to be held for farmers to get familiarised with the tests prior to independent usage.

	Time	Cost/Accessibility	Reliability	Ease of use	Interpretability	Applicability	Destructiveness	Sensitivity
	Measurement and interpretation either occurs only once a year and take <i>A</i> or occur multiple times annually and take <i>B</i> each time.	In addition to a spade, a trowel and a smartphone, the test requires <i>A</i> , which is readily accessible in Ethiopia.	<i>A</i> studies show <i>B</i> between the listed method and laboratory methods in terms of error and reliability in detecting changes.	After a training day, farmers require <i>A</i> help to conduct the test without compromising the accuracy and reliability of the results.	After conducting the test, farmers require <i>A</i> help to interpret the results without mistakes.	The test and its protocol can be used in Ethiopia <i>A</i> adaptations.	The test requires <i>A</i> .	The indicator and its test results change <i>A</i> in response to land-use changes or management practices.
Green (G)	<i>A</i> = < 30 min <i>B</i> = < 15 min	<i>A</i> = no further equipment	<i>A</i> = Several <i>B</i> = no significant difference	<i>A</i> = no	<i>A</i> = no	<i>A</i> = without	<i>A</i> = no soil sample.	<i>A</i> = within one year
Amber (A)	<i>A</i> = < 60 min <i>B</i> = < 30 min	<i>A</i> = equipment not exceeding \$3.0	<i>A</i> = Several/Few <i>B</i> = no significant difference/similar trends	<i>A</i> = some	<i>A</i> = some	<i>A</i> = with few/small	<i>A</i> = some soil that could be returned after the tests.	<i>A</i> = within five years
Red (R)	<i>A</i> = > 60 min <i>B</i> = < 60 min	<i>A</i> = equipment exceeding \$3.0	<i>A</i> = Several/Few <i>B</i> = no agreement	<i>A</i> = extensive	<i>A</i> = extensive	<i>A</i> = with significant	<i>A</i> = digging a hole and extract an important amount of soil that disturbs the concerned area.	<i>A</i> = not within five years

Table 4. Soil health indicators with corresponding science-based and farmer knowledge-based assessment methods. Each test is linked directly to soil functions (marked with a X) and thus to ecosystem services and threats in Ethiopian farmlands (see table 1). Indirect effects of indicators or tests on soil functions are not considered. The tests, which are mainly in-field assessments (see section 2.3.2), with the exception of the sensor-based Slakes and SOCit app (see section 2.3.3), are evaluated using a heatmap based on the criteria (see section 2.2) as shown in table 3. The costs of a spade/trowel and a smartphone are not considered under 'Cost' as these items can be used for other purposes. The tests are categorised into the most relevant indicators as described in section 2.2 due to the multitude of influences and overlaps. Tests based on farmers' knowledge are framed in black, and tests colour-coded according to their category offer Ethiopian farmers particularly high value. Indicators based on static soil properties are included for completeness and marked with *. VESS refers to visual evaluation of soil structure. TSBF test refers to tropical soil biology and fertility, compact. refers to compaction, stab. refers to stability, and decomp. refers to decomposition.

Category	Indicator	Link						Test	Farmer accessible kit	Criteria								Source(s)
		Habitat provision	Element cycling	Structure main.	Bio. pop. reg.	Water cycling	OM decomp.			Time	Cost/Accessibility	Reliability	Ease of use	Interpretability	Applicability	Destructiveness	Sensitivity	
Physical	Structure			X				VESS	Spade	A	G	G	G	G	A	R	R	[113, 168, 169]
				X				Bulk density	PVC ring, oven, phone, bag, knife, scale	A	R	G	R	G	G	A	R	[170–172]
				X				Workability	n/a	G	G	R	G	R	G	G	R	
	Colour			X				Colour	n/a	A	G	A	G	G	G	G	R	[161–163, 173–181]
	Stone content*			X				Stoniness	Spade	A	G	A	G	A	G	G	R	
	Depth*			X				Depth	Spade	R	G	A	G	G	G	R	R	
	Depth* (& compact.)			X				Penetration resistance	Fencing wire	G	A	A	G	A	G	G	R	[173, 182]
	Horizon* (& depth*)		X	X				Soil profiling	Spade	R	G	G	G	R	G	R	A	[183, 184]
	Texture*			X				Texture triangle test	Texture triangle, jar, tape measure, phone	G	A	A	G	A	G	A	R	[185–187]
				X				Ribbon test	Water	G	G	A	G	G	G	G	R	[188, 189]
	Infiltration			X		X		Beerkan test	PVC ring, marker, tape measure, bottle	A	A	G	G	G	G	G	R	[190–192]
	Aggregate stab.			X				Slake test	Spade, glass, wire mesh, phone	A	A	G	G	A	G	A	R	[193–195]
				X				Emerson test	Tray, water	A	G	A	G	A	G	A	R	[196–198]
				X				Slakes app	Tray, water, can, smartphone, app	A	G	G	G	G	G	A	R	[195, 199, 200]
				X				Drop shatter test	Spade, tray, plastic bag	A	G	A	G	A	G	R	R	[186]
Chemical	pH		X				X	pH test	pH strip	G	A	G	G	G	G	A	A	[201, 202]
			X				X	Vinegar test	Vinegar, baking soda	G	A	R	G	G	G	A	A	[202]
			X				X	SOCit app	Smartphone	G	G	A	G	A	G	R	A	[138, 139, 201, 203–205]
	SOC		X				X	SOCit app	Smartphone	G	G	A	G	A	G	R	A	
			X				X	Colour, feel, smell test	n/a	G	G	R	G	R	G	A	A	[206, 207]
			X				X	LOI field test	Spade, tray, scale, fan, oven, sieve, bowl, tin, tongs, stove	R	R	A	R	A	G	R	A	[79]
	Labile C		X				X	KMnO4 test	Colour chart, glass, bottle, trays, spatula, dropper	A	R	A	A	A	G	G	A	[182, 208]

(Continued.)

Table 4. (Continued.)

Category	Indicator	Link						Test	Farmer accessible kit	Criteria								Source(s)
		Habitat provision	Element cycling	Structure main.	Bio. pop. reg.	Water cycling	OM decomp.			Time	Cost/Accessibility	Reliability	Ease of use	Interpretability	Applicability	Destructiveness	Sensitivity	
Biological	Soil fauna	X			X			Simplified TSBF test	Spade, tray	R	G	A	G	A	G	R	G	[209–211]
	Roots/Compaction	X		X				Root development	Spade	A	G	A	G	G	G	A	G	[103, 212, 213]
Function	OM decomp.						X	Teabag index	Marker, teabags, scale	G	A	G	G	A	G	G	G	[214–216]
							X	Levabag test	Marker, straw, nylon bag, scale	G	R	A	G	G	G	G	G	[217, 218]
							X	Fabric decomposition	Marker, fabric material	G	A	A	G	G	R	G	G	[219–221]
	Bio. pop. reg.				X			Bait lamina strips	Marker, bait lamina stripes, organic substrate	G	R	G	A	A	G	G	G	[222, 223]
Service	Pest & disease control	X			X			Weed infestation	n/a	A	G	A	G	A	G	G	G	[161–163, 173–181]
		X			X			Crop appearance	n/a	A	G	A	G	R	G	G	G	
	Biomass production	X	X	X	X	X	X	Crop yield	n/a	A	G	A	G	A	G	G	G	
	Erosion control	X		X		X	X	Rills & Gullies	n/a	A	G	A	G	G	G	G	A	
		X		X		X	X	Gully retreat	Pegs	G	G	G	G	G	G	G	G	[224, 225]

indicators are interpreted in a consistent and uniform way [42]. Analysing soil health can be compared to a medical examination of humans, where basic indicators, such as blood pressure should be within certain bands [41, 42]. This analogy can also be applied to the concept of soil health, where specific reference values, sometimes referred to as targets, baselines, thresholds or benchmarks [227], are used to establish specific ranges.

2.4.1. Benchmarks

Due to the large global variability of extrinsic factors influencing static and dynamic soil properties, it is considered impossible to establish absolute, universal reference values for soil health indicators. Hence, the development of benchmarks applicable across different scales, e.g. local level [228], across certain soil districts [229] or soil units [230], has gained increased traction in recent years. The reference values themselves within these districts, e.g. from a soil with maximum biomass production and/or environmental performance [27, 38], are then adopted to contextualise measured values and identify differences.

Matson *et al* [227] distinguish between four quantitative methods to evaluate soil health indicators: (i) 'fixed', (ii) 'reference', (iii) 'distribution' and (iv) 'relative change'. The fixed method (i) uses published, robust values to set a fixed target value. The reference method (ii) is based on a static value which is calculated as a percentage of what would be found in a single reference situation (such as the soil under natural conditions). The distribution approach (iii) is based on a value resulting from the distribution of multiple soils in the same region. The relative change approach (iv), which is the most feasible and widely used, requires less data and works on the basis of monitoring change relative to a target value, assuming a clear target direction for the value exists. An example of this is obtaining a desired pH value between six and seven for many soils so that crop productivity and other soil functions are optimised.

2.4.2. Calibration curves

After setting target or threshold values, many researchers apply them in standard non-linear scoring functions to evaluate soil health indicators against curves based on data from the literature and expert knowledge. By doing this, qualitative and quantitative attributes can be converted into unitless scores. The curves, sometimes referred to as calibration curves, usually take the form of (a) 'more is better', (b) 'optimal range', (c) 'less is better' on a scale of 0–1 (or 0–100) [27]. For example, Lima *et al* [144] and Lenka *et al* [231] assigned the shape "more is better" (a) to indicators, such as SOC, meaning the higher the SOC the better. Furthermore, they assigned the 'optimal range' (b) to pH as near neutral values usually optimise functions in agricultural soils. On the other hand, Lima *et al* [144] and Lenka *et al* [231]

used the curve 'less is better' (c) for bulk density as lower density, within reason, indicates better soil structure. The curve shapes are dependent on the targeted soil functions and the overall societal goals for a given landscape [42]. In contrast, some researchers also used linear scoring functions, where the measured indicators are divided by the maximum or minimum value, but these estimates are usually outperformed by the suitability of the non-linear approach [5].

Interpretation mechanisms, particularly universal approaches, are often controversial due to data scarcity and differing expert judgements [27]. Methods, such as relative change assessment, allow the need for changes in agricultural management to be quickly assessed; these are often urgent in many countries of SSA. In instances where data and information on relevant indicators from comparable sites are available, fixed, reference or distribution methods may be used for a more accurate analysis [232]. In cases of insufficient data, the relative change method should be used to prevent delays. Once the tests have been interpreted individually, they are integrated into a single result for easier communication.

2.5. Integrating soil health indicators

As multiple indicators and tests are needed to assess the health of the soil adequately, simply listing a multitude of numerical values of the indicators may be too overwhelming for non-soil experts to draw a conclusion about soil management. Moreover, the results may be difficult to assess or translate into something meaningful, such as mitigation concerns and options. Once the individual soil health indicators are interpreted, the results should be integrated or aggregated to effectively inform farmers about the impact of management practices on soil health. This integration can be done either numerically or graphically.

2.5.1. Numerical representation

According to Rinot *et al* [81], the integration of multiple quantitative indicators into a numerical soil health index can be achieved using either (i) additive or (ii) weighted-additive methods. Additive approaches, which yield a number between 1 and 10, treat each indicator as an equal contributor. However, this is not truly representative of the complexity of the soil [156, 233]. The more sophisticated approach is to assign specific weights to individual indicators, depending on (a) expert judgement, (b) the indicators' importance in contributing to a specific soil function or service, or (c) based on quantitative techniques used to derive the minimum data set [142, 145, 154]. Hussain *et al* [5] compared various soil health indicator integration methods. Common numerical methods include the Comprehensive Assessment of Soil Health [234], the Soil Management Assessment Framework [148], and

others [201, 235–242]. Several visual soil examination methods can also be used to provide a soil health index, such as the Muenchberg Soil Quality Rating [106]. Despite all of these approaches, the creation of numerical soil health indices remains controversial. The aggregation of multiple indicators into a single index can result in the loss of detailed information. Given the vast array of functions provided by soils, the trade-offs and synergies between indicators would no longer be apparent. Huge threats, such as erosion, could be overlooked if the relevant indicators of aggregate stability or root biomass are combined into a single score alongside other indicators designed to assess different threats. Moreover, challenges arise when converting quantitative and qualitative, or continuous and categorical data points, as assigning absolute and exact values is subjective and can have a significant impact on the index [27].

2.5.2. Graphical representation

Instead of developing an overall soil health index, some studies use a mixed approach combining numerical and graphical representations of soil health. Here, the individual numerical values of the soil health indicators are supplemented by colour coding, e.g. according to a traffic light system in which the colours indicate low, sufficient or excessive values. Beyond the colour-coded numerical values, various types of diagrams are available for the simple visualization of individual soil health indicators, such as spider or amoeba diagrams, or parallel coordinate plots [243, 244]. These compact and concise ways allow individual indicators to be compared with reference values in a simple graphical manner without the need to summarise the indicators into indices. This might facilitate communication of the soil health assessment between diverse stakeholders without losing information, but might also be a source of confusion. The impact of management decisions on individual attributes can be readily conveyed through such diagrams, for instance, by comparing the area coverage of polygons in the spider diagram [27].

To summarise, this section covers requirements, indicators, interpretation and integration schemes. This study examines the latest, farmer-centred, soil health tests, and discusses tests that are more suitable for use by farmers. The in-depth analysis and comparison in table 4 summarises the current state of research on soil health tests suitable for farmers in SSA. While scientifically based tests tend to provide more accurate and reliable results, tests based on farmers' knowledge have advantages in terms of time and cost efficiency. Reflecting on the current challenges and limitations farmers face in SSA outlined in section 1.5 and the findings of this section, a significant research gap remains; the provision of more accurate yet accessible methods to enable farmers to assess soil health more reliably. This applies in particular to key soil health indicators, such as SOC.

3. Nature-based Solutions to improve soil health in sub-Saharan African farmlands

Once farmers have assessed the health of their soil, they must be able to make appropriate management decisions to maintain or improve soil health and biomass production. Alongside the amount of C input, agricultural management is considered the most important factor for SOC accumulation, which influences nutrient cycling, soil structure and plant growth. This is crucial for maintaining a healthy soil [67]. In order to enable sustainable development in SSA, agriculture must shift from unsustainable and intensive farming methods to accessible and sustainable NbS.

3.1. Benefits of Nature-based Solutions

As described by Seddon *et al* [59, 245], NbSs have the capacity to reduce the vulnerability of interconnected ecological and socio-economic systems to environmental shocks in three different ways; the reduction of exposure to climate hazards, the reduction of sensitivity to adverse impacts and the building of adaptive capacity [59, 246, 247]. These benefits have already been demonstrated in many case studies around the world. For example, farmers in Malawi increased yield and resilience to soil threats through intercropping maize with nitrogen-fixing trees at minimal cost, saving the equivalent of USD11.6 million annually in nitrogen-based inorganic fertiliser [248, 249]. A review of NbS in arid and semi-arid areas of Africa, including countries such as Ethiopia, Kenya and South Africa, highlighted the successful adoption of NbS in terms of socio-economic benefits, sustainable resource use, resource enhancement and conservation, as well as infrastructure sustainability and ecosystem resilience. The measures included solutions, such as water harvesting structures, afforestation and agroforestry, some of which were developed jointly with local communities, incorporating their indigenous knowledge through stakeholder engagement workshops. This involvement significantly increased the effectiveness of the NbS interventions [250]. These examples demonstrate that in the agricultural context, NbS can:

- offer low-cost solutions, especially to a vulnerable society [20, 245];
- maintain and enhance yield in drier, more variable climates [245, 251, 252];
- benefit the water supply [245, 253];
- benefit soil health, soil moisture and biodiversity [75, 254];
- reduce risks, such as flooding, erosion and water scarcity, mitigate climate change, and conserve land, water and biodiversity [254, 255];
- provide multiple other benefits [20, 44, 59, 67, 75, 248, 256–268].

When implemented appropriately, NbS can offer a ‘triple-win’ effect for people, planet and economy [20, 44, 245]. This makes NbS extremely valuable pathways for SSA farmers, both now and in the future.

3.2. Evaluation of Nature-based Solutions

Despite the variety of solutions and their numerous benefits, there is no single silver bullet that can increase farmers’ resilience to all soil threats in isolation. Each location (in different climate zones, ecosystems or regions) and each farmer (with different pressures and priorities) requires specific practices; in essence, solutions which are suitable for a particular farm in Malawi may not be applicable elsewhere in SSA, or even for their nearest neighbour [44].

In order to identify the range of suitable, sustainable agricultural practices, table 5 lists criteria that are used to evaluate the practices [269]. The criteria reflect possible motivations and key aspects for farmers to implement NbS, also drawing on the soil-related ecosystem services from table 1. These criteria can be used to determine whether a practice can be classified as a NbS and what benefits it offers. Using these metrics, potential NbS, which are listed in table 6 and compiled from various resources ([268–271]), can be assessed. In this example for Ethiopia, each NbS is compared to the baseline, corresponding to conventional agriculture as usually practised by Ethiopian smallholder farmers. The baseline refers to mixed crop-livestock agricultural systems, where livestock mainly graze on communal land, and crop rotation is practised on arable land using a small amount of chemical fertilisers, herbicides and pesticides. In the baseline case, the work on the farm is carried out without bringing in external labour. Similar to the assessment of soil health tests, it can be assumed that, although this assessment refers to suitability in the Ethiopian context, NbS are equally applicable and effective in other SSA contexts. This assumption is supported by the fact that the sources provided in table 6 is compiled from literature sources that mainly cover SSA. Hence, there are multiple practices with great potential for farmers in SSA (RQ2), particularly those with strong advantages in terms of the categories labour, finance and biomass production in table 6. However, it must be emphasised that the most suitable solution for each individual farmer (and each individual field) depends on the farmer’s circumstances, constraints and priorities. Accordingly, the choice is highly context-dependent.

If a household is short of labour, reduced tillage or residue retention are beneficial. The reduction of tillage to no or minimal tillage saves time compared to conventional agriculture and reduces erosion. Furthermore, the practice could benefit soil health, (above-ground) biodiversity conservation, and water quality and cycling. This improves the climate resilience of the farming system [275, 278, 279]. Countering this, however, is the value of tillage

for weed control and the incorporation of organic residues that may otherwise be consumed by grazing livestock or surface fauna. Similar to the reduction of tillage, the retention of residues, e.g. straw incorporation, reduces the amount of labour needed and soil erosion. However, retaining residues in the field could lead to income losses, as feed and/or fodder for animals may have to be purchased externally [283, 317].

On the other hand, if a farming household is short of finances, mixed cropping, companion cropping, intercropping, terracing or biochar made on-farm using a soil pit should be prioritised. The use of mixed cropping systems provides farmers with numerous benefits, as has already been demonstrated in Ethiopia [318, 319] and Rwanda [320], ranging from higher crop and biomass yields to reduced erosion, and pest and disease control [271, 272]. Although additional crops or legumes would need to be purchased initially, seeds could be retained for subsequent years to reduce costs. When implemented appropriately, this type of NbS offers a unique opportunity to increase the resource use efficiency of the area, as other plants, such as legumes (e.g. clover or cowpeas), can be grown alongside main crops, such as maize, without requiring additional space or extensive nutrients. However, avoidance of competition between plants for resources, such as water, light and nutrients, requires careful selection of plants and appropriate cropping design [321]. If a household is short of finances and practices agriculture on (steep) hillsides, terracing could be particularly beneficial. Terraces do not require large investments, but do involve a considerable amount of work for construction and maintenance. The terraces are built by digging soil from down to upper slopes, in some cases stabilising them with stones or branches. They are of great benefit to the soil, reducing erosion, increasing water retention, regulating the climate and improving resilience through C storage [280–282]. Biochar can improve retention of water and nutrients, so improving crop yield. If biochar is made on-farm using a no-cost soil pit from unused residues, such as rice husks, it can significantly reduce the pressure on household finances. However, this also requires considerable labour to produce the biochar so more costly options might be preferred. In cases where financial resources are scarce, smallholders could draw on climate finance mechanisms or micro-credit programmes to support the introduction of NbS, thereby reducing upfront costs while improving climate resilience [322–324].

If the key priority is to maximise production, different methods can be considered depending on the availability of resources and location. Soil-conservation methods, such as terraces, offer advantages for a farm located in hilly terrain. In these locations, agroforestry systems could be of even greater benefit, not only reducing erosion, but also diversifying sources of income [34, 273]. These systems

Table 5. Heatmap legend indicating the suitability of NbSs. The suitability of the solutions is compared with a baseline. This baseline refers to mixed crop-livestock agricultural systems, where livestock mainly graze on communal land, and crop rotation is practised on arable land involving chemical/inorganic fertilizers, herbicides and pesticides. In the baseline case, the work on the farm is carried out without external assistance. The positive (P) and negative (N) category refer to solutions which provide advantages and disadvantages over the baseline, whereas a neutral (0) rating indicates same behaviour and a variable (V) rating an unclear behaviour compared to the baseline. The description (part 1) is to be combined with only one of the categories, either P, 0, N or V, and completed with description (part 2).

Category	Criterion	Description (Part 1)	Positive (P)	Neutral (0)	Negative (N)	Variable (V)	Description (Part 2)
Labour	Establishment time	The establishment of the practice	reduces	does not affect	increases	reduces or increases	the total working time (labour).
	Operating time	The operating of the practice	reduces	does not affect	increases	reduces or increases	the total working time (labour).
Finance	Establishment cost	The establishment of the practice	reduces	does not affect	increases	reduces or increases	the total cost excluding labour to avoid double-counting.
	Operating cost	The operating of the practice	reduces	does not affect	increases	reduces or increases	the total cost excluding labour to avoid double-counting.
	Net income	The practice	increases	does not affect	reduces	increases or reduces	total net income after costs.
Production	Crop production area	The practice	increases	does not affect	reduces	increases or reduces	the total area available for cropping.
	Crop yield	The practice	increases	does not affect	reduces	increases or reduces	the total crop yield across the farmed area.
	Biomass production	The practice	increases	does not affect	reduces	increases or reduces	the total biomass production, including crops, fruit, etc.
	Pest & disease control	The practice	reduces	does not affect	increases	reduces or increases	the chance of pest and disease.
	Climate resilience	The practice	increases	does not affect	reduces	increases or reduces	the farm's resilience to extreme weather events.
Soil & water	Soil health	The practice	increases	does not affect	reduces	increases or reduces	the health of soil.
	Erosion control	The practice	reduces	does not affect	increases	reduces or increases	the chance of erosion.
	Water quality & supply	The practice	increases	does not affect	reduces	increases or reduces	the quality, cycling and supply of water.
Environment	Biodiversity conservation	The practice	increases	does not affect	reduces	increases or reduces	the above-ground biodiversity.
	Climate regulation	The practice	helps	does not affect	hinders	helps or hinders	climate regulation through e.g. carbon sequestration.

Table 6. On-farm practices potentially considered as nature-based solutions in their specific setting evaluated against criteria relevant for the uptake of nature-based solutions in Ethiopian farmlands. The suitability of the solutions is compared with a baseline. This baseline refers to mixed crop-livestock agricultural systems, where livestock mainly graze on communal land, and crop rotation is practised on arable land involving chemical/inorganic fertilizers, herbicides and pesticides. In the baseline case, the work on the farm is carried out without external assistance. The positive (P) and negative (N) categories refer to solutions which provide advantages and disadvantages over the baseline, whereas a neutral (0) rating indicates same behaviour and a variable (V) rating an unclear behaviour compared to the baseline. The practices are sorted by their impact from top to bottom, with more positive and less negative impacts being better. The corresponding legend can be found in table 5.

Potential nature-based solution	Labour		Finances			Production					Soil & water			Environment		Source(s)
	Establishment time	Operating time	Establishment cost	Operating cost	Net income	Crop production area	Crop yield	Biomass production	Pest & Disease control	Climate resilience	Soil health	Erosion control	Water quality & supply	Biodiversity conservation	Climate regulation	
Mixed crop, companion cropping, intercropping	0 ¹	N ²	0 ¹	0 ³	P ⁴	P ⁵	P ⁶	P ⁷	P ⁸	P ⁹	P ¹⁰	P ¹¹	P ¹²	P ¹³	P ¹⁴	[271, 272]
Agroforestry	N ¹⁵	P ¹⁶	N ¹⁷	P ¹⁸	P ¹⁹	N ²⁰	P ²¹	P ²²	P ²³	P ²⁴	P ²⁵	P ²⁶	P ²⁷	P ²⁸	P ²⁹	[34, 273]
Crop diversification	0 ¹	0 ³⁰	N ³¹	0 ³¹	P ³²	0 ³³	P ³⁴	P ³⁴	P ³⁵	P ³⁶	P ³⁷	P ³⁸	P ³⁹	P ⁴⁰	P ⁴¹	[274, 275]
Cover and catch crops	0 ¹	N ⁴²	0 ¹	N ⁴²	P ⁴	0 ⁴³	P ⁴⁴	P ⁴⁵	P ⁸	P ⁹	P ¹⁰	P ¹¹	P ¹²	P ¹³	P ¹⁴	[276, 277]
No (zero-tillage) or minimal (conservation) tillage ^a	0 ¹	P ⁴⁶	0 ¹	P ⁴⁶	V ⁴⁷	0 ³³	V ⁴⁸	V ⁴⁸	V ⁴⁹	P ³⁶	P ⁵⁰	P ⁵¹	P ⁵²	P ⁵³	P ⁵⁴	[275, 278, 279]
Soil-water conservation—Terraces	N ⁵⁵	N ⁵⁶	0 ⁵⁷	0 ⁵⁷	P ⁵⁸	P ⁵⁹	P ⁵⁹	P ⁵⁹	0 ⁶⁰	P ³⁶	P ⁶¹	P ¹¹	P ⁶²	P ⁶³	P ⁴¹	[280–282]
Biochar (made on farm)	0 ⁶⁴	N ⁶⁵	V ⁶⁶	0 ⁶⁷	P ⁶⁸	0 ³³	P ⁶⁹	P ⁷⁰	0 ⁷¹	P ³⁶	P ⁷²	P ⁷³	P ⁷⁴	0 ⁷⁵	P ⁴¹	[283–285]
Organic manure/compost (made on farm)	0 ⁷⁶	N ⁶⁵	0 ⁷⁶	V ⁷⁷	V ⁷⁸	0 ³³	P ⁶⁹	P ⁷⁰	0 ⁷¹	P ³⁶	P ⁷²	P ⁷³	P ⁷⁴	0 ⁷⁵	P ⁴¹	[286–288]
Residue retention	0 ¹	P ⁷⁹	0 ¹	N ⁸⁰	V ⁸¹	0 ³³	P ⁶⁹	P ⁷⁰	N ⁸²	P ³⁶	P ⁷²	P ⁷³	P ⁷⁴	0 ⁷⁵	P ⁴¹	[289, 290]
Water harvesting (e.g. ponds, tanks, gutters)	N ⁵⁵	N ⁵⁶	V ⁸³	0 ⁸³	P ⁸⁴	0 ⁸⁵	P ⁸⁴	P ⁸⁴	0 ⁶⁰	P ³⁶	P ⁸⁶	0 ⁸⁷	P ⁸⁸	P ⁸⁹	P ⁴¹	[291–293]
Biochar (bought in)	0 ¹	N ⁶⁵	0 ¹	N ⁹⁰	V ⁹¹	0 ³³	P ⁶⁹	P ⁷⁰	0 ⁷¹	P ³⁶	P ⁷²	P ⁷³	P ⁷⁴	0 ⁷⁵	P ⁴¹	[283–285]
Forestry	N ¹⁵	P ¹⁶	N ¹⁷	P ¹⁸	V ⁹²	N ²⁰	N ²⁰	V ⁹³	P ²³	P ²⁴	P ²⁵	P ²⁶	P ²⁷	P ²⁸	P ²⁹	[294, 295]
Organic manure/compost (bought in)	0 ¹	N ⁶⁵	0 ¹	N ⁹⁰	V ⁹⁴	0 ³³	P ⁶⁹	P ⁷⁰	0 ⁷¹	P ³⁶	P ⁷²	P ⁷³	P ⁷⁴	0 ⁷⁵	P ⁴¹	[286–288]
Field margins, buffer or grass strips	0 ⁹⁵	P ¹⁶	0 ⁹⁶	P ⁹⁷	N ⁹⁸	N ²⁰	N ²⁰	N ²⁰	V ⁹⁹	P ¹⁰⁰	P ²⁵	P ¹¹	P ⁶²	P ²⁸	P ¹⁰¹	[296, 297]
Soil-water conservation—Bunds, trenches	N ⁵⁵	N ⁵⁶	0 ¹⁰²	0 ¹⁰²	V ¹⁰³	N ¹⁰⁴	V ¹⁰⁵	V ¹⁰⁶	0 ⁶⁰	P ³⁶	P ⁶¹	P ¹¹	P ⁶²	P ⁶³	P ⁴¹	[280–282]
No herbicides/pesticides	0 ¹	P ¹⁰⁷	0 ¹	P ¹⁰⁸	N ¹⁰⁹	0 ³³	N ¹¹⁰	N ¹¹⁰	N ¹¹¹	N ¹¹²	N ¹¹²	N ¹¹²	N ¹¹²	P ¹¹³	V ¹¹⁴	[67, 298]
No inorganic fertilizer	0 ¹	P ¹⁰⁷	0 ¹	P ¹⁰⁸	N ¹⁰⁹	0 ³³	N ¹¹⁵	N ¹¹⁶	0 ¹¹⁷	N ¹¹²	N ¹¹²	N ¹¹²	N ¹¹²	0 ¹¹⁸	V ¹¹⁴	[67, 299]

^a Assuming increase in SOC

¹ No need for establishment

² More labour required to plant multiple crop species

³ Different costs for seeds than if only monoculture is sown, but seeds can be retained for future years

⁴ Higher biomass production and crop yield through improved soil health, water cycling and biodiversity

⁵ Use of area between main crops which cannot be used otherwise

⁶ Use of area between crops improves crop yield as well as the impact of additional crops on main crops

⁷ Use of area between crops improves total biomass

(Continued.)

Table 6. (Continued.)

8	Pest and diseases reduced through breaking up monoculture or arable crop cycles [29, 300]
9	Increased OM inputs and storage of nutrients stabilise soils and improve their resilience against weather extremes
10	Increased OM inputs and storage of nutrients allow soil structure to develop
11	Reduce run-off during heavy rainfall periods
12	Reduced evaporation and run-off through soil cover and therefore increased amount of infiltrated water to replenish groundwater
13	Greater crop variety harbours more diverse flora and fauna
14	Increased OM and nutrient input and reduced susceptibility to erosion lead to increased carbon storage
15	Requires more labour to establish than with arable crops
16	Less labour required to maintain than with arable crops
17	Trees need to be purchased
18	Forestry does not have annual fertilizer, seed or pesticide costs unlike arable farming
19	Higher biomass production
20	Arable land used up by non-arable areas or plants other than main crops
21	Shading from trees can provide a better environment for crop growth as well as improving soil structure [252, 301, 302]
22	Better crop yield and tree growth
23	Blocks of forestry break up habitats for arable pest and diseases in surrounding arable land [294, 303]
24	Deep roots and perennial growth of trees make them more climate resilient to e.g. droughts and floods, than annual crops
25	Increased OM inputs and reduced disturbance through tillage allow soil structure to develop
26	Roots stabilise soil structure [34, 303, 304]
27	Trees increase water retention, reduce evaporation, increase water cycling and supply through deeper roots (depends on tree species, e.g. eucalyptus is water-hungry) [303, 305]
28	Non-arable areas provide habitat for more species than arable monoculture
29	Trees store carbon, both in plant and soil
30	Similar labour required as for sowing usual crops
31	Seeds bought in initially, but could be retained for following years
32	Greater soil health supports growth of plants
33	No impact on land available for cropping
34	Greater diversity reduces erosion and improves soil health as well as reducing pests and diseases
35	Greater diversity disrupts pest cycles from main crops
36	Improved soil health
37	Different root architectures stabilise soil structure and support belowground biodiversity
38	Different root architectures stabilise soil structure
39	Different root architectures and improved soil health support cycling of water
40	Greater diversity provides habitat for more/different fauna
41	Improved soil health supports carbon sequestration
42	Higher labour and costs due to extra seed than if arable land were left fallow
43	Crops grown sequentially with arable crops
44	Capture of nutrients through cover/catch crops supports growth of main crops
45	Improved crop yield
46	Less equipment required than for conventional tillage
47	Reduced operating time and cost as well as improved soil health can increase profit, however diseases such as weed can reduce short-term crop yield
48	No direct impact, however potential indirect impact through greater weed infestation or positive impact on soil health
49	With good management, reduced tillage could reduce pests and diseases, but can also result in increased weed infestation and harbour more pests [306, 307]
50	Reduced soil disturbance and increased OM input through crop residue on surface allow soil structure to develop

(Continued.)

Table 6. (Continued.)

51	Reduced soil disturbance allows soil structure to develop
52	Reduced run-off improves water retention and reduces evaporation
53	Crop residues left on the surface harbour fauna
54	Reduced soil disturbance minimizes SOC to be released
55	Initial implementation requires labour
56	Maintenance of structures requires labour
57	Constructed by digging soil from down slope to upper slopes, sometimes stabilised by stones from the field
58	Potentially higher crop yield and biomass production and improved soil health benefits crop growth especially in the long-term
59	Increases area available for cropping on steep slopes
60	Minimal impact on pest and diseases
61	Reduced run-off supports soil structure and therefore soil health
62	Reduce run-off and therefore increased water retention and higher amount of infiltration
63	Improved soil structure through reduced run-off provides more habitat for flora and fauna
64	Requires minimal labour to set up biochar production, e.g. in biochar kiln or in soil pit
65	Labour required for application
66	Extra cost is required if biochar kiln is used, but not if biochar is prepared in soil pit
67	Biochar made from material/residues on the farm not needed for other purposes (e.g. animal feeds)
68	Increased crop production increases annual net income
69	Increased OM input and cation exchange capacity improve crop growth by increasing the ability to hold more nutrients [308, 309]
70	Increased crop growth improves biomass production
71	No effect on pests and diseases
72	Increased OM input supports soil health
73	Increased OM input improves soil structure [286, 310, 311]
74	Improved soil structure supports water cycling
75	No direct impact on aboveground biodiversity
76	Most composting methods do not require additional equipment
77	No costs when applying manures/composts, but decreased benefits of using manures for other purposes
78	Reduces options for other uses of manures, but improved soil health could improve crop yield especially in the long-term
79	Reduced labour as crop residues remain on arable land
80	Reduced benefits as residues cannot be used for other purposes
81	Income could be reduced through not selling or using crop residues, but improved soil health could improve crop yield especially over the long-term
82	Can result in increase in pests such as rodents [312]
83	Tanks and guttering requires initial investment, but ponds are usually constructed by simply digging to impermeable layer, with subsequent operating costs likely to be minimal
84	Collected water could be used to irrigate plants, thereby increasing crop yields
85	Measures are usually built on communal ground, therefore they do not affect the land available for individual farmers

(Continued.)

Table 6. (Continued.)

86	Improved water supply increases plant growth and so also plant inputs increasing soil OM
87	No direct impact on erosion
88	Improved water cycling and supply through increased water availability, unless water is used for other means
89	Ponds harbour diverse flora and fauna, whereas water available from guttering could improve soil health and therefore harbour more/different flora and fauna
90	Purchase increases annual costs
91	Increased soil health and crop production can outweigh additional costs and labour, but this is not always the case (e.g. depends on economy, market, material etc.) [313]
92	Income from annual crop yield is reduced, however, trees provide a long-term investment; annual income could be increased if fruit trees are grown;
93	Annual biomass production could be more or less, depends on tree species, growth profile, and other factors
94	Costs of buying manure reduces income, but improved soil health could improve crop yield especially in the long-term
95	If sown, similar labour required as for arable crops
96	If sown, similar cost of seeds as for arable crops
97	If not ploughed up, no need for additional seeding in subsequent years
98	Lower biomass production
99	Strips could harbour pests or break up habitats to reduce pests in arable monocultures
100	Perennial plants are more resilient to climate extremes than annual crops
101	Reduced cultivated areas increases carbon storage
102	Compared to other water-related measures, they require only a small amount of material available on farms such as stones or branches for initial construction and maintenance
103	Reduced due to less land available for cropping or improved due to improved water retention, crop growth and additional biomass on bunds or in trenches
104	Takes away land from cropping
105	Less land available for cropping but improved water retention is likely to benefit crop growth
106	Less land available for cropping but improved water retention could benefit crop growth and additional biomass can be produced on bunds or in trenches
107	No application required
108	No need for annual purchases
109	Less money spent on fertilizers, but reduced crop production
110	Increased pests and diseases
111	No control of pests and diseases
112	Reduced crop growth results in lower organic inputs which breaks down soil structure
113	Other species are able to proliferate
114	Reduced carbon emissions due to fertilizer production but increased loss of carbon due to lower yields
115	Less nutrients available
116	Less nutrients available for crop growth
117	No direct impact on pests and diseases
118	Controlled manually, no direct impact of fertilizers [314–316]

provide benefits in all topographies by regulating field-scale microclimate, improving soil structure and reducing water evaporation. For example, agroforestry in Ethiopia has been observed to increase water infiltration and reduce catchment runoff by up to 81% [34]. In many cases in SSA, it is reported that trees increase the crop or livestock production, thereby improving nutrition and livelihoods [34]. Regardless of the terrain, cropping regimes, such as mixed cropping, companion cropping, intercropping, diversification of crops, or cover and catch crops, can help to improve crop yield and biomass production. Diversifying crops can make better use of the land area and provide alternative sources of nutrients, which has a positive impact on yield, water, soil and income [274, 275, 325]. Diverse crops might include underutilised, native or indigenous plants, which can have significant additional benefits for biodiversity [274, 275]. Crop diversification can require the purchase of additional seeds, so increasing costs in the first year, but retention of seeds in subsequent years can avoid these extra costs over the long-term. Cover and catch crops are grown between main crops, with cover crops avoiding bare soil, so reducing erosion and loss of nutrients, while catch crops grow rapidly to take up excess nutrients and prevent them from leaching into the groundwater [67, 326–328]. These practices help to reduce erosion, increase resilience to pests and diseases and can increase crop productivity, income, livelihoods, and climate resilience and regulation [329–331]. In addition to the various cropping regimes, organic fertilisers, such as biochar, manure or compost, can also lead to higher crop yields, thanks to healthier soil resulting from the supply of organic matter and nutrients [283–288]. If they are derived from unused wastes on-farm, this can be done with no additional cost to the farm. Despite the numerous advantages and their potential, studies show that organic resources are applied in insufficient quantities in SSA [332], and across a farm they tend to be applied more in home gardens where higher value crops are grown [226]. Further expansion of this practice could be challenging. This is because organic resources, such as manure, are often important energy sources in SSA [286–288, 332, 333], and so availability may be limited.

In addition to the nature-based methods listed in table 6, which are aimed directly at individual farms, there are also other valuable community level methods from which farmers can benefit indirectly. These include, exclosures on degraded non-productive arable land, and community ponds. These methods are often implemented on communal ground to benefit the entire village. Farmers can use water from the pond or might be given permission to collect biomass from exclosure areas to feed their animals during periods of extreme drought [334, 335].

Despite the many advantages of these nature-based practices, it is important to acknowledge that it is impossible to determine in advance whether these practices can be classified as NbS according to the IUCN's standard [67]. This is because the classification depends on the final implementation of each measure in a specific context. Sensitive aspects, such as cultural integrity, depend entirely on the respective community and its circumstances. For instance, valuable resources, such as organic manures, often remain unused in Borana, southern Ethiopia, because of cultural and traditional beliefs [336]. Accordingly, a practice must be designed to ensure the rights, values and knowledge of local and indigenous peoples are fully respected [59, 245, 337, 338]. This could potentially be achieved through co-design tailored to the specific needs of communities through participatory engagement, with scientists and community members working together to identify problems, assess causes and share knowledge in order to take action. This could lead to community-driven change in land management, as has been implemented in an agropastoral community in East Africa [339]. Conversely, the availability of necessary resources (e.g. specific tree seedlings for agroforestry systems) could influence the final selection of practices. Furthermore, the applicability and suitability of solutions, such as soil and water conservation measures, depend on how much erosion is taking place. In flat areas, for example, stone bunds are less suitable than other measures to improve soil health.

In conclusion, it is essential to determine the cultural integrity, availability and applicability alongside the criteria in table 5 for each practice in each individual location [44]. Beyond that, each farmer's socio-economic conditions require tailored solutions depending on factors, such as financial resources, available livestock and the amount of biomass used as fuel [44]. Reflecting on the threatened ecosystem services and the cycle of soil degradation and social deprivation described in section 1, as well as the findings from this section, methods must be found to identify the most suitable (set of) NbS for improving the soil for a farmer in advance. To this end, modelling the impact of NbS on soil and the entire farm can be extremely valuable. Provided it is technologically feasible and smartphones are available, the models could be used to advise farmers. If smartphones or similar devices are not available, the results of modelling exercises could be communicated to farmers through agricultural advisors.

4. Predicting the impact of Nature-based Solutions on soil health and farmers in sub-Saharan African farmlands

When selecting and implementing specific practices from a wide range of NbSs in LMICs, where vital

services, such as long-term sustainable food production, are at stake, risks need to be minimised to the greatest extent possible. Therefore, it is valuable to predict the potential impact of a NbS before it is implemented. This enables the most suitable solution for a given situation to be identified in advance, therefore reducing the need for experimentation with different practices in each location. This can help farmers make more informed decisions by providing actionable recommendations on which sustainable agricultural practices to adopt. As a result, management risk is potentially reduced whilst environmental and social outcomes are improved. For NbS to be beneficial for long-term sustainability, not only environmental but also societal challenges need to be considered. Approaches that bridge benefits of mitigation strategies, with socio-economic and cultural impacts to farmers, are scant but an important step going forward [340]. This is because the demand for ecosystem services is primarily determined by socio-economic factors, and its quantification has mainly been attempted by economists [83]. However, the aim of this review is to identify ways to predict the effects of NbS on soil health and on ecosystem services and functions. Although purely socio-economic models are beyond the scope of this review, interdisciplinary models are considered here, as they could provide important information to the farmer.

4.1. Ecological modelling

The ecological impact of NbS can be obtained through a variety of models. These vary based on:

- ecosystem services and soil functions of main concern [82, 83, 341],
- soil health indicators of major concern,
- scale [82, 84, 341],
- underlying scope and formulation [82, 84, 341, 342],
- underlying principle [83, 343],
- contained environmental covariates.

The selection of suitable models depends primarily on the local context in terms of the targeted services, the modelling scale and the available soil health indicators. Therefore, models should address ecosystem services that are of central concern to farmers in SSA and are to be tackled by NbS. Models which relate to biomass production and soil erosion are of primary interest. Particularly, organic matter and element cycling models are of great relevance due to their importance in the production of biomass and erosion control [82, 84, 344].

The soil functions and ecosystem services dictate which soil health indicators are suitable for modelling. In turn, the methods available to farmers in SSA for assessing soil health indicators govern the selection of applicable models. For example, models, such

as Agro-C [345], which require total soil nitrogen levels as input data, cannot be considered as it is difficult to estimate nitrogen using only simple methods. The models benefit from parameterisation with the simple soil health indicators that are accessible to farmers, alongside more quantitative approaches. By integrating data obtained from those soil health indicators assessed across multiple farms, improved regional decisions would be feasible, and could help to guide interventions and policy changes.

Beyond that, the modelling scale needs to be carefully considered to maximise the relevance of the models for smallholder farmers. Microsite, ecosystem, plot or field scale approaches allow for detailed modelling of processes in the field. By contrast, models on farm, landscape, catchment, regional, national or global scales offer assessments of large-scale dynamics, prediction of climate change with dynamic soil feedback and the simulation of global scenarios [85]. In this context, models on a scale up to the farm level are most relevant to smallholder farmers, in particular, those capable of considering processes across the entire farm.

While the aforementioned factors determine the range of models to be considered, the formulation of the models and the availability of data are decisive in identifying which models are ultimately considered most valuable.

4.1.1. Process-based models

In soil science, the term ‘process-based models’ refers to models that are based on a conceptual understanding of soil mechanisms. Each important process is simulated, usually by one or more equations, and these equations are combined to provide an overall representation of the system. These mechanistic models serve to identify, extrapolate and predict long-term temporal and spatial changes in the soil based on internal, external, natural and anthropogenic factors [127]. For example, RothC is a process-based model, which simulates the carbon flows between different soil pools based on decomposition rates [346]. It can be used to predict SOC dynamics under different climate and land management scenarios [346]. A non-exhaustive compilation of such models can be found in table 7. Their application to represent ecological phenomena offers specific advantages in terms of transparency, physical consistency and suitability for specific cases [347]. However, due to the complexity of soil and computational efficiency, some process-based models are limited in their accuracy or generalizability, e.g. in predicting SOC changes over time on larger scale or across different climatic zones [127, 348–350]. In line with the rise in AI-based soil health assessment methods, the adoption of AI methods for modelling has gained notable momentum in recent years.

Table 7. Selection of process-based models to assess the ecological impact of nature-based solutions on soil health and soil-based ecosystem services and soil functions, in the context of soil organic carbon (SOC), crop yield, erosion, nitrogen (N) and phosphorus (P), based on Vereecken *et al* [83], Neil *et al* [341] and Aitkenhead [82]. To assess the potential for meeting the needs of Ethiopian farmers, models are selected based on their age, stage of development, specificity with respect to soil, their use cases, the spatial and temporal scale and the required soil health indicators as inputs for modelling.

Model	Use case					Source
	SOC	Crop yield	Erosion	N	P	
APSIM		X		X		[351]
CANDY	X			X		[352]
CENTURY	X			X	X	[353]
CERES		X				[354]
CN-SIM	X			X		[355]
DAISY	X	X		X		[356]
DAYCENT	X	X		X		[357]
DNDC	X	X		X		[358]
ECOSSE	X			X		[359]
EPIC	X	X	X	X	X	[360]
EUROSEM			X			[361]
ICBM	X					[362]
KINEROS2			X			[363]
LISEM			X			[364]
MONICA		X		X		[365]
N14C	X			X		[366]
RothC	X					[346]
RUSLE			X			[367]
SOMKO	X					[368]
STICS	X	X		X		[369]
SUNDIAL	X			X		[370]
WEPP			X			[371]
YASSO	X			X		[372]

4.1.2. Models based on artificial intelligence

In contrast to the soil health assessment, most of the AI-based methods for modelling the changes of soil health indicators belong to the category of learning and predicting. Algorithms in this category are purely data-driven approaches [124].

4.1.2.1. Data-based artificial intelligence algorithms

This type of model is trained solely on data, e.g. to recognise patterns or establish relationships between independent and dependent variables, and do not take into account any inherent understanding of the underlying processes. The training of such models depends heavily on the architecture of the model. The data is used to adjust selected parameters to minimise the difference between the predicted and the actual data values. If the model architecture and the selected parameters provide a good representation of the system, the model can be used for predictions on unseen data. A wide range of such methods have been employed, including statistical models (e.g. generalised additive models), widely used Machine Learning approaches (e.g. random forest), and Deep Learning models (e.g. neural networks) for a range of different uses as shown in table 8 (alongside many others not mentioned here) [373–378]. A multitude of models have been implemented to predict changes in crop yield or erosion severity in different locations across different spatial and temporal scales. Furthermore, data-driven AI models, such as random forest, support vector machines and neural networks,

and can be used to predict the dynamics of SOC, nitrogen and phosphorus. In many cases, AI methods are reported to be promising approaches for future research [379]. Some researchers have demonstrated greater prediction performance in classification and regression tasks [347, 348, 375, 380] through data-based AI methods compared to process-based methods. Others observe better computational efficiency in approximating behaviour of complex systems, enabling fast virtual experimentation [127, 375]. The simplicity, the uniformity in application across different tasks and the greater flexibility in architecture and parameterisation are key advantages of AI models, such as neural networks, over process-based methods when dealing with large amounts of data [126, 127, 381]. Machine Learning-based methods often operate within non-linear domains, establishing relationships between variables based solely on data rather than information derived from potentially inaccurate or incomplete process-based models. As a result, they avoid making unrealistic and/or linear assumptions. They can also infer missing data based on patterns in existing data and avoid the need for time-consuming expert annotations through automation and training methods, such as self-supervised learning [382].

However, AI that is purely based on data mostly overlooks the underlying soil processes (which could reveal new insights) [127]. It is worth noting that the performance of these AI algorithms strongly depends on the size and quality of the dataset [348]. Lastly, the limited interpretability of advanced, data-driven

Table 8. Selection of data-driven AI models to assess the ecological impact of nature-based solutions on soil organic carbon (SOC), crop yield, erosion, nitrogen (N) and phosphorus (P). The selection of widely used AI models in soil science, such as tree-based methods such as decision trees or random forests, through to neural networks, is narrowed down based on their ability to simulate various impacts of NbS on key aspects such as SOC.

Model	Use case				
	SOC	Crop yield	Erosion	N	P
Decision tree		X [383]			
Random forest	X [384]	X [373, 383, 385, 386]	X [380]	X [387]	X [388, 389]
Quantile regression forest	X [390]				
(Boosted) regression trees			X [391]		
Support vector machine	X [384]			X [387]	
Support vector regression			X [392]		
Multiple linear regression		X [373]	X [391]		
Logistic regression				X [387]	
Neural network	X [384, 393]	X [385]	X [391, 394]	X [387]	X [395]

AI methods, such as neural networks (sometimes referred to as a ‘black box’) poses a significant challenge [126]. However, they constitute a powerful toolbox, particularly due to their multimodal capabilities and the integration of systems, such as large language models.

4.1.2.2. Knowledge-based artificial intelligence algorithms

A rather unexplored area for AI-supported modelling of soil health indicators, but one that promises great potential for improving the interpretability of AI methods, is the domain of reasoning and decision making [83, 124, 127]. Unlike data-based approaches, which analyse statistical patterns, knowledge-based models are rooted in logic and rely heavily on tacit and explicit knowledge. This area includes conceptual schemes, such as symbolic rule inference and expert systems. Here, knowledge is captured in the form of facts and rules that are used to construct a (partially) fixed knowledge graph model [396] or neuro-symbolic AI models [397]. A common approach is the integration of diverse knowledge into graphs, including scientific as well as local and indigenous knowledge. This could be achieved through participatory and collaborative workshops between indigenous peoples and researchers documenting lived experiences or traditions, which are usually not captured in text but rather in cultural practices, such as stories, songs, dance and lore. These are then systematically translated into formalized rules [398, 399]. The graph can then be used to classify unknown data and make informed predictions according to its logic. One of the few applications of knowledge-based systems with respect to soil health is the recognition of patterns through fuzzy logic to assess soil erosion in Ethiopia [400]. This approach aims to rank the potential for erosion rather than model potential changes resulting from erosion. Another example is the application of a conceptual model to incorporate prior knowledge into the model development for predicting the soil bulk density [401]. Again, the approach is not intended to model the changes of soil

bulk density based on external or internal influences over time and/or space, but it promises to provide a more interpretable and rational form compared to classical data-mining techniques [401]. A key benefit of such approaches over classical methods is that they can typically be trained using smaller datasets, as the embedded domain knowledge restricts the search space of the optimisation task [402].

4.1.2.3. Hybrid modelling

The idea behind hybrid models is to combine different aspects, often theoretical foundations and interpretable components of physical models with the adaptability of AI methods, making them extremely relevant and versatile [403–405]. Previous research has already shown that the assimilation of biophysics and data in models is advantageous, e.g. in physics-informed neural networks or soil science-informed Machine Learning models [116, 124, 127, 347, 390, 403], differentiable programming [347], structural equation modelling [406], Bayesian belief networks [407] and geographically weighted regression kriging models [350, 408]. There are also specific applications of soil erosion models, such as the Revised Universal Soil Loss Equation (RUSLE), combined with random forests and neural networks [409], long short-term memory networks [377] or extreme gradient boosting [376]. Other models include APSIM in conjunction with random forests and multiple linear regression [373] and RothC with random forests [350, 410]. The combination of process-based and AI models could also serve as an emulator for regional crop yield changes. This allows issues, such as limited upscaling of process-based models and hallucinations of AI models, to be overcome [411, 412]. Although there is no literature currently available on the combination of process- and knowledge-based models, let alone the combination of all three (process, data and knowledge), integrating expert knowledge into modelling could be extremely valuable to future applications, especially in the expanded use of AI [124, 127, 348]. This could enable the use of previously untapped sources of information, such

as the knowledge of soil scientists, but also, more importantly, the knowledge of farmers. This could be simple information about crop productivity in different agricultural areas, which could be used to evaluate process-based models or to supervise the training of data-based AI methods. It could also yield profound insights from the experiences of predecessors regarding specific practices.

The application of ecological models, including process-based models and AI models, whether data- or knowledge-based or hybrid models, offers certain advantages in terms of their accuracy and suitability for specific processes. However, their perception is limited, as soil management practices and measures are determined by overarching conditions for the entire farm. Therefore, models must consider the different flows within the farm, such as food production, animal production, labour and income. To holistically determine entire feedback loops and interrelationships, as well as effects on the soil, models should be applied in an interdisciplinary manner.

4.2. Interdisciplinary modelling

The necessity to evaluate the impact of NbS, not only in terms of specific ecosystem services, such as crop production or soil erosion and C cycling, but also on the entire agricultural operation in a so-called '*system model*', is based on the strong interrelationships between the different parts of a system. Each part of the farming system affected by management decisions is modelled, usually using very simple approaches. The impacts of one part of the system on another are described explicitly, allowing the wider consequences of decisions to be understood. For instance, the implementation of NbS to enhance soil health through the incorporation of organic matter or modifications in tillage practices influences the use of organic material or the labour available in other parts of the farm [413]. In resource-limited households, such as many smallholder farms in SSA, this influence can lead to conflicts over resources. For example, the increased use of organic fertiliser versus the amount available for other purposes, including use as fuel for households [414]. In turn, this can result in perceived or actual costs which may prevent farmers from following recommendations to improve their soils [340, 415].

Smith *et al* [340] identified a number of different modelling frameworks linking multiple modules in order to represent cross-sectional relationships. However, most of these remain focused on crop and/or livestock production [416–419]. To close this research gap, Smith *et al* [340] developed the ORATOR model, which aims to reflect household decision-making and its consequences in other areas of smallholder farms. Modified versions of the process-based models, RothC and ECOSSE, combined with other climate, water and nutrient models, establish links between soil, plant and animal

production, and their joint impact on water and fuel consumption. The aforementioned factors are conditioned by and condition (i.e. bi-causal relationship) both on-farm and off-farm labour, which in turn lead to external expenditures and the total income of the farm [340].

In summary, in order to predict the impacts of NbS on soil health (RQ3), a range of models could be used. The analysis provided in this section highlights specific strengths and limitations of different models. While process-based models can provide accurate simulations within certain boundaries as they incorporate physical laws and soil processes, they are often limited in their scalability and generalizability due to incomplete description of processes. Although data-driven AI models can solve these issues by making use of large datasets, they often overlook underlying processes and are usually more difficult to interpret than process-based models. Knowledge-based models can offer greater interpretability and transparency and can be trained with smaller datasets by incorporating knowledge, but may demand time-consuming knowledge acquisition to ensure their reliability. System models are valuable methods as they integrate biophysical and socio-economic factors affecting the overall farm management, but they can be over-complex and inevitably require more assumptions to be made that affect the accuracy of modelling.

In view of the pressing need to enhance soil health, taking into account the constraints and socio-economic challenges for farmers outlined in section 1 and the findings of this section, a specific gap needs to be addressed promptly; the provision of accurate, scalable and interpretable models to predict the impact of NbS on soil and the entire farm.

5. Potential future research directions

Farmers in SSA urgently need readily accessible and affordable soil health assessments, coupled with actionable advice for NbS to enhance soil health and livelihoods. As demonstrated in previous sections, a vast landscape of potential tools and methods is available. Nevertheless, notable constraints and research gaps remain within individual domains, that need to be tackled to deliver tailored assistance to farmers.

5.1. Soil health assessment

To resolve the research gap identified in section 2 concerning the provision of more accurate yet accessible methods to assess soil health in SSA, this study outlines two prospective research directions that hold considerable promise to resolve the issue.

The first is the exploration of so far untapped indigenous knowledge. Since many of the soil health tests listed in table 4 are based on indigenous or farmer knowledge, there may be many other indicators that have yet to be recorded that could

prove helpful. This argument is supported by the fact that centuries ago, indigenous communities had already adopted sustainable land management practices as systems, such as observed in Terra Preta soils [420]. Similarly, there is likely to be an equally extensive body of knowledge about accurate, accessible tests to assess soil health indicators that has simply not yet been discovered or documented.

The second is the use of intelligent, holistic and technological solutions for future applications. Given the predicted increase in the availability of smartphones and similar devices, there is also a compelling rationale for exploring the potential of these tools. There are a few successful examples, such as the virtual agronomist [421], which demonstrate the potential use of smartphones in agricultural contexts to improve the soil management together with farmers in SSA. Table 4 lists solutions that could be used to assess individual soil indicators with a reasonable degree of accuracy. If smartphones become more widely available in SSA, such solutions should assess multiple indicators rather than just single factors to enable a streamlined approach and holistically assess soil health.

5.2. Use of models to inform soil health improvement through Nature-based Solutions

To successfully fill the research gap in the provision of accurate, scalable and interpretable models to predict the impacts of NbS on soils and the farm (section 4), a range of perspectives and conditions must be considered. Therefore, the ecological model of soil health should be embedded within a socio-economic framework to describe the impacts on the whole farm. This will allow soil health to be linked to social and economic factors to ensure that quantifiable benefits outweigh costs. Ideally, the ecological modelling component itself would consist of a suite of methods with different functions, including process-based models (see table 7) and AI methods based on data (see table 8) and farmers' knowledge. Such a hybrid modelling approach allows the weaknesses of individual methods to be mitigated and their strengths to be combined.

The gaps outlined in the assessment and the modelling of soil health can, of course, be addressed in isolation. However, the farmers need to gain a holistic understanding by the results of their own soil health assessment being fed directly into the models to provide actionable recommendations for improving overall soil health. This study therefore suggests that research should focus on the assessment and modelling of soil health as components of a holistic decision-support framework [421–423]. With access to smartphones, such a framework could facilitate participatory soil testing for site-specific diagnoses and long-term impact assessments, giving farmers greater autonomy. In instances where farmers lack access to smart devices, agricultural advisors could

provide assistance to provide recommendations for farmers.

6. Conclusion

Soils provide a wide range of ecosystem services that are of great importance to humanity, such as biomass production, C sequestration, erosion control and climate change mitigation. Soils around the world are at risk, especially in many countries in SSA, which are threatened by severe soil degradation due to factors including climate change, soil erosion and suboptimal land management. The health of soil needs to be improved to ensure long-term, sustainable crop production and livelihoods. Nature-based solutions offer significant benefits for soil health while also improving crop production and livelihoods.

A series of steps must be taken to translate the findings of this review into actionable policy directives and advice for farmers and other key stakeholders. This can only be achieved through long-term integrative and participatory initiatives involving farmers, agricultural advisors, non-governmental organisations and policymakers. To start this process, farmers must first be able to easily assess soil health; this could be done using the tests described in this study. When combined with modelling, this could provide farmers with actionable, farm-specific recommendations for improving soil health. This process should be facilitated through governments and researchers by providing farmers with practical field guides containing simple instructions on tests and NbSs, or offering support in using smartphones for soil assessment and improvement. In order to guarantee fair and appropriate environmental as well as social outcomes, interdisciplinary studies should be conducted to establish an evidence base for the impacts of recommended practices.

First and foremost, the focus on farmer-oriented soil health tests should serve to advance decision-making regarding sustainable management practices and raise awareness of the importance of soil health. Beyond that, the subsequent adoption of NbSs is likely to bring a number of key benefits. Primarily, there is the potential to improve soil health, which in turn increases crop yields and reduces soil erosion. These effects lead to improved food security and climate resilience, respectively. Last but not least, the introduction of NbS contributes to both climate change mitigation and adaptation. This overview aims not only to facilitate the use of sustainable and climate-resilient NbSs, but also to improve farmers' livelihoods, particularly in SSA and other LMICs worldwide.

Data availability statement

No new data were created or analysed in this study.

Acknowledgments

We would like to express our sincere gratitude to Moses Kimani for his support and insightful advice during the preparation of the manuscript. This work was supported by the UKRI AI Centre for Doctoral Training in Sustainable Understandable agri-food Systems Transformed by Artificial Intelligence (SUSTAIN) [Grant Reference: EP/Y03063X/1]. We also acknowledge the generous support of alumni and friends in establishing the University of Aberdeen's Interdisciplinary Institute, which partly enabled this research, including Dr Jane Hellman Caseley (MBChB 1956), Professor Patrick Meares (DSc 1959), Nancy Miller (MA 1942), Norman Robertson, Dr Ian Slessor (MBChB 1956) and Anne Young (MA 1957). Furthermore, we recognise the support from Global Centre on Biodiversity for Climate (GCBC) Grant Number RG2-009244. GCBC is a UK Official Development Assistance (ODA) programme that aims to support developing countries to better value, protect, restore, and sustainably manage biodiversity in ways that tackle climate change and poverty alleviation. The views expressed do not necessarily reflect the UK government's official policies.

Author contributions

Dominik Bittner  0000-0001-6583-6288
Conceptualization (lead), Formal analysis (lead), Investigation (lead), Methodology (lead), Visualization (lead), Writing – original draft (lead), Writing – review & editing (lead)

Jo Smith  0000-0001-6984-6766
Conceptualization (equal), Funding acquisition (equal), Methodology (equal), Project administration (equal), Supervision (equal), Writing – review & editing (equal)

Georgios Leontidis  0000-0001-6671-5568
Conceptualization (equal), Funding acquisition (equal), Methodology (equal), Project administration (equal), Supervision (equal), Writing – review & editing (equal)

Grant A Campbell  0009-0009-6796-1598
Writing – review & editing (equal)

Jeanne Biegel  0009-0003-9805-7522
Investigation (equal), Writing – original draft (equal), Writing – review & editing (equal)

Pete Smith
Writing – review & editing (equal)

Matthias Kuhnert
Writing – review & editing (equal)

Rastislav Skalský
Writing – review & editing (equal)

Licida M Giuliani  0000-0001-6668-7501
Writing – review & editing (equal)

Abdul Walid Salik  0000-0001-7836-2368
Writing – review & editing (equal)

Paul Hallett
Writing – review & editing (equal)

David F R P Burslem
Writing – review & editing (equal)

Getahun Yakob
Writing – review & editing (equal)

Euan Phimister
Writing – review & editing (equal)

Amare Hailelassie
Writing – review & editing (equal)

Desalegn Tegegne
Writing – review & editing (equal)

Sarem Norouzi  0000-0003-2255-8657
Writing – review & editing (equal)

Heyu Chen
Writing – review & editing (equal)

Cecile Gubry-Rangin
Writing – review & editing (equal)

Adnan Khan  0000-0002-0342-1957
Writing – review & editing (equal)

References

- [1] Keesstra S D et al 2016 The significance of soils and soil science towards realization of the United Nations sustainable development goals *Soil* **2** 111–28
- [2] Evangelista S J, Field D J, McBratney A B, Minasny B, Ng W, Padarian J, Román Dobarco M and Wadoux A M J-C 2023 A proposal for the assessment of soil security: soil functions, soil services and threats to soil *Soil Secur.* **10** 100086
- [3] Amundson R, Berhe A A, Hopmans J W, Olson C, Sztein A E and Sparks D L 2015 Soil and human security in the 21st century *Science* **348** 1261071
- [4] Braimoh A K and Vlek P L G 2008 *Land Use and Soil Resources* (<https://doi.org/10.1007/978-1-4020-6778-5>)
- [5] Hussain Z, Deng L, Wang X, Cui R and Liu G 2022 A review of farmland soil health assessment methods: current status and a novel approach *Sustain* **14** 9300
- [6] Drobnik T, Greiner L, Keller A and Grêt-Regamey A 2018 Soil quality indicators—from soil functions to ecosystem services *Ecol. Indic.* **94** 151–69
- [7] Stewart Z P, Pierzynski G M, Middendorf B J and Prasad P V V 2020 Approaches to improve soil fertility in sub-Saharan Africa *J. Exp. Bot.* **71** 632–41
- [8] Smith P et al 2024 Status of the World's soils *Annu. Rev. Environ. Resour.* **49** 73–104
- [9] Smith P et al 2016 Global change pressures on soils from land use and management *Glob. Change Biol.* **22** 1008–28
- [10] Kopittke P M, Menzies N W, Wang P, McKenna B A and Lombi E 2019 Soil and the intensification of agriculture for global food security *Environ. Int.* **132** 105078
- [11] Borrelli P et al 2017 An assessment of the global impact of 21st century land use change on soil erosion *Nat. Commun.* **8** 1–13

- [12] Wang L, Lu P, Feng S, Hamel C, Sun D, Siddique K H M and Gan G Y 2024 Strategies to improve soil health by optimizing the plant–soil–microbe–anthropogenic activity nexus *Agric., Ecosyst. Environ.* **359** 108750
- [13] Gomiero T 2016 Soil degradation, land scarcity and food security: reviewing a complex challenge *Sustainability* **18** 1–41
- [14] Solomon T B, Snyman H A and Smit G N 2007 Cattle-rangeland management practices and perceptions of pastoralists towards rangeland degradation in the Borana zone of southern Ethiopia *J. Environ. Manage.* **82** 481–94
- [15] Nkonya E, Mirzabaev A and von Braun J 2015 *Economics of Land Assessment for Improvement—a Global Degradation and Sustainable Development* (https://doi.org/10.1007/978-3-319-19168-3_3)
- [16] Leakey R 2017 Socially modified organisms in multifunctional agriculture—addressing the needs of smallholder farmers in Africa *Arch. Crop Sci.* **1** 20–29
- [17] Leakey R 2013 Addressing the causes of land degradation, food/nutritional insecurity and poverty: a new approach to agricultural intensification in the tropics and subtropics *Wake up Before it is too Late Make Agric. Truly Sustainable Now for Food Security in a Changing Climate* pp 192–8
- [18] Douxchamps S et al 2016 Linking agricultural adaptation strategies, food security and vulnerability: evidence from West Africa *Reg. Environ. Change* **16** 1305–17
- [19] Smith J et al 2019 Treatment of organic resources before soil incorporation in semi-arid regions improves resilience to El Niño, and increases crop production and economic returns *Environ. Res. Lett.* **14** 085004
- [20] Reid H, Bourne A, Muller H, Podvin K, Scorgie S and Orindi V 2018 *A Framework for Assessing the Effectiveness of Ecosystem-based Approaches to Adaptation* (Elsevier Inc.) (<https://doi.org/10.1016/B978-0-12-811891-7.00016-5>)
- [21] Tadesse A and Hailu W 2024 Causes and consequences of land degradation in Ethiopia: a review *Int. J. Sci. Qual. Anal.* **10** 10–21
- [22] Tully K, Sullivan C, Weil R and Sanchez P 2015 The state of soil degradation in sub-Saharan Africa: baselines, trajectories, and solutions *Sustain* **7** 6523–52
- [23] Carr T W, Balković J, Dodds P E, Folberth C and Skalský R 2021 The impact of water erosion on global maize and wheat productivity *Agric., Ecosyst. Environ.* **322** 107655
- [24] Solomon N, Birhane E, Tilahun M, Schauer M, Gebremedhin M A, Gebremariam F T, Gidey T and Newete S W 2024 Revitalizing Ethiopia's highland soil degradation: a comprehensive review on land degradation and effective management interventions *Discov. Sustain.* **5** 106
- [25] Guder A C and Kabeta W F 2025 Evaluation of future land use change impacts on soil erosion for holota watershed, Ethiopia pp 1–21
- [26] Moges D M, Kmoch A, Bhat H G and Uuemaa E 2020 Future soil loss in highland Ethiopia under changing climate and land use *Reg. Environ. Change* **20** 32
- [27] Bünemann E K et al 2018 Soil quality—a critical review *Soil Biol. Biochem.* **120** 105–25
- [28] Costanza R et al 1997 The value of the world's ecosystem services and natural capital *Nature* **387** 253–60
- [29] Kihara J, Bolo P, Kinyua M, Nyawira S S and Sommer R 2020 Soil health and ecosystem services: lessons from sub-Saharan Africa (SSA) *Geoderma* **370** 114342
- [30] Turner K G et al 2015 A review of methods, data, and models to assess changes in the value of ecosystem services from land degradation and restoration *Ecol. Model.* **319** 190–207
- [31] McElwee P 2021 The role of soils in learning and inspiration, physical and psychological experiences, and in supporting identities *Phil. Trans. R. Soc. B* **376** 20200184
- [32] Egoh B N, O'Farrell P J, Charef A, Josephine Gurney L, Koellner T, Nibam Abi H, Egoh M and Willemen L 2012 An African account of ecosystem service provision: use, threats and policy options for sustainable livelihoods *Ecosyst. Serv.* **2** 71–81
- [33] Tamire C, Elias E and Argaw M 2023 A systematic review of ecosystem services assessments, trends, and challenges in Ethiopia *Watershed Ecol. Environ.* **5** 38–45
- [34] Rosenstock T S et al 2019 A planetary health perspective on agroforestry in sub-Saharan Africa *One Earth* **1** 330–44
- [35] Glenk K, McVittie A and Moran D 2012 Deliverable D3. 1 : soil and soil organic carbon within an ecosystem service approach linking biophysical and economic data
- [36] Tefera M, Solomon B, Guadie A, Lakew W, Sewachen B and Shumye D 2025 Evaluating the contamination of soil, water and vegetables with heavy metals along with the estimation of transfer factor and human health risk in Gondar city, Ethiopia *Food Saf. Risk* **12** 5
- [37] Baritz R, Prokop G and Trombetti M 2021 Soil monitoring in Europe: indicators and thresholds for soil health assessments p 181
- [38] Doran J W and Parkin T B 1994 Defining and assessing soil quality *Defin. Soil Qual. Sustain. Environ.* **35** 1–21
- [39] Doran J W, Sarrantonio M, Sarrantonio M and Liebig M A 1996 Soil health and sustainability *Adv. Agron.* **56** 2–55
- [40] Hannam J A, Harris M, Deeks L, Hoskins H, Hutchison J, Withers A J, Harris J A, Way L and Rickson R J R 2025 Developing a multifunctional indicator framework for soil health *Ecol. Indic.* **175** 113515
- [41] Pankhurst C E, Doube B M and Gupta V V 1997 Biological indicators of soil health
- [42] Doran J W and Safley M 1997 Defining and assessing soil health and sustainable productivity
- [43] Sileshi G W et al 2025 A review of organic inputs to inform soil health advice for African smallholder farmers : localization matters *npj Sustain. Agric.* **3** 10
- [44] Keesstra S, Nunes J, Novara A, Finger D, Avelar D, Kalantari Z and Cerdà A 2018 The superior effect of nature based solutions in land management for enhancing ecosystem services *Sci. Total Environ.* **610–611** 997–1009
- [45] Giuliani L M, Warner E, Campbell G A, Lynch J, Smith A C and Smith P 2024 Advancing nature-based solutions through enhanced soil health monitoring in the United Kingdom *Soil Use Manag.* **40** 1–18
- [46] Lal R et al 2021 Soils and sustainable development goals of the United Nations: an international union of soil sciences perspective *Geoderma Reg.* **25** e00398
- [47] de Paul Obade V and Lal R 2014 Soil quality evaluation under different land management practices *Environ. Earth Sci.* **72** 4531–49
- [48] Smith P et al 2021 Soil-derived nature's contributions to people and their contribution to the un sustainable development goals *Phil. Trans. R. Soc. B* **376** 20200185
- [49] Efthimiou N 2025 Governance and degradation of soil in the EU. An overview of policies with a focus on soil erosion *Soil Tillage Res.* **245** 106308
- [50] Eyob G, "Ethiopia to grow 5 billion trees in the second green legacy campaign (available at: www.worldagroforestry.org/blog/2020/06/09/ethiopia-grow-5-billion-trees-second-green-legacy-campaign) (Accessed 26 June 2025)
- [51] FarmAfrica 2023 Nature-based solutions for sustainable and inclusive development
- [52] Pienaaah C K A, Mosonsieyiri M, Arku G and Luginaah I 2025 Community-led nature-based solutions for enhancing climate change preparedness and resilience in semi-arid environments *World Dev. Sustain.* **7** 100237
- [53] Ogisi O D and Begho T 2023 Adoption of climate-smart agricultural practices in sub-Saharan Africa: a review of the progress, barriers, gender differences and recommendations *Farming Syst.* **1** 100019
- [54] Curran P, Dougill A and Vincent K 2018 Policy coherence for sustainable development in sub-Saharan Africa
- [55] Collins N et al 2025 Growing resilience: unlocking the potential of nature-based solutions for climate resilience in sub-Saharan Africa

- [56] Auriacombe C and van der Walt G 2010 Fundamental policy challenges influencing sustainable development in Africa
- [57] Deal J Land degradation in Ethiopia calls for community-led soil interventions (available at: www.iwmi.org/news/land-degradation-in-ethiopia/) (Accessed 26 June 2025)
- [58] Mekuria W, Gedle A, Tesfaye Y and Phimister E 2023 Implications of changes in land use for ecosystem service values of two highly eroded watersheds in Lake Abaya Chamo sub-basin, Ethiopia *Ecosyst. Serv.* **64** 101564
- [59] Seddon N, Smith A, Smith P, Key I, Chausson A, Girardin C, House J, Srivastava S and Turner B 2021 Getting the message right on nature-based solutions to climate change *Glob. Change Biol.* **27** 1518–46
- [60] Gómez Martín E, Giordano R, Pagano A, van der Keur P and Máñez Costa M 2020 Using a system thinking approach to assess the contribution of nature based solutions to sustainable development goals *Sci. Total Environ.* **738** 139693
- [61] No P B 2006 Soil carbon policies in Ethiopia and Kenya : evolution, challenges, and opportunities introduction impacts of soil carbon policy and development deterioration through land degradation context of soil carbon no 54
- [62] Benton T G, Bieg C, Harwatt H, Pudasaini R and Wellesley L 2021 Food system impacts on biodiversity loss: three levers for food system transformation in support of nature (Energy, Environment and Resources Programme)
- [63] Gwiazdowska D, Marchwińska K and Juś K 2024 Sustainable food production and processing—sustainable agriculture and biotechnological approaches in food chain *Sustainable Food: Production and Consumption Perspectives* pp 13–28
- [64] Ramankutty N *et al* Trends in global agricultural land use : implications for environmental health and food security pp 789–815
- [65] Robinson G M 2024 Global sustainable agriculture and land management systems *Geogr. Sustain.* **5** 637–46
- [66] Oberč B P and Schnell A A 2020 Approaches to sustainable agriculture: exploring the pathways towards the future of farming (<https://doi.org/10.2305/iucn.ch.2020.07.en>)
- [67] Demozzi T, Oberč B P, López A P, Larbodièrè L, Borges M A and Schnell A A 2024 Sustainable agriculture and nature-based solutions
- [68] Muhie S H 2022 Novel approaches and practices to sustainable agriculture *J. Agric. Food Res.* **10** 100446
- [69] Qiao L 2022 *et al* Soil quality both increases crop production and improves resilience to climate change *Nat. Clim. Change* **12** 574–80
- [70] Mrunalini K *et al* 2022 Nature-based solutions in soil restoration for improving agricultural productivity *Land Degrad. Dev.* **33** 1269–89
- [71] Oberč B P, De Jong R, Demozzi T and Schnell A A *Towards a circular economy that begins and ends in nature*
- [72] Chivenge P P, Murwira H K, Giller K E, Mapfumo P and Six J 2007 Long-term impact of reduced tillage and residue management on soil carbon stabilization: implications for conservation agriculture on contrasting soils *Soil Tillage Res.* **94** 328–37
- [73] Fujisaki K *et al* 2018 Soil carbon stock changes in tropical croplands are mainly driven by carbon inputs: a synthesis *Agric. Ecosyst. Environ.* **259** 147–58
- [74] Cohen-Shacham E, Walters G, Janzen C and Maginnis S 2016 Nature-based solutions to address global societal challenges
- [75] FAO 2021 Nature-based solutions in agriculture: the case and pathway for adoption
- [76] Savilaakso S, Westerlaan P, Haddaway N, Perera O, Skidan O and Porciello J 2024 What are the programme level costs and benefits of implementing nature-based solutions for climate change mitigation and adaptation in smallholder agricultural systems? A rapid evidence assessment
- [77] Sonneveld B G J S, Merbis M D, Alfarrá A, Ünver O and Arnal M A 2018 Nature-based solutions for agricultural water management and food security
- [78] Ozturk I 2017 The dynamic relationship between agricultural sustainability and food-energy-water poverty in a panel of selected Sub-Saharan African Countries *Energy Policy* **107** 289–99
- [79] Head J S, Crockatt M E, Didarali Z, Woodward M-J and Emmett B A 2020 The role of citizen science in meeting SDG targets around soil health *Sustainability* **12** 10254
- [80] Warner E *et al* 2025 Integrated above- and below- ground ecological monitoring for nature- based solutions *Ecol. Solut. Evid.* **6** 1–14
- [81] Rinot O, Levy G J, Steinberger Y, Svoray T and Eshel G 2019 Soil health assessment: a critical review of current methodologies and a proposed new approach *Sci. Total Environ.* **648** 1484–91
- [82] Aitkenhead M 2020 Modelling nature based solutions from soil ecosystem services *Modelling Nature-based Solutions: Integrating Computational and Participatory Scenario Modelling for Environmental Management and Planning*
- [83] Vereecken H *et al* 2016 Modeling soil processes: review, key challenges, and new perspectives *Vadose Zone J.* **15** 1–57
- [84] Manzoni S and Porporato A 2009 Soil carbon and nitrogen mineralization: theory and models across scales *Soil Biol. Biochem.* **41** 1355–79
- [85] Campbell E E and Paustian K 2015 Current developments in soil organic matter modeling and the expansion of model applications: a review *Environ. Res. Lett.* **10** 123004
- [86] Kabbiri R, Dora M, Kumar V, Elepu G and Gellynck X 2018 Mobile phone adoption in agri-food sector: are farmers in sub-Saharan Africa connected? *Technol. Forecast. Soc. Change* **131** 253–61
- [87] Chen R A demand-side view of mobile internet adoption in the global South (<https://doi.org/10.1596/1813-9450-9590>)
- [88] Bahia K and Agnoletto F 2020 GSMA mobile connectivity index 2020
- [89] Porciello J, Coggins S, Otunba-Payne G, Mabaya E and Shackford S 2021 How are farmers using digital services in low- and middle-income countries? *Agriculture in the Digital Age* (available at: <https://ecommons.cornell.edu/handle/1813/103771>)
- [90] World Bank 2024 Poverty, prosperity, and planet report 2024: pathways out of the polycrisis (<https://doi.org/10.60572/01JA-MK74>)
- [91] FAO 2020 The state of food and agriculture 2020 *Overcoming Water Challenges in Agriculture*
- [92] Slayi M, Zhou L, Dzvene A R and Mpanyaro Z 2024 Drivers and consequences of land degradation on livestock productivity in Sub-Saharan Africa: a systematic literature review *Land* **13** 1402
- [93] Junior D *et al* 2024 Digitalisation in agriculture: a scoping review of technologies in practice, challenges, and opportunities for smallholder farmers in sub-saharan africa *J. Agric. Food Res.* **18** 101286
- [94] Smidt H J and Jokonya O 2022 Factors affecting digital technology adoption by small-scale farmers in agriculture value chains (AVCs) in South Africa *Inf. Technol. Dev.* **3** 558–84
- [95] Kibblewhite M G, Ritz K and Swift M J 2008 Soil health in agricultural systems *Phil. Trans. R. Soc. B* **363** 685–701
- [96] Schwilch G *et al* 2016 Operationalizing ecosystem services for the mitigation of soil threats: a proposed framework *Ecol. Indic.* **67** 586–97
- [97] Herrick J E Environmental soil management (Range and Animal Sciences and Resources Management) vol I
- [98] Herrick J E *et al* 2016 The land-potential knowledge system (landpks): mobile apps and collaboration for optimizing climate change investments *Ecosyst. Heal. Sustain.* **2** 1–7
- [99] Chiew L Y, Brodie J F, Burslem D F R P, Reynolds G, Vairappan C S and Slade E M 2024 Invertebrate-mediated

- ecosystem processes are resilient to disturbance across a land-use gradient in Borneo *Ecosystems* **27** 710–23
- [100] Feller C and Beare M H 1997 Physical control of soil organic matter dynamics in the tropics *Geoderma* **79** 69–116
- [101] Hughes H M M 2023 *et al* Towards a farmer-feasible soil health assessment that is globally applicable **345** 118582
- [102] Stott D E 2019 Recommended soil health indicators and associated laboratory procedures
- [103] Shepherd T G, Stagnari F, Pisante M and Benites J 2008 *FAO: visual soil assessment*
- [104] Hoogsteen M J J, Lantinga E A, Bakker E J, Groot J C J and Titttonell P A 2015 Estimating soil organic carbon through loss on ignition: effects of ignition conditions and structural water loss *Eur. J. Soil Sci.* **66** 320–8
- [105] Almajmaie A, Hardie M, Acuna T and Birch C 2017 Evaluation of methods for determining soil aggregate stability *Soil Tillage Res.* **167** 39–45
- [106] Mueller R *et al* 2014 The Muencheberg soil quality rating for assessing the quality of global farmland *Novel Measurement and Assessment Tools for Monitoring and Management of Land and Water Resources in Agricultural Landscapes of Central Asia*
- [107] McGarry D 2006 A methodology of a visual soil—field assessment tool to the LADA program p 50
- [108] Seiter S 1999 Soil quality card design guide: a guide to develop locally adapted conservation tools
- [109] Guo M 2021 Soil health assessment and management: recent development in science and practices *Soil Systems* **5** 61
- [110] Haney R L, Haney E B, Smith D R, Harmel R D and White M J 2018 The soil health tool—theory and initial broad-scale application *Appl. Soil Ecol.* **125** 162–8
- [111] Adeyolanu O D and Ogunkunle A O 2016 Comparison of qualitative and quantitative approaches to soil quality assessment for agricultural purposes in South-western Nigeria *Cogent Food Agric.* **2** 1149914
- [112] Chang T, Feng G, Paul V, Adeli A and Brooks J P 2022 Soil health assessment methods: progress, applications and comparison *Adv. Agron.* **172** 129–210
- [113] Ball B C, Guimarães R M L, Cloy J M, Hargreaves P R, Shepherd T G and McKenzie B M 2017 Visual soil evaluation: a summary of some applications and potential developments for agriculture *Soil Tillage Res.* **173** 114–24
- [114] Babaeian E, Sadeghi M, Jones S B, Montzka C, Verecken H and Tuller M 2019 Ground, proximal, and satellite remote sensing of soil moisture *Rev. Geophys.* **57** 530–617
- [115] Norouzi S, Sadeghi M, Liaghat A, Tuller M, Jones S B and Ebrahimian H 2021 Information depth of NIR/SWIR soil reflectance spectroscopy *Remote Sens. Environ.* **256** 112315
- [116] Minasny B, Bandai T, Ghezzehei T A, Huang Y-C, Ma Y, McBratney A B, Ng W, Norouzi S, Padarian J and Sharififar A 2024 Soil science-informed machine learning *Geoderma* **452** 117094
- [117] Woods M, Hemment D, Ajates R, Copley A, Xaver A and Konstantakopoulos G 2020 GROW Citizens' observatory: leveraging the power of citizens, open data and technology to generate engagement, and action on soil policy and soil moisture monitoring *IOP Conf. Ser.: Earth Environ. Sci.* **509** 2–4
- [118] Kovács K Z *et al* 2019 Citizen observatory based soil moisture monitoring—the GROW example *Hungarian Geogr. Bull.* **68** 119–39
- [119] McBratney A B, Mendonça Santos M L and Minasny B 2003 On digital soil mapping *Geoderma* **117** 3–52
- [120] Greiner L, Keller A, Grêt-Regamey A and Papritz A 2017 Soil function assessment: review of methods for quantifying the contributions of soils to ecosystem services *Land Use Policy* **69** 224–37
- [121] Reidy B, Simo I, Sills P and Creamer R E 2016 Pedotransfer functions for Irish soils—estimation of bulk density (pb) per horizon type *Soil* **2** 25–39
- [122] Van Looy K *et al* 2017 Pedotransfer functions in earth system science: challenges and perspectives *Rev. Geophys.* **55** 1199–256
- [123] Minasny B, Mcbratney A B and Bristow K L 1999 Comparison of different approaches to the development of pedotransfer functions for water-retention curves *Geoderma* **93** 225–53
- [124] Wadoux A 2025 Artificial intelligence in soil science *Eur. J. Soil Sci. Rev.* **76** 229–37
- [125] Minsky M 1968 *Semantic Information Processing* (The MIT Press)
- [126] Pichler M and Hartig F 2023 Machine learning and deep learning—a review for ecologists *Methods Ecol. Evol.* **14** 994–1016
- [127] Minasny B and McBratney A B 2025 Machine learning and artificial intelligence applications in soil science *Eur. J. Soil Sci. Opin.* **76** e70093
- [128] Russell S and Norvig P *Artificial Intelligence: A Modern Approach, Global Edition* 4th edn
- [129] Pournader M, Ghaderi H, Hassanzadegan A and Fahimnia B 2021 Artificial intelligence applications in supply chain management *Int. J. Prod. Econ.* **241** 108250
- [130] Doorn N 2021 Artificial intelligence in the water domain: opportunities for responsible use *Sci. Total Environ.* **755** 142561
- [131] Pace R, Schiano Di Cola V, Monti M M, Affinito A, Cuomo S, Loreto F and Ruocco M 2025 Artificial intelligence in soil microbiome analysis: a potential application in predicting and enhancing soil health—a review *Discov. Appl. Sci.* **7** 85
- [132] Wilhelm R C, van Es H M and Buckley D H 2022 Predicting measures of soil health using the microbiome and supervised machine learning *Soil Biol. Biochem.* **164** 108472
- [133] Baghche-Maryam M Z, Sheklabadi M and Ayoubi S 2025 Covariate selection approaches in spatial prediction of soil quality indices using machine learning models at the watershed scale, west of Iran *Soil Tillage Res.* **252** 106571
- [134] Wadoux A, Minasny B and McBratney A B 2020 Machine learning for digital soil mapping: applications, challenges and suggested solutions *Earth-Sci. Rev.* **210** 103359
- [135] Naeimi M, Daggupati P and Biswas A 2024 Image-based soil characterization: a review on smartphone applications *Comput. Electron. Agric.* **227** 109502
- [136] Naeimi M, Krzic M, Scott S, Daggupati P and Biswas A 2025 Optimizing image-based soil organic matter prediction: effects of illumination type and intensity *Smart Agric. Technol.* **11** 100922
- [137] Naeimi M, Porwal V, Scott S, Krzic M, Daggupati P, Vasava H, Saurette D, Biswas A, Roul A and Biswas A 2025 Deep metric learning for soil organic matter prediction: a novel similarity-based approach using smartphone-captured images *Comput. Electron. Agric.* **237** 110728
- [138] Aitkenhead M, Coull M, Gwatkin R and Donnelly D 2016 Automated soil physical parameter assessment using smartphone and digital camera imagery *J. Imaging* **2** 35
- [139] Aitkenhead M J, Poggio L, Wardell-Johnson D, Coull M C, Rivington M, Black H I J, Jacob G, Boke S and Habte M 2020 Estimating soil properties from smartphone imagery in Ethiopia *Comput. Electron. Agric.* **171** 105322
- [140] Pate S *et al* 2025 Toward sustainable compost use: prediction of organic matter via smartphone image analysis *Agronomy* **177** 1–14
- [141] Emmet-Booth J P, Forristal P D, Fenton O, Ball B C and Holden N M 2016 A review of visual soil evaluation techniques for soil structure *Soil Use Manage.* **32** 623–34
- [142] Moncada M P, Gabriels D and Cornelis W M 2014 Data-driven analysis of soil quality indicators using limited data *Geoderma* **235–236** 271–8

- [143] O'Sullivan L, Bampa F, Knights K and Creamer R E 2017 Soil protection for a sustainable future: options for a soil monitoring network for Ireland *Soil Use Manag.* **33** 346–63
- [144] Lima A C R, Brussaard L, Totola M R, Hoogmoed W B and de Goede R G M 2013 A functional evaluation of three indicator sets for assessing soil quality *Appl. Soil Ecol.* **64** 194–200
- [145] Andrews S S, Karlen D L and Mitchell J P 2002 A comparison of soil quality indexing methods for vegetable production systems in Northern California *Agric., Ecosyst. Environ.* **90** 25–45
- [146] Ritz K, Black H I J, Campbell C D, Harris J A and Wood C 2009 Selecting biological indicators for monitoring soils: a framework for balancing scientific and technical opinion to assist policy development *Ecol. Indic.* **9** 1212–21
- [147] Stone D, Ritz K, Griffiths B G, Orgiazzi A and Creamer R E 2016 Selection of biological indicators appropriate for European soil monitoring *Appl. Soil Ecol.* **97** 12–22
- [148] Andrews S S, Karlen D L and Cambardella C A 2004 The soil management assessment framework: a quantitative soil quality evaluation method *Soil Sci. Soc. Am. J.* **68** 1945–62
- [149] Trolldborg M, Aalders I, Towers W, Hallett P D, McKenzie B M, Bengough A G, Lilly A, Ball B C and Hough R L 2013 Application of Bayesian belief networks to quantify and map areas at risk to soil threats: using soil compaction as an example *Soil Tillage Res.* **132** 56–68
- [150] Aalders I, Hough R L and Towers W 2011 Risk of erosion in peat soils—an investigation using Bayesian belief networks *Soil Use Manag.* **27** 538–49
- [151] D'Hose T, Cougnon M, De Vlieghe A, Vandecasteele B, Viaene N, Cornelis W, Van Bockstaele E and Reheul D 2014 The positive relationship between soil quality and crop production: a case study on the effect of farm compost application *Appl. Soil Ecol.* **75** 189–98
- [152] de Paul Obade V and Lal R 2016 Towards a standard technique for soil quality assessment *Geoderma* **265** 96–102
- [153] Sharma K L *et al* 2015 Combined effect of tillage and organic fertilization on soil quality key indicators and indices in alluvial soils of Indo-Gangetic Plains under rainfed maize–wheat system *Arch. Agron. Soil Sci.* **61** 313–27
- [154] Mukherjee A and Lal R 2014 Comparison of soil quality index using three methods *PLoS One* **9** e105981
- [155] Shukla M K, Lal R and Ebinger M 2006 Determining soil quality indicators by factor analysis *Soil Tillage Res.* **87** 194–204
- [156] Fine A K, van Es H M, Schindelbeck R R, van Es H M and Schindelbeck R R 2017 Statistics, scoring functions, and regional analysis of a comprehensive soil health database *Soil Sci. Soc. Am. J.* **81** 589–601
- [157] Musumba M *et al* 2017 Guide for the sustainable intensification assessment framework *Feed Future* pp 1–46
- [158] Bezabih B, Lemenih M and Regassa A 2016 Farmers' perception on soil fertility status of small-scale farming system in southwestern Ethiopia *J. Soil Sci. Environ. Manage.* **7** 143–53
- [159] Karlton E, Lemenih M and Tolera M 2013 Comparing farmers' perception of soil fertility change with soil properties and crop performance in beseku, Ethiopia *Land Degrad. Dev.* **24** 228–35
- [160] Luján Soto R, Cuéllar Padilla M and de Vente J 2020 Participatory selection of soil quality indicators for monitoring the impacts of regenerative agriculture on ecosystem services *Ecosyst. Serv.* **45** 101157
- [161] Eze S, Dougill A J, Banwart S A, Sallu S M, Smith H E, Tripathi H G, Mgohele R N and Senkoro C J 2021 Farmers' indicators of soil health in the African highlands *Catena* **203** 105336
- [162] Tesfahunegn G B, Tamene L and Vlek P L G 2011 A participatory soil quality assessment in Northern Ethiopia's Mai-Negus catchment *Catena* **86** 1–13
- [163] Abera W, Assen M and Satyal P 2021 Synergy between farmers' knowledge of soil quality change and scientifically measured soil quality indicators in Wanka watershed, northwestern highlands of Ethiopia *Environ. Dev. Sustain.* **23** 1316–34
- [164] Asfaw Z and Ågren G I 2007 Farmers' local knowledge and topsoil properties of agroforestry practices in Sidama, Southern Ethiopia *Agrofor. Syst.* **71** 35–48
- [165] Kuria A W, Barrios E, Pagella T, Muthuri C W, Mukuralinda A and Sinclair F L 2018 Farmers' knowledge of soil quality indicators along a land degradation gradient in Rwanda *Geoderma Reg.* **15** e00199
- [166] Khumalo T A, Chakale M V, Asong J A, Aremu A O and Amoo S O 2025 Indigenous farming methods and crop management practices used by local farmers in Madibeng local municipality, South Africa *Sci. Rep.* **15** 1–13
- [167] Foster E, Jolliffe D M, Ibarra G L, Lakner C and Tettah-Baah S 2025 Global poverty revisited using 2021 PPPs and new data on consumption (available at: <http://hdl.handle.net/10986/43300>)
- [168] Mutuku E A, Vanlauwe B, Roobroeck D, Boeckx P and Cornelis W M 2021 Visual soil examination and evaluation in the sub-humid and semi-arid regions of Kenya *Soil Tillage Res.* **213** 105135
- [169] Franco H H S, Guimarães R M L, Tormena C A, Cherubin M R and Favilla H S 2019 Global applications of the visual evaluation of soil structure method: a systematic review and meta-analysis *Soil Tillage Res.* **190** 61–69
- [170] Brown K and Wherrett A 2025 Bulk density—measurement (available at: www.soilquality.org.au/factsheets/bulk-density-measurement) (Accessed 23 July 2025)
- [171] FAO 2023 Standard operating procedure for soil bulk density, cylinder method (<https://doi.org/10.4060/cc7568en>)
- [172] Klopp H 2023 Bulk density is an indicator of soil health (available at: <https://extension.sdstate.edu/bulk-density-indicator-soil-health>) (Accessed 23 July 2025)
- [173] Chaudhary R S *et al* 2021 Participatory soil quality assessment using low-cost tools under contrasting management practices in a vertisol *Agric. Res.* **11** 642–51
- [174] Endale T, Diels J, Tsegaye D, Kassaye A, Belayneh L and Verdoodt A 2023 Farmer-science-based soil degradation metrics guide prioritization of catchment-tailored control measures *Environ. Dev.* **45** 100783
- [175] Getahun M, Adgo E, Nigussie Z, Regassa A and Rosenlund J 2025 Exploring the integration of local and scientific soil knowledge for sustainable management in Ethiopia's highlands *Reg. Environ. Change* **25** 1–14
- [176] Jibat M, Getachew W, Getu A and Kiflew H 2019 Survey and identification of major weeds of seeds spice in Ethiopia *J. Plant Pathol. Microbiol.* **10** 1–4
- [177] Laekemariam F, Kibret K and Mamo T 2017 Farmers' soil knowledge, fertility management logic and its linkage with scientifically analyzed soil properties in southern Ethiopia *Agric. Food Secur.* **6** 1–12
- [178] Mati B M 2012 Soil and water conservation structures for smallholder agriculture. Training manual 5 *Nile Basin Initiat. Nile Equatorial Lakes Subsid. Action Program.—Reg. Agric. Trade Program. Bujumbura, Burundi* pp 3–60
- [179] Mesfin D, Assefa E and Simane B 2023 Farmers and scientific-based assessment of soil quality in the Choke agroecosystem, Upper Blue Nile Basin, Ethiopia *Environ. Sustain. Indic.* **19** 100279
- [180] NorthernRivers 2024 Soil health card
- [181] Wawire A W, Csorba Á, Kovács E, Mairura F S, Tóth J A and Michéli E 2021 Comparing farmers' soil fertility knowledge systems and scientific assessment in Upper Eastern Kenya *Geoderma* **396** 115090
- [182] Thoumazeau A *et al* 2019 Biofunctool®: a new framework to assess the impact of land management on soil quality. Part A: concept and validation of the set of indicators *Ecol. Indic.* **97** 100–10
- [183] Bockheim J G, Gennadiyev A N, Hammer R D and Tandarich J P 2005 Historical development of key concepts in pedology *Geoderma* **124** 23–36

- [184] Hartemink A E and Minasny B 2014 Towards digital soil morphometrics *Geoderma* **230–231** 305–17
- [185] Endrias M, Assen M and Legass A 2024 Impacts of land use and management methods on soil quality dynamics in central highlands of Ethiopia *Environ. Monit. Assess.* **196** 872
- [186] FAO 2020 Soil testing methods—global soil doctors programme—a farmer-to-farmer training programme (<https://doi.org/10.4060/ca2796en>)
- [187] Rajendra T C, Gomez C, Dharumarajan S and Kumar D N 2025 Assessing soil texture classification accuracy based on VNIR lab spectroscopy *Chemom. Intell. Lab. Syst.* **263** 105419
- [188] NSW Department of Primary Industries 2014 Determining soil texture using the ribboning technique *Agric. NSW Water Unit.* **1** 1–3
- [189] Norton M, Barrie D, Campbell E, Officer W and McEwen D Valuing your soils
- [190] Basset C et al 2025 Review of conceptual and empirical approaches to characterize infiltration *Vadose Zone J.* **24** e20393
- [191] Angulo-jaramillo R, Bagarello V, Di Prima S, Gosset A, Iovino M and Lassabatere L 2019 Beerkan estimation of soil transfer parameters (BEST) across soils and scales *J. Hydrol.* **576** 239–61
- [192] FAO Global soil doctors programme—field exercises (available at: www.fao.org/global-soil-partnership/soil-doctors-programme/educational-material/field-exercises/en/) (Accessed 23 July 2025)
- [193] Herrick J E, Whitford W G, De Soyza A G and Van Zee J W 2001 Field soil aggregate stability kit for soil quality and rangeland health evaluations *Catena* **44** 27–35
- [194] ThoumazEAU A et al 2018 Biofunctool®: a new framework to assess the impact of land management on soil soil quality with the Biofunctool® index *Ecol. Indic.* **97** 429–37
- [195] Rieke E L et al 2022 Evaluation of aggregate stability methods for soil health *Geoderma* **428** 116156
- [196] Patterson R 2020 Technical note : T20-1 Emerson aggregate stability test for wastewater : an interpretation for consultants and regulators pp 1–5
- [197] Department of Sustainable Natural Resources Soil survey standard test method. Emerson aggregate test *Dep. Sustain. Nat. Resour. Resour* p 4 (available at: www.environment.nsw.gov.au/resources/soils/testmethods/eat.pdf)
- [198] Raine R J and Loch S R 2003 What is a sodic soil? Identification and management options for construction sites and disturbed lands *Roads, Struct. Soils Rural Queensl.* 29–30 July Toowoomba. *Queensl. Dep. Main Roads, Brisbane* 14
- [199] Flynn K D, Bagnall D K and Morgan C L S 2020 Evaluation of SLAKES, a smartphone application for quantifying aggregate stability, in high-clay soils *Soil Sci. Soc. Am. J.* **84** 345–53
- [200] SoilHealthInstitute 2025 Slakes: a free smartphone app to measure aggregate stability
- [201] Mulat Y, Kibret K, Bedadi B and Mohammed M 2021 Soil quality evaluation under different land use types in Kersa sub-watershed, eastern Ethiopia *Environ. Syst. Res.* **10** 19
- [202] RHS Soil: understanding pH and testing soil (available at: www.rhs.org.uk/soil-composts-mulches/ph-and-testing-soil) (Accessed 23 July 2025)
- [203] Aitkenhead M SOCiT (available at: <https://ics.hutton.ac.uk/socit/>) (Accessed 23 July 2025)
- [204] Aitkenhead M J, Coull M C, Towers W, Hudson G and Black H I J 2012 Predicting soil chemical composition and other soil parameters from field observations using a neural network *Comput. Electron. Agric.* **82** 108–16
- [205] Aitkenhead M, Cameron C, Gaskin G, Choisy B, Coull M and Black H 2018 Digital RGB photography and visible-range spectroscopy for soil composition analysis *Geoderma* **313** 265–75
- [206] Quansah C, Drechsel P and Yirenkyl B B 2001 Farmers' perceptions and management of soil organic matter—a case study from West Africa *Nutr. Cycling Agroecosyst.* **64** 205–13
- [207] De Bruyn L L 2014 Characterisation of farmers' soil sense and the implications for on-farm monitoring of soil health Characterisation of farmers' soil sense and the implications for on-farm monitoring of soil health *Aust. J. Exp. Agric.* **43** 285–305
- [208] Islam R and Sundermeier A Soil quality test kit (available at: <https://ohioline.osu.edu/factsheet/SAG-4>) (Accessed 23 July 2025)
- [209] Baillie I C, Anderson J M and Ingram J S I 1990 Tropical soil biology and fertility: a handbook of methods *J. Ecol.* **78** 547
- [210] Anderson J P E and Domsch K H 1978 A physiological method for the quantitative measurement of microbial biomass in soils *Soil Biol. Biochem.* **10** 215–21
- [211] Masebo N, Birhane E, Takele S, Belay Z, Lucena J J, Perez-Sanz A and Anjulo A 2024 The diversity and abundance of soil macrofauna under different agroforestry practices in the drylands of southern Ethiopia *Agrofor. Syst.* **98** 441–59
- [212] Anbarasan S and Ramesh S 2021 The role of plant roots in nutrient uptake and soil health *Plant Sci. Arch.* **6** 5–8
- [213] ROC In field soil testing: guidance and support document pp 1–18
- [214] Mori T, Ono K and Sakai Y 2023 Testing the tea bag index as a potential indicator for assessing litter decomposition in aquatic ecosystems *Ecol. Indic.* **152** 110358
- [215] Pino V, McBratney A, O'Brien E and Ng W 2021 Boosting soil citizen-science using Tea Bag Index method towards soil security in Australia *Soil Secur.* **5** 100016
- [216] Tresch S and Fließbach A 2021 Decomposition study using tea bags: fertilCrop technical note (Tech. Guide) pp 1–4 (available at: www.teatime4science.org/%0Ahttp://orgprints.org/30717/)
- [217] Guilbaud M The Leva-bag assesses the biological activity of soils (available at: www.tema-agriculture-terroirs.fr/cultivar-grandes-cultures/cultures/le-leva-bag-evalue-lactivite-biologique-des-sols-864800.php) (Accessed 23 July 2025)
- [218] Levabag The levabag method (available at: www.levabag.com/la-methode-levabag/) (Accessed 23 July 2025)
- [219] Nachimuthu G, Hundt A, Palmer B, Schwenke G D and Knox O G G 2022 Cotton strip assay detects soil microbial degradation differences among crop rotation and tillage experiments on Vertisols *J. Microbiol. Methods* **200** 106558
- [220] SurreyWildlifeTrust Soil your undies for science (available at: www.surreywildlifetrust.org/news/soil-your-undies-science) (Accessed 23 July 2025)
- [221] USDA Soil your undies challenge (available at: www.nrcs.usda.gov/state-offices/montana/soil-your-undies-challenge) (Accessed 23 July 2025)
- [222] André A, Antunes S C, Gonçalves F and Pereira R 2009 Bait-lamina assay as a tool to assess the effects of metal contamination in the feeding activity of soil invertebrates within a uranium mine area *Environ. Pollut.* **157** 2368–77
- [223] Vorobeichik E L and Bergman I E 2023 Modification of the bait-lamina test to estimate soil macrofauna and mesofauna feeding activity *Soil Biol. Biochem.* **183** 109047
- [224] Ibrahim H, Zuni A and Abdulkadir A U 2018 Assessment of rate of gully retreat using fixed-benchmark technique in Dutsin-Ma, Katsina State, Nigeria *Sokoto J. Soc. Sci.* **8** 186–94
- [225] Mekuria W et al 2024 Gully rehabilitation in Southern Ethiopia—value and impacts for farmers 1–26
- [226] Yakob G, Habte M, Smith J U, Hallett P D, Phimister E, Rivington M, Black H and Mekuria W 2023 Changes in soil properties with long-term organic inputs due to distance from homestead and farm characteristics in Southern Ethiopian Farmlands *Geoderma Reg.* **35** e00710

- [227] Matson A *et al* 2024 Four approaches to setting soil health targets and thresholds in agricultural soils *J. Environ. Manage.* **371** 123141
- [228] Luca M *et al* 2015 Status of the World's Soil resources (available at: www.fao.org/3/a-i5199e.pdf)
- [229] Wadoux A *et al* 2024 On soil districts *Geoderma* **452** 117065
- [230] Gutierrez S, Greve M H, Møller A B, Beucher A, Arthur E, Normand S, Wollesen de Jonge L and Gomes L D C 2024 A systematic benchmarking framework for future assessments of soil health: an example from Denmark *J. Environ. Manage.* **366** 121882
- [231] Lenka N K, Meena B P, Lal R, Khandagle A, Lenka S and Shirale A O 2022 Comparing four indexing approaches to define soil quality in an intensively cropped region of Northern India *Front. Environ. Sci.* **10** 1–10
- [232] Styc Q, Minasny B, Jang H J and McBratney A 2025 Modelling soil organic carbon dynamics at the continental extent using pedogenon mapping *Eur. J. Soil Sci.* **76** 1–16
- [233] Svoray T, Michailov E, Cohen A, Rokah L and Sturm A 2012 Predicting gully initiation: comparing data mining techniques, analytical hierarchy processes and the topographic threshold *Earth Surf. Process. Landforms* **37** 607–19
- [234] Moebius-Clune B N *et al* 2016 *Comprehensive Assessment of Soil Health—The Cornell Framework Manual* 3.1 edn
- [235] Hubbs M D, Norfleet M L and Lightle D T 2003 Interpreting the soil conditioning index: a tool for measuring soil organic matter trends pp 344–8
- [236] Karlen D L and Stott D E 1994 A framework for evaluating physical and chemical indicators of soil quality *Defining Soil Quality for a Sustainable Environment. Proc. Symp. (Minneapolis, MN, January 1992)* pp 53–72
- [237] Xue R, Wang C, Liu M, Zhang D, Li K and Li N 2019 A new method for soil health assessment based on analytic hierarchy process and meta-analysis *Sci. Total Environ.* **650** 2771–7
- [238] Lilburne L, Sparling G and Schipper L 2004 Soil quality monitoring in New Zealand: development of an interpretative framework *Agric. Ecosyst. Environ.* **104** 535–44
- [239] Debeljak M *et al* 2019 A field-scale decision support system for assessment and management of soil functions *Front. Environ. Sci.* **7** 455631
- [240] Bai Z *et al* 2018 Effects of agricultural management practices on soil quality: a review of long-term experiments for Europe and China *Agric., Ecosyst. Environ.* **265** 1–7
- [241] Liebig M A, Miller M E, Varvel G E, Doran J W and Hanson J D 2004 AEPAT: software for assessing agronomic and environmental performance of management practices in long-term agroecosystem experiments *Agron. J.* **96** 109–15
- [242] Karlen D L, Veum K S, Sudduth K A, Obrycki J F and Nunes M R 2019 Soil health assessment: past accomplishments, current activities, and future opportunities *Soil Tillage Res.* **195** 104365
- [243] Rutgers M, Schouten A J, Bloem J, Van Eekeren N, De Goede R G M, Jagersop Akkerhuis G A J M, Van der Wal A, Mulder C, Brussaard L and Breure A M 2009 Biological measurements in a nationwide soil monitoring network *Eur. J. Soil Sci.* **60** 820–32
- [244] Rutgers M, van Wijnen H J, Schouten A J, Mulder C, Kuiten A M P, Brussaard L and Breure A M 2012 A method to assess ecosystem services developed from soil attributes with stakeholders and data of four arable farms *Sci. Total Environ.* **415** 39–48
- [245] Seddon N, Chausson A, Berry P, Girardin C A J, Smith A and Turner B 2020 Understanding the value and limits of nature-based solutions to climate change and other global challenges *Phil. Trans. R. Soc. B* **375** 20190120
- [246] Valenzuela R B, Yeo-chang Y and Park M S 2020 Local People's participation in mangrove restoration projects and impacts on social capital and livelihood : a case study in the philippines
- [247] Lin A 2025 The contribution of community forestry to climate change adaptive capacity in tropical dry forests : lessons from myanmar the contribution of community forestry to climate change adaptive capacity in tropical dry forests : lessons from Myanmar *Int. For. Rev.* **21** 324–40
- [248] Jones H P, Hole D G and Zavaleta E S 2012 Harnessing nature to help people adapt to climate change *Nat. Clim. Change* **2** 504–9
- [249] Critchley W, Harari N, Mollee E, Mekdaschi-Studer R and Eichenberger J 2023 Sustainable land management and climate change adaptation for small-scale land users in sub-Saharan Africa *Land* **12** 1206
- [250] Okello C, Githiora Y W, Sithole S and Owuor M A 2024 Nature-based solutions for water resource management in Africa's Arid and sem-arid lands (ASALs): a systematic review of existing interventions *Nat.-Based Solut.* **6** 100172
- [251] Torralba M, Fagerholm N, Burgess P J, Moreno G and Plieninger T 2016 Do European agroforestry systems enhance biodiversity and ecosystem services? A meta-analysis *Agric. Ecosyst. Environ.* **230** 150–61
- [252] Tschardt T *et al* 2011 Multifunctional shade-tree management in tropical agroforestry landscapes—a review *J. Appl. Ecol.* **48** 619–29
- [253] Filoso S, Bezerra M O, Weiss K C B and Palmer M A 2013 Impacts of forest restoration on water yield: a systematic review *Int. J. Water* **7** 191–205
- [254] FAO 2021 Nature-based solutions in agriculture: sustainable management and conservation of land, water, and biodiversity
- [255] Seddon N *et al* 2020 Global recognition of the importance of nature-based solutions to the impacts of climate change *Glob. Sustain.* **3** e15
- [256] Chausson A *et al* 2020 Mapping the effectiveness of nature-based solutions for climate change adaptation *Glob. Change Biol.* **26** 6134–55
- [257] Vignola R, Harvey C A, Bautista-Solis P, Avelino J, Rapidel B, Donatti C and Martinez R 2015 Ecosystem-based adaptation for smallholder farmers: definitions, opportunities and constraints *Agric. Ecosyst. Environ.* **211** 126–32
- [258] Tamburini G *et al* 2020 Agricultural diversification promotes multiple ecosystem services without compromising yield *Sci. Adv.* **6** eaba1715
- [259] Temmerman S, Meire P, Bouma T J, Herman P M J, Ysebaert T and De Vriend H J 2013 Ecosystem-based coastal defence in the face of global change *Nature* **504** 79–83
- [260] Nel J L *et al* 2014 Natural hazards in a changing world: a case for ecosystem-based management *PLoS One* **9** e95942
- [261] Guerrero-ramírez N R *et al* 2021 Global root traits (GRooT) database *Glob. Ecol. Biogeogr.* **30** 25–37
- [262] Laughlin D C and Hooper D 2014 Applying trait-based models to achieve functional targets for theory-driven ecological restoration *Ecol. Lett.* **17** 771–84
- [263] Sharma N K, Singh R J, Mandal D, Kumar A, Alam N M and Keesstra S 2017 Increasing farmer's income and reducing soil erosion using intercropping in rainfed maize-wheat rotation of Himalaya, India *Agric. Ecosyst. Environ.* **247** 43–53
- [264] Novara A, Gristina L, Guaitoli F, Santoro A and Cerdà A 2013 Managing soil nitrate with cover crops and buffer strips in Sicilian vineyards *Solid Earth* **4** 255–62
- [265] Vancampenhout K, Nyssen J, Gebremichael D, Deckers J, Poesen J, Haile M and Moeyersons J 2006 Stone bunds for soil conservation in the northern Ethiopian highlands: impacts on soil fertility and crop yield *Soil Tillage Res.* **90** 1–15
- [266] Parras-Alcántara L, Lozano-García B, Keesstra S, Cerdà A and Brevik E C 2016 Long-term effects of soil management on ecosystem services and soil loss estimation in olive grove top soils *Sci. Total Environ.* **571** 498–506

- [267] Nunes J P, Naranjo Quintanilla P, Santos J M, Serpa D, Carvalho-Santos C, Rocha J, Keizer J J and Keesstra S D 2018 Afforestation, subsequent forest fires and provision of hydrological services: a model-based analysis for a mediterranean mountainous catchment *Land Degrad. Dev.* **29** 776–88
- [268] Miralles-Wilhelm F 2023 Nature-based solutions in agricultural landscapes for reducing tradeoffs between food production, climate change, and conservation objectives *Front. Water* **5** 1247322
- [269] Palomo-campesino S, Gonz A and García-Llorente M 2018 Exploring the connections between agroecological practices and ecosystem services : a systematic literature review *Sustainability* **10** 4339
- [270] Id A C et al 2024 Harnessing nature-based solutions for economic recovery : a systematic review *PLoS Clim.* **3** e0000281
- [271] Himmelstein J, Ares A, Gallagher D and Myers J 2017 A meta-analysis of intercropping in Africa : impacts on crop yield, farmer income, and integrated pest management effects **15** 1–10
- [272] Fuchs K, Kraus D, Houska T, Kermah M, Haas E, Kiese R, Butterbach-Bahl K and Scheer C 2024 Intercropping legumes improves long term productivity and soil carbon and nitrogen stocks in sub - Saharan Africa *Glob. Biogeochem. Cycles* **38** e2024GB008159
- [273] Amare D, Wondie M, Mekuria W and Darr D 2019 Agroforestry of smallholder farmers in ethiopia: practices and benefits *Small-Scale For.* **18** 39–56
- [274] Makate C, Wang R, Makate M and Mango N 2016 Crop diversification and livelihoods of smallholder farmers in Zimbabwe : adaptive management for environmental change *Springerplus* **5** 1135
- [275] Paula P and Fern D 2022 The impact of crop diversification, tillage and fertilization type on soil total microbial, fungal and bacterial abundance: a worldwide meta-analysis of agricultural sites *Agric. Ecosyst. Environ.* **329** 107867
- [276] Khan Z R et al 2020 Achieving food security for one million sub-Saharan African poor through push—pull innovation by 2020
- [277] Midega C A O, Bruce T J A, Pickett J A, Pittchar J O, Murage A and Khan Z R 2015 Field crops research climate-adapted companion cropping increases agricultural productivity in East Africa *Field Crops Res.* **180** 118–25
- [278] Tulema B, Aune J B, Johnsen F H and Vanlauwe B 2008 The prospects of reduced tillage in tef (*Eragrostis tef* Zucca) in Gare Arera, West Shawa Zone of Oromiya, Ethiopia *Soil Tillage Res.* **99** 58–65
- [279] Githongo M W, Kiboi M N, Ngetich F K, Musafiri C M, Muriuki A and Fließbach A 2021 The effect of minimum tillage and animal manure on maize yields and soil organic carbon in sub - Saharan Africa : a meta - analysis *Environ. Challenges* **5** 100340
- [280] Gelaye Y, Shumie M and Musie S 2025 Impact of biophysical soil and water conservation practices on soil quality and bread wheat yield in Ethiopia: a bibliographic review *Discov. Sustain.* **6** 716
- [281] Hishe S, Lyimo J and Bewket W 2017 Soil and water conservation effects on soil properties in the Middle Silluh Valley, northern Ethiopia *Int. Soil Water Conserv. Res.* **5** 231–40
- [282] Amare T, Zegeye A D, Yitaferu B, Steenhuis T S, Hurni H and Zeleke G 2014 Combined effect of soil bund with biological soil and water conservation measures in the northwestern Ethiopian highlands *Ecolohydrol. Hydrobiol.* **14** 192–9
- [283] Baudron F, Jaleta M, Okitoi O and Tegegn A 2014 Conservation agriculture in African mixed crop-livestock systems: expanding the niche *Agric. Ecosyst. Environ.* **187** 171–82
- [284] Gwenzi W, Chaukura N, Mukome F N D, Machado S and Nyamasoka B 2015 Biochar production and applications in sub-Saharan Africa : opportunities, constraints, risks and uncertainties *J. Environ. Manage.* **150** 250–61
- [285] Mahmoud Y, Njenga M, Sundberg C and Roing K 2021 Soils, sinks, and smallholder farmers: examining the benefits of biochar energy transitions in Kenya *Energy Res. Soc. Sci.* **75** 102033
- [286] Bayu W, Rethman N F G and Hammes P S 2005 The role of animal manure in sustainable soil fertility management in sub-saharan africa: a review *J. Sustain. Agric.* **0046**
- [287] Washaya S and Washaya D D 2023 Benefits, concerns and prospects of using goat manure in sub-Saharan Africa *Pastoralism* **13** 28
- [288] Ndambi O A, Pelster D E and Owino J O 2019 Manure management practices and policies in sub-Saharan Africa : implications on manure quality as a fertilizer *Front. Sustain. Food Syst.* **3** 1–14
- [289] Valbuena D et al 2012 Conservation Agriculture in mixed crop–livestock systems: scoping crop residue trade-offs in sub-Saharan Africa and South Asia *Field Crops Res.* **132** 175–84
- [290] Guto S N, Pypers P, Vanlauwe B, De Ridder N and Giller K E 2012 Socio-ecological niches for minimum tillage and crop-residue retention in continuous maize cropping systems in smallholder farms of Central Kenya *Agron. J.* **104** 188–98
- [291] Brempong M B, Amankwaa-Yeboah P, Yeboah S, Owusu Danquah E, Agyeman K, Keteku A K, Addo-Danso A and Adomako J 2023 Soil and water conservation measures to adapt cropping systems to climate change facilitated water stresses in Africa *Front. Sustain. Food Syst.* **6** 1091665
- [292] Raheem A and Oyewale O 2025 Physical management strategies for enhancing soil resilience to climate change : insights From Africa **1–23**
- [293] Filho W L and Gomez J D T 2018 *Rainwater-Smart Agriculture in Arid and Semi-Arid Areas* (<https://doi.org/10.1007/978-3-319-66239-8>)
- [294] Muthee K, Duguma L, Majale C, Mucheru-Muna M, Wainaina P and Minang P 2022 A quantitative appraisal of selected agroforestry studies in the sub-Saharan Africa *Heliyon* **8** e10670
- [295] Mbow C, Van Noordwijk M, Luedeling E, Neufeldt H, Minang P A and Kowero G 2014 ScienceDirect Agroforestry solutions to address food security and climate change challenges in Africa *Curr. Opin. Environ. Sustain.* **6** 61–67
- [296] Ndakidemi B J et al 2022 Field margin plants support natural enemies in sub-Saharan Africa smallholder common bean farming systems *Plants* **12** 1–11
- [297] Haddaway N R et al 2018 The multifunctional roles of vegetated strips around and within agricultural fields *Environ. Evid.* **7** 1–43
- [298] Zhou W, Li M and Achal V 2025 A comprehensive review on environmental and human health impacts of chemical pesticide usage *Emerg. Contam.* **11** 100410
- [299] Kabato W, Hailegnaw N, Mutum L and Molnar Z 2025 Managing soil health for climate resilience and crop productivity in a changing environment *Sci. Total Environ.* **1000** 180460
- [300] Al-Musawi Z K, Vona V and Kulmány I M 2025 Utilizing different crop rotation systems for agricultural and environmental sustainability: a review *Agronomy* **15** 1–31
- [301] Wolde A and Desalegn S 2025 Impact of nitrogen-fixing trees in agroforestry on soil physicochemical properties and crop productivity in Ethiopia: a systematic review *Arch. Agron. Soil Sci.* **71** 1–16
- [302] Luedeling E et al 2016 Field-scale modeling of tree—crop interactions : challenges and development needs *AGSY* **142** 51–69
- [303] Kuyah S et al 2016 Trees in agricultural landscapes enhance provision of ecosystem services in sub-Saharan Africa *Int. J. Biodivers. Sci. Ecosyst. Serv. Manage.* **12** 255–73

- [304] Rolo V, Rivest D, Maillard É and Moreno G 2023 Agroforestry potential for adaptation to climate change: a soil-based perspective *Soil Use Manag.* **39** 1006–32
- [305] Bargués-Tobella A, Hasselquist N J, Bazié H R, Bayala J, Laudon H and Ilstedt U 2020 Trees in African drylands can promote deep soil and groundwater recharge in a future climate with more intense rainfall *Land Degrad. Dev.* **31** 81–95
- [306] Mashingaidze N, Madakadze C, Twomlow S, Nyamangara J and Hove L 2012 Crop yield and weed growth under conservation agriculture in semi-arid Zimbabwe *Soil Tillage Res.* **124** 102–10
- [307] Mhlamba B, Ercoli L, Thierfelder C and Pellegrino E 2022 Conservation agriculture practices lead to diverse weed communities and higher maize grain yield in Southern Africa *Field Crops Res.* **289** 108724
- [308] Mashamaite C V, Motsi H, Manyevere A and Poswa S B 2024 Assessing the potential of biochar as a viable alternative to synthetic fertilizers in sub-Saharan Africa smallholder farming: a review *Agronomy* **14** 1215
- [309] Motsi H, Phiri E E and Mothapo P N 2025 Manure utilisation under smallholder farmers in sub-Saharan Africa. A systematic review and meta-analysis *Nutr. Cycl. Agroecosyst.* **131** 613–34
- [310] Bolan S *et al* 2024 Biochar modulating soil biological health: a review *Sci. Total Environ.* **914** 169585
- [311] Das S, Liptzin D and Maharjan B 2023 Long-term manure application improves soil health and stabilizes carbon in continuous maize production system *Geoderma* **430** 116338
- [312] Araya T, Ochsner T E, Mkeni P N S, Hounkpatin K O L and Amelung W 2024 Challenges and constraints of conservation agriculture adoption in smallholder farms in sub-Saharan Africa: a review *Int. Soil Water Conserv. Res.* **12** 828–43
- [313] Dickinson D, Balduccio L, Buysse J, Ronsse F, van Huylenbroeck G and Prins W 2015 Cost-benefit analysis of using biochar to improve cereals agriculture *GCB Bioenergy* **7** 850–64
- [314] Ortiz A M D, Outhwaite C L, Dalin C and Newbold T 2020 Review A review of the interactions between biodiversity, agriculture, climate change, and international trade: research and policy priorities *One Earth* **4** 88–101
- [315] Mozumder P and Berrens R P 2007 Inorganic fertilizer use and biodiversity risk: an empirical investigation *Ecol. Econ.* **62** 538–43
- [316] Wan L, Liu G and Su X 2025 Organic fertilization balances biodiversity maintenance, grass production, soil storage, nutrient cycling and greenhouse gas emissions for sustainable grassland development in China: a meta-analysis *Agric. Ecosyst. Environ.* **381** 109473
- [317] Paul B K *et al* 2013 Medium-term impact of tillage and residue management on soil aggregate stability, soil carbon and crop productivity *Agric. Ecosyst. Environ.* **164** 14–22
- [318] Alemseged Y B, King G W, Coppock L R and Tohill J C 1996 A preliminary investigation of the potential for maize—legume intercropping in the semi-arid area of Sidamo region, Ethiopia. I. Maize response *S. Afr. J. Plant Soil* **13** 120–4
- [319] Tamado T, Fininsa C and Worku W 2007 Agronomic performance and productivity of common bean (*Phaseolus vulgaris* L.) varieties in double intercropping with maize (*Zea mays* L.) in Eastern Ethiopia *Asian J. Plant Sci.* **6** 749–56
- [320] Léonidas H, Toru T, Fidèle N, Watabaji M D, Gelete T B and Hubert H 2024 Monocropping vs mixed cropping systems under a changing climate: smallholder farmers' perceptions and farm profitability in Eastern Rwanda *Environ. Sustain. Indic.* **24** 100527
- [321] Mugi-Ngenga E, Bastiaans L, Anten N P R, Zingore S, Baijukya F and Giller K E 2023 The role of inter-specific competition for water in maize-legume intercropping systems in northern Tanzania *Agric. Syst.* **207** 103619
- [322] Bensadi A 2025 Subnational heterogeneity in climate-smart agriculture green finance and irrigation efficiency across seven African nations 1–15
- [323] Stewart R 2012 The impact of microfinance in sub-Saharan Africa: a systematic review of the evidence *World Dev.* **40** 2249–62
- [324] Gatto A and Sadik-zada E R 2023 Access to microfinance as a resilience policy to address sustainable development goals: a content analysis *Heliyon* **8** e10860
- [325] Estrada-carmona N 2022 Complex agricultural landscapes host more biodiversity than simple ones: a global meta-analysis 1–10
- [326] Lal R 1987 Managing the soils of sub-Saharan Africa *Science* **236** 1069–76
- [327] Atakoun A M, Tovihoudji P G, Diogo R V C, Yemadje P L, Balarabe O, Akponikpè P B I, Sekloka E, Hougni A and Tittone P 2023 Evaluation of cover crop contributions to conservation agriculture in northern Benin *Field Crops Res.* **303** 109118
- [328] Smit E H, Strauss J A and Swanepoel P A 2021 Utilisation of cover crops: implications for conservation agriculture systems in a mediterranean climate region of South Africa *Plant Soil* **462** 207–18
- [329] Wanic M, Żuk-Golaszewska K and Orzech K 2019 Catch crops and the soil environment—a review of the literature *J. Elem.* **24** 31–45
- [330] Fageria N K, Baligar V C and Bailey B A 2005 Role of cover crops in improving soil and row crop productivity *Commun. Soil Sci. Plant Anal.* **36** 2733–57
- [331] Demissie S *et al* 2024 Cover crops improve soil condition and barley yield in a subtropical highland agroecosystem *Nutr. Cycl. Agroecosyst.* **129** 257–75
- [332] Atoloye I A 2024 Revitalizing soil health for climate resilience in Africa: the need for high rates of carbon inputs *Farming Syst.* **2** 100096
- [333] Agegnehu G, Nelson P N and Bird M I 2016 Science of the total environment the effects of biochar, compost and their mixture and nitrogen fertilizer on yield and nitrogen use efficiency of barley grown on a Nitisol in the highlands of Ethiopia *Sci. Total Environ.* **569–570** 869–79
- [334] Baudron F, Mamo A, Tirfessa D and Argaw M 2015 Impact of farmland enclosure on the productivity and sustainability of a mixed crop-livestock system in the central rift valley of Ethiopia *Agric., Ecosyst. Environ.* **207** 109–18
- [335] Mathewos M and Mamo Z 2023 Enclosure effects on soil physicochemical properties and woody species diversity in the south Rift valley basin of Ethiopia *Environ. Res. Commun.* **5** 115001
- [336] Jagisso Y, Aune J and Angassa A 2019 Unlocking the agricultural potential of manure in agropastoral systems: traditional beliefs hindering its use in Southern Ethiopia *Agriculture* **9** 1–17
- [337] Seddon N 2022 Harnessing the potential of nature-based solutions for mitigating and adapting to climate change *Science* **1416** 1410–6
- [338] Mekonnen Z, Kidemu M, Abebe H, Semere M, Gebreyesus M, Worku A, Tesfaye M and Chernet A 2021 Traditional knowledge and institutions for sustainable climate change adaptation in Ethiopia *Curr. Res. Environ. Sustain.* **3** 100080
- [339] Blake W H *et al* 2021 Integrating land-water-people connectivity concepts across disciplines for co-design of soil erosion solutions *Land Degrad. Dev.* **32** 3415–30
- [340] Smith J *et al* 2020 A systems model describing the impact of organic resource use on farming households in low to middle income countries *Agric. Syst.* **184** 102895
- [341] Neil S and Ode Sang Å 2015 A review on the state of the art in scenario modelling for environmental management
- [342] Shoyama K, Kamiyama C, Morimoto J, Ooba M and Okuro T 2017 A review of modeling approaches for ecosystem services assessment in the Asian region *Ecosyst. Serv.* **26** 316–28

- [343] Kumar P et al 2021 Nature-based solutions efficiency evaluation against natural hazards: modelling methods, advantages and limitations *Sci. Total Environ.* **784** 147058
- [344] Borrelli P et al 2021 Soil erosion modelling: a global review and statistical analysis *Sci. Total Environ.* **780** 146494
- [345] Huang Y, Yu Y, Zhang W, Sun W and Liu S 2009 Agro-C : a biogeophysical model for simulating the carbon budget of agroecosystems **149** 106–29
- [346] Coleman K and Jenkinson D S 2014 RothC—a model for the turnover of carbon in soil
- [347] Shen C et al 2023 Differentiable modelling to unify machine learning and physical models for geosciences *Nat. Rev. Earth Environ.* **4** 552–67
- [348] Bernardini L G, Rosinger C, Bodner G, Keiblinger K M, Izquierdo-Verdiguier E, Spiegel H, Retzlaff C O and Holzinger A 2024 Learning vs. understanding: when does artificial intelligence outperform process-based modelling in soil organic carbon prediction? *Nat. Biotechnol.* **81** 20–31
- [349] Wadoux A, Heuvelink G B M, Lark R M, Lagacherie P, Bouma J, Mulder V L, Libohova Z, Yang L and McBratney A B 2021 Ten challenges for the future of pedometrics *Geoderma* **401** 115155
- [350] Zhang L, Heuvelink G B M, Mulder V L, Chen S, Deng X and Yang L 2024 Using process-oriented model output to enhance machine learning-based soil organic carbon prediction in space and time *Sci. Total Environ.* **922** 170778
- [351] Mccown R L, Hammer G L, Hargreaves J N G, Holzworth D P and Freebairn D M 1996 APSIM: a novel software system for model development, model testing and simulation in agricultural systems research *Agric. Syst.* **50** 255–71
- [352] Franko U, Oelschlägel B and Schenk S 1995 Simulation of temperature-, water- and nitrogen dynamics using the model CANDY *Ecol. Model.* **81** 213–22
- [353] Parton W J, Schimel D S and Cole C V 1987 Analysis of factors controlling soil organic matter levels in great plains grasslands *Soil Sci. Soc. Am. J.* **51** 1173–9
- [354] Laryea K B, Monteith J L and Smrth G D Modelling soil physical processes and crop growth in the semi-arid tropics
- [355] Petersen B M, Berntsen J, Hansen S and Jensen L S 2005 CN-SIM—a model for the turnover of soil organic matter. I. Long-term carbon and radiocarbon development *Soil Biol. Biochem.* **37** 359–74
- [356] Hansen S, Jensen H E, Nielsen N E and Svendsen H 1991 Simulation of nitrogen dynamics and biomass production in winter wheat using the Danish simulation model DAISY *Fertilizer Res.* **27** 245–59
- [357] Hartman M D 2015 Simulated interaction of carbon dynamics and nitrogen trace gas fluxes using the DAYCENT model (<https://doi.org/10.1201/9781420032635.ch8>)
- [358] Giltrap D L, Li C and Sagar S 2010 DNDC: a process-based model of greenhouse gas fluxes from agricultural soils *Agric. Ecosyst. Environ.* **136** 292–300
- [359] Res C et al 2010 Estimating changes in Scottish soil carbon stocks using ECOSSE. I. Model description and uncertainties *Clim. Res.* **45** 179–92
- [360] Williams J R, Jones C A, Kiniry J R and Spanel D A 1989 The EPIC crop growth model *Trans. Am. Soc. Agric. Eng.* **32** 497–511
- [361] Area N P, Collins F and Leuven K U 1998 The European soil erosion model (EUROSEM): a dynamic approach for predicting sediment transport from vol 544 pp 527–44
- [362] E. Society and E. Applications 2025 ICBM : the Introductory Carbon Balance Model for Exploration of Soil Carbon Balances **7** 1226–36
- [363] Al-Qurashi A, McIntyre N, Wheeler H and Unkrich C 2008 Application of the Kineros2 rainfall-runoff model to an arid catchment in Oman *J. Hydrol.* **355** 91–105
- [364] De Roo A D P J 1996 LISEM: a physically-based hydrological and soil erosion model incorporated in a GIS (IAHS-AISH Publication) pp 395–403
- [365] Nendel C, Berg M, Kersebaum K C, Mirschel W, Specka X, Wegehenkel M, Wenkel K O and Wieland R 2011 The MONICA model: testing predictability for crop growth, soil moisture and nitrogen dynamics *Ecol. Model.* **222** 1614–25
- [366] Tipping E, Rowe E C, Evans C D, Mills R T E, Emmett B A, Chaplow J S and Hall J R 2012 N14C : a plant—soil nitrogen and carbon cycling model to simulate terrestrial ecosystem responses to atmospheric nitrogen deposition *Ecol. Model.* **247** 11–26
- [367] Renard K G, Foster G R, Weesies G A, McCool D K and Yoder D C 1997 Predicting soil erosion by water: a guide to conservation planning with the Revised universal soil loss equation (RUSLE) (Accessed 9 September 2025)
- [368] Gignoux J et al 2001 Design and test of a generic cohort model of soil organic matter decomposition : the SOMKO model pp 639–60
- [369] Brisson N, Ruppin E, Goodall S and Reggia J A 1998 STICS : a generic model for the simulation of crops and their water and nitrogen balances. 1. Theory and parameterization applied to wheat and corn *J. Cereb. Blood Flow Metab.* **18** 998
- [370] Smith J U, Bradbury N J and Addiscott T M 1996 SUNDIAL: a PC-based system for simulating nitrogen dynamics in arable land *Agron. J.* **88** 38–43
- [371] Lafen J M, Elliot W J, Flanagan D C, Meyer C R and Nearing M A 1997 WEPP-predicting water erosion using a process-based model *J. Soil Water Conserv.* **52** 96–102
- [372] Liski J, Palosuo T, Peltoniemi M and Siev R 2005 Carbon and decomposition model Yasso for forest soils *Ecol. Model.* **189** 168–82
- [373] Feng P, Wang B, Liu D L, Waters C and Yu Q 2019 Incorporating machine learning with biophysical model can improve the evaluation of climate extremes impacts on wheat yield in south-eastern Australia *Agric. For. Meteorol.* **275** 100–13
- [374] Bayissa Y, Srinivasan R, Hunink J, Nyolei D, Moges S, de Andrade Costa D, Tadesse D, Melesse A and Tilahun S 2025 Evaluating the potential of Nature-based solutions to mitigate land use and climate change impacts on the hydrology of the Gefersa and Legedadi watersheds in Ethiopia *J. Hydrol. Reg. Stud.* **57** 102130
- [375] Gumiere S J, Camporese M, Botto A, Lafond J A, Paniconi C, Gallichand J and Rousseau A N 2020 Machine learning vs. physics-based modeling for real-time irrigation management *Front. Water* **2** 1–13
- [376] Samarinas N, Tsakiridis N L, Kalopesa E and Zalidis G C 2024 Soil loss estimation by water erosion in agricultural areas introducing artificial intelligence geospatial layers into the RUSLE model *Land* **13** 174
- [377] Senanayake S, Pradhan B, Alamri A and Park H 2022 A new application of deep neural network (LSTM) and RUSLE models in soil erosion prediction *Sci. Total Environ.* **845** 157220
- [378] Segura C, Neal A L, Castro-Sardiña L, Harris P, Rivero M J, Cardenas L M and Irisarri J G N 2024 Comparison of direct and indirect soil organic carbon prediction at farm field scale *J. Environ. Manage.* **365** 121573
- [379] Paudel D, Boogaard H, de Wit A, Janssen S, Osinga S, Pylaniadis C and Athanasiadis I N 2021 Machine learning for large-scale crop yield forecasting *Agric. Syst.* **187** 103016
- [380] Jimeno-Sáez P, Martínez-España R, Casalí J, Pérez-Sánchez J and Senent-Aparicio J 2022 A comparison of performance of SWAT and machine learning models for predicting sediment load in a forested Basin, Northern Spain *Catena* **212** 105953

- [381] Shahzad K, Mardare A I and Hassel A W 2024 Accelerating materials discovery: combinatorial synthesis, high-throughput characterization, and computational advances *Sci. Technol. Adv. Mater. Methods* **4** 2292486
- [382] Thessen A E 2016 Adoption of machine learning techniques in ecology and earth science *One Ecosyst.* **1** 1–38
- [383] Shams M Y, Gamel S A and Talaat F M 2024 Enhancing crop recommendation systems with explainable artificial intelligence: a study on agricultural decision-making *Neural Comput. Appl.* **36** 5695–714
- [384] Li T et al 2024 A comprehensive review of soil organic carbon estimates: integrating remote sensing and machine learning technologies *J. Soils Sediments* **24** 3556–71
- [385] Morales A and Villalobos F J 2023 Using machine learning for crop yield prediction in the past or the future *Front. Plant Sci.* **14** 1–13
- [386] Filippi P, Jones E J, Wimalathunge N S, Somarathna P D S N, Pozza L E, Ugbaje S U, Jephcott T G, Paterson S E, Whelan B M and Bishop T F A 2019 An approach to forecast grain crop yield using multi-layered, multi-farm data sets and machine learning *Precis. Agric.* **20** 1015–29
- [387] Kee S, Pan B, Qin A K and Chen D 2024 Advancing agroecosystem modelling of nitrogen losses with machine learning *Earth Crit. Zo* **1** 100006
- [388] Jiang L, Wen G, Lu J, Yang H, Jin Y, Nie X, Wang Z, Chen M, Du Y and Wang Y 2024 Science of the total environment machine learning in soil nutrient dynamics of alpine grasslands *Sci. Total Environ.* **946** 174295
- [389] Hall R L, de Santana F B, Grunsky E C, Browne M A, Lowe V, Fitzsimons M, Higgins S, Gallagher V and Daly K 2024 A machine learning approach to predicting plant available phosphorus that accounts for soil heterogeneity and regional variability *J. Soils Sediments* **24** 390–401
- [390] Heuvelink G B M et al 2021 Machine learning in space and time for modelling soil organic carbon change *Eur. J. Soil Sci.* **72** 1607–23
- [391] Sahour H, Gholami V, Vazifedan M and Saeedi S 2021 Machine learning applications for water-induced soil erosion modeling and mapping *Soil Tillage Res.* **211** 105032
- [392] Chowdhuri I et al 2020 Implementation of artificial intelligence based ensemble models for gully erosion susceptibility assessment *Remote Sens.* **12** 3620
- [393] Tao F et al 2020 Deep learning optimizes data-driven representation of soil organic carbon in earth system model over the conterminous United States *Front. Big Data* **3** 1–15
- [394] Amanollahi J, Gharibi S and Rastkhadiv A 2025 Prediction of soil erosion control ecosystem service using machine learning based on the ANN model in Asia *Environ. Sustain. Indic.* **26** 100723
- [395] Keshavarzi A and Sarmadian F 2015 RESEARCH PAPER A neural network model for estimating soil phosphorus using terrain analysis *Egypt. J. Remote Sens. Space Sci.* **18** 127–35
- [396] Peng C, Xia F, Naseriparsa M and Osborne F 2023 Knowledge graphs: opportunities and challenges *Artif. Intell. Rev.* **56** 13071–102
- [397] Bhuyan B P, Ramdane-Cherif A, Tomar R and Singh T P 2024 Neuro-symbolic artificial intelligence: a survey *Neural Comput. Appl.* **36** 12809–44
- [398] Fraisl D, See L, Fritz S, Haklay M and McCallum I 2024 Leveraging the collaborative power of AI and citizen science for sustainable development Citizen science and the SDGs *Nat. Sustain.* **8** 125–32
- [399] Lewis J E, Whaanga H and Yolgörmez C 2025 Abundant intelligences: placing AI within Indigenous knowledge frameworks *AI Soc.* **40** 2141–57
- [400] Saadatpour M, Afshar A and Afshar M H 2011 Fuzzy pattern recognition method for assessing soil erosion *Environ. Monit. Assess.* **180** 385–97
- [401] Tranter G et al 2007 Building and testing conceptual and empirical models for predicting soil bulk density *Soil Use Manag.* **23** 437–43
- [402] Norouzi S, Pesch C, Arthur E, Norgaard T and Møldrup P 2024 Physics-informed neural networks for estimating a continuous form of the soil water retention curve from basic soil properties (available at: <http://doi.org/10.22541/essoar.171926389.99753202/v1>)
- [403] Reichstein M, Camps-Valls G, Stevens B, Jung M, Denzler J, Carvalhais N and Prabhat F 2019 Deep learning and process understanding for data-driven Earth system science *Nature* **566** 195–204
- [404] Schweidtmann A M, Zhang D and von Stosch M 2024 A review and perspective on hybrid modeling methodologies *Digit. Chem. Eng.* **10** 100136
- [405] von Rueden L, Mayer S, Sifa R, Bauckhage C and Garcke J 2020 *Combining Machine Learning and Simulation to a Hybrid Modelling Approach: Current and Future Directions* vol 12080 (LNCS. Springer International Publishing)
- [406] Angelini M E, Heuvelink G B M, Kempen B and Morrás H J M 2016 Mapping the soils of an Argentine Pampas region using structural equation modelling *Geoderma* **281** 102–18
- [407] Taalab K, Corstanje R, Mayr T M, Whelan M J and Creamer R E 2015 The application of expert knowledge in Bayesian networks to predict soil bulk density at the landscape scale *Eur. J. Soil Sci.* **66** 930–41
- [408] Xie E, Zhang X, Lu F, Peng Y, Chen J and Zhao Y 2022 Integration of a process-based model into the digital soil mapping improves the space-time soil organic carbon modelling in intensively human-impacted area *Geoderma* **409** 115599
- [409] Avand M, Mohammadi M, Mirchooli F, Kavian A and Tiefenbacher J P 2023 A new approach for smart soil erosion modeling: integration of empirical and machine-learning models *Environ. Model. Assess* **28** 145–60
- [410] Zhang X, Xie E, Chen J, Peng Y, Yan G and Zhao Y 2023 Modelling the spatiotemporal dynamics of cropland soil organic carbon by integrating process-based models differing in structures with machine learning *J. Soils Sediments* **23** 2816–31
- [411] Ippolito T, Balković J, Skalsky R, Folberth C, Krisztin T and Neff J 2023 Predicting spatiotemporal soil organic carbon responses to management using EPIC-IIASA meta-models *J. Environ. Manage.* **344** 118532
- [412] Folberth C, Baklanov A, Balković J, Skalský R, Khabarov N and Obersteiner M 2019 Spatio-temporal downscaling of gridded crop model yield estimates based on machine learning *Agric. For. Meteorol.* **264** 1–15
- [413] De Barros I and Pacheco E P 2017 Integrated emergy and economic performance assessments of maize production in semiarid tropics: comparing tillage systems *J. Environ. Account. Manage.* **5** 207–29
- [414] Smith J U, Fischer A, Hallett P D, Homans H Y, Smith P, Abdul-Salam Y, Emmerling H H and Phimister E 2015 Sustainable use of organic resources for bioenergy, food and water provision in rural Sub-Saharan Africa *Renew. Sustain. Energy Rev.* **50** 903–17
- [415] Iiyama M, Mukuralinda A, Ndayambaje J D, Musana B S, Ndoli A, Mowo J G, Garrity D, Ling S and Ruganzu V 2018 Addressing the paradox—the divergence between smallholders' preference and actual adoption of agricultural innovations *Int. J. Agric. Sustain.* **16** 472–85
- [416] van Ittersum M K et al 2008 Integrated assessment of agricultural systems—A component-based framework for the European Union (SEAMLESS) *Agric. Syst.* **96** 150–65
- [417] Ma S et al 2014 The grassland model intercomparison of the MACSUR (modelling European agriculture with climate change for food security) European knowledge hub *Proc. 7th Int. Congr. Environ. Model. Softw. Bold Visions Environ. Model. iEMSs 2014* vol 4 pp 2218–25
- [418] Monteil C, Moulin C, Penot E and Ryschawy J 2022 Modelling mixed crop-livestock farms for supporting

- farmers' strategic reflections: the CLIFS approach *Comput. Electron. Agric.* **192** 106570
- [419] Kanellopoulos A *et al* 2007 FSSIM, a bio-economic farm model for simulating the response of EU farming systems to agricultural and environmental policies
- [420] Bezerra J, Turnhout E, Vasquez I M, Rittl T F, Arts B and Kuyper T W 2019 The promises of the Amazonian soil: shifts in discourses of Terra Preta and biochar *J. Environ. Policy Plan* **21** 623–35
- [421] Shepherd K D *et al* 2025 Virtual Agronomist—an AI-assisted chatbot for guiding crop management decisions of smallholder farmers in Africa pp 1–33
- [422] Smith P *et al* 2020 How to measure, report and verify soil carbon change to realize the potential of soil carbon sequestration for atmospheric greenhouse gas removal *Glob. Change Biol.* **26** 219–41
- [423] Paustian K *et al* 2019 Quantifying carbon for agricultural soil management : from the current status toward a global soil information system *Carbon Manage.* **10** 567–87